

## FINAL EXAM

A. Sign your bubble sheet on the back at the bottom in ink.

B. In pencil, write and encode in the spaces indicated:

- 1) Name (last name, first initial, middle initial)
- 2) UF ID number
- 3) **SKIP** Section number

C. Under “special codes” code in the test ID numbers 4, 1.

1	2	3	●	5	6	7	8	9	0
●	2	3	4	5	6	7	8	9	0

D. At the top right of your answer sheet, for “Test Form Code”, encode A.

● B C D E

E. 1) This test consists of 18 multiple choice questions

- 2) The time allowed is 120 minutes.
- 3) You may write on the test.
- 4) Raise your hand if you need more scratch paper or if you have a problem with your test. **DO NOT LEAVE YOUR SEAT UNLESS YOU ARE FINISHED WITH THE TEST.**

**F. KEEP YOUR BUBBLE SHEET COVERED AT ALL TIMES.**

G. When you are finished:

- 1) Before turning in your test **check carefully for transcribing errors**. Any mistakes you leave in are there to stay.
- 2) You must turn in your scantron and tearoff sheets to your discussion leader or exam proctor. Be prepared to show your picture I.D. with a legible signature.
- 3) The answers will be posted in Canvas within one day after the exam.

**Summary of Integration Formulas**

- Fundamental Theorem of Calculus

$$\int_a^b F'(x) dx = F(b) - F(a)$$

- Fundamental Theorem of Line Integrals

$$\int_C \nabla f \cdot d\vec{r} = f(\vec{r}(b)) - f(\vec{r}(a))$$

- Green's Theorem (circulation form)

$$\iint_D \text{curl } \vec{F} \cdot \hat{k} dA = \oint_C \vec{F} \cdot d\vec{r}$$

- Stokes' Theorem

$$\iint_S \text{curl } \vec{F} \cdot \hat{n} dS = \oint_C \vec{F} \cdot d\vec{r}$$

- Green's Theorem (flux form)

$$\iint_D \text{div } \vec{F} dA = \oint_C \vec{F} \cdot \hat{n} ds$$

- Divergence Theorem

$$\iiint_E \text{div } \vec{F} dV = \oiint_S \vec{F} \cdot \hat{n} dS$$

Questions 1 – 18 are worth 6 points each.

1. Let  $C$  be a simple, smooth, closed curve in the  $xy$ -plane, for how many of the following vector fields will the circulation  $\oint_C \vec{F} \cdot d\vec{r}$  be equal to zero?

i.  $\vec{F}(x, y) = \langle 3, \pi^2 \rangle$

ii.  $\vec{F}(x, y) = \langle 2y + x^2, \sin y + 2x \rangle$

iii.  $\vec{F}(x, y) = \langle x + y, x - y \rangle$

iv.  $\vec{F}(x, y) = \langle \cos y, \sin x \rangle$

(a) 0

(b) 1

(c) 2

(d) 3

(e) 4

- 
2. Let  $C$  be a line segment with initial point  $(1, 2)$  and terminal point  $(-3, 5)$ . Compute the line integral  $\int_C 2x - y \, ds$ .

(a) 0

(b)  $-\frac{55}{2}$

(c)  $-\frac{11}{2}$

(d)  $-\frac{33}{2}$

(e)  $-\frac{25}{2}$

3. If  $\vec{F}(x, y, z) = \langle xe^y, z \ln x, xyz \rangle$ , then which of the following vectors is parallel to  $\nabla \times \vec{F}$  at the point  $(1, 0, -2)$ .

(a)  $\langle 2, 0, 3 \rangle$

(b)  $\langle 2, 1, 0 \rangle$

(c)  $\langle 0, 0, 1 \rangle$

(d)  $\langle 2, -1, 0 \rangle$

(e) none of the above

---

4. Evaluate  $\int_C (e^x y + x^2) dx + (e^x + \cos(y)) dy$  where  $C$  is any smooth curve from  $(1, 0)$  to  $(0, \pi)$ .

(a)  $\frac{2}{3}$

(b)  $\pi - \frac{1}{3}$

(c) 0

(d)  $\pi$

(e)  $-\pi$

5. Let  $S$  be the part of the plane  $z = 4 + 2x + 6y$  defined over the region  $\{(x, y) | 0 \leq x \leq 2, 0 \leq y \leq 1\}$  and oriented so that the  $z$  component of the normal vector is positive. If  $\vec{F}(x, y, z) = \langle z, z, 12xy \rangle$ , what is the flux of  $\vec{F}$  across  $S$ ?

- (a) 0
- (b) -132
- (c) -144
- (d) -84
- (e) -72

- 
6. Let  $S$  be the surface defined by  $\vec{r}(u, v) = \langle 2u, -3v, v - 2u \rangle$  with domain  $\{(u, v) | 0 \leq u \leq 1, -2 \leq v \leq 0\}$ , then compute  $\iint_S xy + 2z \, dS$ .

- (a)  $14\sqrt{76}$
- (b)  $6\sqrt{76}$
- (c)  $12\sqrt{76}$
- (d)  $20\sqrt{19}$
- (e) none of the above

7. Evaluate the line integral  $\int_C \vec{F} \cdot d\vec{r}$  where  $\vec{F}(x, y) = \langle ye^{xy}, xe^{xy} \rangle$  and if the curve  $C$  is  $\vec{r}(t) = t\hat{i} - 8\hat{j}$ ,  $0 \leq t \leq 1$ .

- (a)  $e^{-8} - 2$
  - (b) 2
  - (c) 0
  - (d)  $e^{-8} - 1$
  - (e) None of the above
- 

8. Evaluate  $\int_C \sin(x) dx + z \cos(y) dy + \sin(y) dz$  where  $C$  is the ellipse  $4x^2 + 9y^2 = 36$ , oriented clockwise.

- (a) 4
- (b) 24
- (c) 0
- (d) 9
- (e) 1

9. Find the surface area of the part of the surface  $z = x^2 + y^2$  below the plane  $z = 9$ .

(a)  $\frac{\pi}{6}(3\sqrt{3} - 1)$

(b)  $\frac{\pi}{6}(3\sqrt{3} - 2\sqrt{2})$

(c)  $\frac{\pi}{6}(37^{3/2} - 1)$

(d)  $\frac{\pi}{6}(29^{3/2} - 1)$

(e)  $\frac{\pi}{6}(27^{3/2} - 1)$

---

10. Compute the tangent plane to the surface parametrized by  $\vec{r} = u\hat{i} + uv\hat{j} + (u + v)\hat{k}$  at the point  $(1,2,3)$ .

(a)  $3x + 2y + z = 10$

(b)  $x - y + z = 2$

(c)  $x + 2y + 3z = 14$

(d)  $\langle x, y, z \rangle = \langle 1 + u, 2 + uv, 3 + u + v \rangle$

(e)  $\frac{x-1}{1} = \frac{y-2}{2} = \frac{z-3}{3}$

11. Let  $\vec{F}(x, y, z) = \langle 2y^3, 1, e^z \rangle$ . Use Stokes' Theorem to set up an integral to find the circulation of  $\vec{F}$  along  $C$ , where  $C$  is the curve of intersection of the plane  $y + 2z = 3$  and the cylinder  $x^2 + y^2 = 4$ . Orient  $C$  to be counterclockwise when viewed from above.

(a)  $\int_0^{2\pi} \int_0^2 -12r^3 \sin^2 \theta \, dr \, d\theta$

(b)  $\int_0^{2\pi} \int_0^2 12r^2 \cos^2 \theta \, dr \, d\theta$

(c)  $\int_0^{2\pi} \int_0^2 12r^3 \cos^2 \theta \, dr \, d\theta$

(d)  $\int_0^{2\pi} \int_0^2 -6r^2 \sin^2 \theta \, dr \, d\theta$

(e)  $\int_0^{2\pi} \int_0^2 -6r^3 \sin^2 \theta \, dr \, d\theta$

12. Suppose  $\vec{F}(x, y, z) = \langle xy^2, yz^2, zx^2 \rangle$  and let  $S$  be the union of the upper hemisphere of radius 2 centered at the origin with the disc of radius 2 in the  $xy$ -plane centered at the origin such that  $S$  is positively oriented. Evaluate  $\iint_S \vec{F}(x, y, z) \cdot d\vec{S}$ .

(a) 0

(b)  $\frac{8\pi^2}{3}$

(c)  $\frac{16\pi^2}{3}$

(d)  $\frac{64\pi}{5}$

(e)  $\frac{128\pi}{5}$








13. Find the work done by the force field  $\vec{F}(x, y) = \langle x, y + 2 \rangle$  in moving an object along an arch of the cycloid  $r(t) = \langle t - \sin t, 1 - \cos t \rangle$ ,  $0 \leq t \leq 2\pi$ .

- (a)  $\pi^2$
  - (b)  $2\pi^2$
  - (c)  $3\pi^2$
  - (d)  $4\pi^2$
  - (e) None of the Above
- 

14. Let  $f(x, y, z)$  be a potential function of the vector field  $\vec{F} = \langle 2x, z \cos(yz), y \cos(yz) \rangle$ . If  $f(0, 0, 0) = -1$ , find  $f\left(2, 1, \frac{\pi}{2}\right)$

- (a) 1
- (b) 2
- (c) 3
- (d) 4
- (e) 5

15. Sketch the gradient field of  $f(x, y) = x - y^2$ .



16. Let  $\vec{F} = \langle yz + 1, xz + 1, xy + 1 \rangle$ . Which of the following statements must be correct?

P.  $\vec{F}$  is conservative.

Q.  $\int_{C_1} \vec{F} \cdot d\vec{r} = \int_{C_2} \vec{F} \cdot d\vec{r}$  for any two smooth curves  $C_1$  and  $C_2$ .

R. The circulation of  $\vec{F}$  along a smooth curve is zero.

- a. P and Q only
- b. P and R only
- c. Q and R only
- d. P, Q, and R
- e. P only

17. Let  $\vec{F} = \langle z - 2y, z + 2x, e^{-xy} \rangle$  and  $S$  be the part of the paraboloid  $z = 9 - x^2 - y^2$  with  $z \geq 0$  and oriented so that  $z$  component of the normal vector is positive. Calculate

$$\iint_S \nabla \times \vec{F} \cdot \vec{n} \, dS =$$

- a.  $9\pi$
  - b.  $18\pi$
  - c.  $36\pi$
  - d.  $48\pi$
  - e. 0
- 

18. You attend the University of

- (a) Florida
- (b) Missouri
- (c) Arkansas
- (d) Kansas