

A. Sign your bubble sheet on the back at the bottom in ink.

B. In pencil, write and encode in the spaces indicated:

- 1) Name (last name, first initial, middle initial)
- 2) UF ID number
- 3) Section number

C. Under “special codes” code in the test ID numbers 4, 1.

1	2	3	●	5	6	7	8	9	0
●	2	3	4	5	6	7	8	9	0

D. At the top right of your answer sheet, for “Test Form Code”, encode A.

● B C D E

E. 1) This test consists of 24 multiple choice questions. The test is counted out of 100 points, and there are 12 bonus points available.

2) The time allowed is 120 minutes.

3) Raise your hand if you need more scratch paper or if you have a problem with your test. **DO NOT LEAVE YOUR SEAT UNLESS YOU ARE FINISHED WITH THE TEST.**

F. KEEP YOUR BUBBLE SHEET COVERED AT ALL TIMES.

G. When you are finished:

1) Before turning in your test **check carefully for transcribing errors**. Any mistakes you leave in are there to stay.

2) You must turn in your scantron to your discussion leader or exam proctor. Be prepared to show your picture I.D. with a legible signature.

3) The answers will be posted in Canvas within one day after the exam.

University of Florida Honor Pledge:

On my honor, I have neither given nor received unauthorized aid doing this exam.

Signature: _____

Summary of Integration Formulas

- Fundamental Theorem of Calculus

$$\int_a^b F'(x) dx = F(b) - F(a)$$

- Fundamental Theorem of Line Integrals

$$\int_C \nabla f \cdot d\vec{r} = f(\vec{r}(b)) - f(\vec{r}(a))$$

- Green's Theorem (circulation form)

$$\iint_D \text{curl } \vec{F} \cdot \hat{k} dA = \oint_C \vec{F} \cdot d\vec{r}$$

- Stokes' Theorem

$$\iint_S \text{curl } \vec{F} \cdot \hat{n} dS = \oint_C \vec{F} \cdot d\vec{r}$$

- Green's Theorem (flux form)

$$\iint_D \text{div } \vec{F} dA = \oint_C \vec{F} \cdot \hat{n} ds$$

- Divergence Theorem

$$\iiint_E \text{div } \vec{F} dV = \iint_S \vec{F} \cdot \hat{n} dS$$

NOTE: Be sure to bubble the answers to questions 1–24 on your scantron.

Questions 1 – 20 are worth 5 points each.

1. Let $\vec{F}(x, y, z) = \langle yz^2, x^2z, xy^2 \rangle$. Find $\text{curl } \vec{F}$ at $(1, 2, -1)$.

- a. $\langle -3, 8, -3 \rangle$
- b. $\langle 3, -8, 3 \rangle$
- c. $\langle 3, 8, -3 \rangle$
- d. $\langle 3, -8, -3 \rangle$
- e. $\langle 3, 8, 3 \rangle$

2. Which of the following vector fields are **conservative** on their respective given regions?

$$\vec{F} = \langle -y, x \rangle \text{ on } \mathbb{R}^2$$

$$\vec{G} = \left\langle \frac{y}{x}, \ln(x) \right\rangle \text{ on } \{(x, y) \mid x > 1 \text{ and } y > -1\}$$

$$\vec{H} = \langle ye^{xy}, xe^{xy} \rangle \text{ on } \mathbb{R}^2$$

- a. $\vec{F}$ and $\vec{H}$ only
- b. $\vec{G}$ only
- c. $\vec{G}$ and $\vec{H}$ only
- d. $\vec{F}$ and $\vec{G}$ only
- e. $\vec{F}$, $\vec{G}$, and $\vec{H}$

3. The figure shows a vector field $\vec{F}$ and two curves C_1 and C_2 . Which of the following is correct?


a. $\int_{C_1} \vec{F} \cdot d\vec{r} > \int_{C_2} \vec{F} \cdot d\vec{r} > 0$

b. $\int_{C_1} \vec{F} \cdot d\vec{r} > 0 > \int_{C_2} \vec{F} \cdot d\vec{r}$

c. $0 > \int_{C_2} \vec{F} \cdot d\vec{r} > \int_{C_1} \vec{F} \cdot d\vec{r}$

d. $\int_{C_2} \vec{F} \cdot d\vec{r} > 0 > \int_{C_1} \vec{F} \cdot d\vec{r}$

e. None of the above



4. Suppose $F(x, y) = \langle -y, x \rangle$. Let C_1, C_2 and C_3 be the linear paths respectively parameterized by

$$\vec{r}_1(t) = \langle 2t, 2t \rangle, \quad \vec{r}_2(t) = \langle 2 - 4t, 2 \rangle, \quad \vec{r}_3(t) = \langle -2 + 2t, 2 - 2t \rangle$$

for $0 \leq t \leq 1$. If $C = C_1 \cup C_2 \cup C_3$, then evaluate $\int_C \vec{F} \cdot d\vec{r}$.

a. -8

b. -2

c. 0

d. 2

e. 8

5. Let S be the part of the cone $z = \sqrt{2x^2 + 2y^2}$ with $0 \leq z \leq 2$. Then the area of the surface $S =$

a. $\int_0^{2\pi} \int_0^1 \sqrt{2} \, dr \, d\theta$

b. $\int_0^{2\pi} \int_0^2 \sqrt{5} \, r \, dr \, d\theta$

c. $\int_0^{2\pi} \int_0^2 \sqrt{3} \, dr \, d\theta$

d. $\int_0^{2\pi} \int_0^{\sqrt{2}} \sqrt{2} \, r \, dr \, d\theta$

e. $\int_0^{2\pi} \int_0^{\sqrt{2}} \sqrt{3} \, r \, dr \, d\theta$

6. Suppose $f(x, y, z)$ is a potential function of $\vec{F}(x, y, z) = \langle 2x - z, z, -x + y \rangle$. If $f(1, 1, 0) = 3$, then what is $f(-1, 2, 3)$?

a. 12

b. 6

c. 10

d. 2

e. -2

For Questions 7–8: Consider the surface S parameterized by

$$S : \vec{r}(u, v) = \langle 3 \cos(u), v, \sin(u) \rangle, \quad 0 \leq u \leq 2\pi, \quad -1 \leq v \leq 1.$$

7. Identify the surface S .

- a. Ellipse
- b. Cylinder
- c. Cone
- d. Paraboloid
- e. Ellipsoid

8. If $\hat{n}$ is the unit normal vector (with non-negative z component) to the surface S at $\vec{r}(\pi/4, 0)$, then which of the following vectors is in the same direction as $\hat{n}$?

- a. $\langle \sqrt{2}, 0, \sqrt{2} \rangle$
- b. $\langle -\sqrt{2}, 0, \sqrt{2} \rangle$
- c. $\langle -1, 0, 3 \rangle$
- d. $\langle 1, 0, 3 \rangle$
- e. $\langle 1, 0, \sqrt{2} \rangle$

9. Suppose $\vec{F}(x, y, z) = \langle xy^2, yz^2, zx^2 \rangle$ and let S be the union of the upper hemisphere of radius 2 centered at the origin with the disc of radius 2 in the xy -plane centered at the origin such that S is positively oriented. Evaluate $\iint_S \vec{F}(x, y, z) \cdot d\vec{S}$.

- a. 0
- b. $8\pi^2/3$
- c. $16\pi^2/3$
- d. $128\pi/5$
- e. $64\pi/5$

10. Which of the following regions is/are simply connected on $\mathbb{R}^2$?

$$P = \{(x, y) \mid x^2 + y^2 < 1\}$$

$$Q = \{(x, y) \mid 0 < x^2 + y^2 < 1\}$$

$$R = \{(x, y) \mid 1 < x^2 + y^2 < 9 \text{ and } y > 0\}$$

- a. P only
- b. Q only
- c. R only
- d. P and Q
- e. P and R

11. If S is the part of the paraboloid $z = 4 - x^2 - y^2$ with $z \geq 0$, $\vec{F}(x, y, z) = \langle -y, x, z \rangle$, and $\hat{n}$ is the upward unit normal on S , then $\iint_S (\text{curl } \vec{F}) \cdot \hat{n} \, dS =$

- a. 0
- b. 2π
- c. 4π
- d. 8π
- e. 16π

12. Suppose $\vec{F}(x, y)$ is a **conservative** vector field on $\mathbb{R}^2$. Let C_1 be parameterized by $\vec{r}_1(t) = \langle \cos(t), \sin(t) \rangle$ with $0 \leq t \leq \pi$, let C_2 be the line segment from $(0, -2)$ to $(-1, 0)$, and let C_3 be parameterized by $\vec{r}_3(t) = \langle \cos(t), 2 \sin(t) \rangle$ with $-\pi/2 \leq t \leq 0$. If $\int_{C_1} \vec{F} \cdot d\vec{r} = 4$ and $\int_{C_2} \vec{F} \cdot d\vec{r} = 3$, then what is $\int_{C_3} \vec{F} \cdot d\vec{r}$?

Hint: Draw the curves.

- a. 12
- b. -7
- c. -1
- d. 7
- e. 0

13. Let $E = \{(x, y, z) \mid x^2 + y^2 + z^2 \leq 1 \text{ and } z \geq 0\}$, the unit ball with non-negative z . Which of the following describes ∂E (the boundary of E)?

- a. $\{(x, y, z) \mid x^2 + y^2 + z^2 = 1, z \geq 0\}$
 - b. $\{(x, y, z) \mid x^2 + y^2 \leq 1, z = 0\}$
 - c. $\{(x, y, z) \mid x^2 + y^2 + z^2 = 1, z \geq 0\} \cup \{(x, y, z) \mid x^2 + y^2 \leq 1, z = 0\}$
 - d. $\{(x, y, z) \mid x^2 + y^2 = 1, z = 0\}$
 - e. $\{(x, y, z) \mid x^2 + y^2 = 1, z = 0\} \cup \{(x, y, z) \mid x^2 + y^2 + z^2 = 1, z > 0\}$
-

14. Find the work done by the force field $\vec{F} = yz\hat{i} + xz\hat{j} + xy\hat{k}$ in moving a particle along the curve $\vec{r}(t) = \langle \cos(t), \cos^2(t), \cos^5(2t) \rangle$, $0 \leq t \leq \pi/2$.

- a. -1
- b. -2
- c. 0
- d. 1
- e. 2

15. Evaluate $\int_C y \sin(z) \, ds$, where C is the circular helix given by $\vec{r}(t) = \langle \cos(t), \sin(t), t \rangle$, $0 \leq t \leq 4\pi$.

- a. π
- b. 2π
- c. $\sqrt{2} \pi$
- d. $2\sqrt{2} \pi$
- e. $4\sqrt{2} \pi$

16. Suppose $\vec{F}(x, y, z)$ is a vector field with continuous second partial derivatives on $\mathbb{R}^3$. Which of the following must be true?

P. $\text{curl } F = \vec{0}$

Q. If $\text{div } F = 0$, then there exists a vector field G such that $F = \text{curl } G$.

R. $\text{curl}(\text{curl } F) = \vec{0}$

- a. Q only
- b. R only
- c. P and R only
- d. P, Q, and R
- e. None of them

17. Use Stoke's Theorem to set up an iterated double integral for $\int_C \vec{F} \cdot d\vec{r}$ if


$$\vec{F}(x, y, z) = \langle x + y, -yz, xz \rangle$$

and C is the positive oriented curve of the intersection of the plane $z = x + 1$ and the cylinder $x^2 + y^2 = 1$.

- a. $\int_0^{2\pi} \int_0^1 r(r \sin \theta + 1) dr d\theta$
- b. $\int_0^{2\pi} \int_0^1 r(r \cos \theta + 1) dr d\theta$
- c. $\int_0^{2\pi} \int_0^1 r(-r \sin \theta - 1) dr d\theta$
- d. $\int_0^{2\pi} \int_0^1 r^2(-r \cos \theta - 1) dr d\theta$
- e. $\int_0^{2\pi} \int_0^1 r^2 \cos \theta dr d\theta$

18. The vector field $\vec{F}(x, y)$ is shown below. Determine the sign of $\text{div } \vec{F}$ at $(0, 0)$ and $(8, 8)$.

- a. Both are positive
- b. Both are negative
- c. $\text{div } \vec{F} > 0$ at $(0, 0)$ and $\text{div } \vec{F} < 0$ at $(8, 8)$
- d. $\text{div } \vec{F} < 0$ at $(0, 0)$ and $\text{div } \vec{F} > 0$ at $(8, 8)$
- e. Cannot determine the sign



19. Let $\vec{F} = \langle xy, x^2 + y^2 \rangle$ be the velocity field of a two-dimensional fluid flow. If D is the semiannular region in the upper half-plane between circles $x^2 + y^2 = 1$ and $x^2 + y^2 = 4$ with a positively oriented boundary, ∂D . Then the flux of $\vec{F}$ across the curve ∂D is $\oint_{\partial D} \vec{F} \cdot \vec{n} ds =$


a. $\int_0^{2\pi} \int_1^2 3r^2 \cos \theta dr d\theta$

b. $\int_0^\pi \int_1^2 3r^2 \sin \theta dr d\theta$

c. $\int_0^\pi \int_0^2 r^2 \cos \theta dr d\theta$

d. $\int_0^\pi \int_1^2 r^2 \sin \theta dr d\theta$

e. $\int_0^{2\pi} \int_1^2 3r \cos \theta dr d\theta$



20. Which of the following is correct?

- a. If $\vec{F}(x, y, z) = \langle 1, 0, 0 \rangle$, then the flux of $\vec{F}$ across the yz -plane is zero.
- b. If S is a surface parameterized by $\vec{r}(u, v)$, then the vector $\vec{r}_u \times \vec{r}_v$ lies in the tangent plane of S at a given point.
- c. The vector field $\vec{F}(x, y, z) = \langle x, x, x \rangle$ is independent of path.
- d. There is a vector field $\vec{F}$ on $\mathbb{R}^3$ such that $\text{curl } \vec{F} = \langle 2x, -y, -z \rangle$.
- e. All of the above are correct.

Bonus Questions 21 – 24 are worth 3 points each.

21. Let $\vec{u} = \langle 3, 6, -9 \rangle$ and $\vec{v} = \langle 1, 1, -1 \rangle$. If $\vec{u} = \vec{v}_{//} + \vec{v}_{\perp}$, where $\vec{v}_{//}$ is parallel to $\vec{v}$ and $\vec{v}_{\perp}$ is perpendicular to $\vec{v}$, and $\vec{v}_{\perp} = \langle a, b, c \rangle$, then what is a ?

- a. -3
 - b. 0
 - c. 6
 - d. -6
 - e. 3
-

22. Consider the surface $4x^2 + y^2 - z = 0$. Which of the following is/are correct?

P. The graph of the surface is a paraboloid.

Q. The level curves are hyperbolas.

R. The vertical trace in the the yz -plane is a parabola.

- a. P only
- b. Q only
- c. P and R only
- d. Q and R only
- e. P, Q, and R

23. Let E be the solid region bounded by the paraboloid $z = 2x^2 + 2y^2$ and the plane $z = 10$. Which of the following integrals represents the volume of E ?

a. $\int_0^{2\pi} \int_0^5 (10r - 2r^3) \, dr \, d\theta$

b. $\int_0^{2\pi} \int_0^{\sqrt{5}} (10r - 2r^3) \, dr \, d\theta$

c. $\int_0^{2\pi} \int_0^{10} (2r^3 - 10r) \, dr \, d\theta$

d. $\int_0^{2\pi} \int_0^{\sqrt{5}} (2r^2 - 10) \, dr \, d\theta$

e. $\int_0^{2\pi} \int_0^5 (10 - 2r^2) \, dr \, d\theta$

24. Find the absolute maximum value of $f(x, y) = 1 + xy^2$ on the set $D = \{(x, y) \mid x^2 + y^2 \leq 3, x \geq 0, y \geq 0\}$.

a. 1

b. 2

c. 3

d. 4

e. 5