

Module 7 Lecture Notes

MAC1105

Summer B 2019

7 Rational Functions

7.1 Domain of Rational Functions

Definition


A _____ is a vertical line $x = a$ where the graph tends toward positive or negative infinity as the input approaches a . We write

As $x \rightarrow a$, _____

or

As $x \rightarrow a$, _____

Example 1. Consider the graph of the function $f(x) = \frac{1}{x}$:



We cannot divide by zero, so as $x \rightarrow 0^-$, _____. That is, as x approaches zero from the left, $f(x)$ approaches $-\infty$. Similarly, as $x \rightarrow 0^+$, _____. That is, as x approaches zero from the right, $f(x)$ approaches ∞ . So, there is a vertical asymptote at _____.

Note 1. A vertical asymptote represents a value at which a rational function is _____, so the value is not in the domain of the function. In general, to find the domain of a rational function we need to find the values which cause the _____ to be equal to zero.

How to Find the Domain of a Rational Function

1. Set the _____ equal to zero.
2. Solve to find the x -values that make the _____ equal to zero.
3. The domain is _____ except those found in step 2.

Example 2. Find the domain of the rational function:

$$f(x) = \frac{1}{x+3}$$

Note 2. It does not matter what type of real number makes the denominator equal to zero. When we have two or more fractions in the function, the domain is the _____ of the domains of each function separately. In other words, we will remove EVERY value of the function that makes the denominator equal to zero from the domain of the function.

Example 3. Find the domain of the following function:

$$f(x) = \frac{1}{3x + 2} - \frac{1}{-3x - 5}$$

Note 3. If we are given a rational function which has a denominator with degree larger than 1, we will first need to _____ the denominator and then solve for the x -values that make the denominator equal to _____.

Example 4. Find the domain of the rational function:

$$f(x) = \frac{1}{x^2 + 4x + 3}$$

Example 5. Find the domain of the rational function:

$$f(x) = \frac{1}{2x^3 - 3x^2 - 9x}$$

Example 6. Find the domain of the rational function:


$$f(x) = \frac{1}{x^3 + 13x}$$

Definition

A _____ is a horizontal line $y = b$ where the graph approaches the line as the inputs increase or decrease without bound. We write

As $x \rightarrow \infty$ or $x \rightarrow -\infty$, _____

Example 7. Recall the graph of $f(x) = \frac{1}{x}$:



As $x \rightarrow \infty$, _____. That is, as x approaches infinity, $f(x)$ approaches zero.


Similarly, as $x \rightarrow -\infty$, _____. That is, as x approaches ∞ , $f(x)$ approaches zero.

So, there is a horizontal asymptote at _____.

7.2 Graphing Rational Functions

Graphs of Basic Rational Functions To graph rational functions in this section, we will work

with two basic rational functions: $f(x) = \frac{1}{x}$ and $f(x) = \frac{1}{x^2}$.



Graphing Rational Functions With Translations

Consider the function $f(x) = \frac{a}{x-h} + k$.

Horizontal Shifts

Vertical Shifts

Stretch Factor

Note 4. So, we can now use the graph of $\frac{1}{x}$ to graph $f(x) = \frac{a}{x-h} + k$. When translating the graph, we must remember to shift horizontal and/or vertical asymptotes.

Note 5. To find the value of a , plug in a point on the graph after finding the value of h and k .

Graphing Rational Functions With Translations

Consider the function $f(x) = \frac{a}{(x-h)^2} + k$.

Horizontal Shifts

Vertical Shifts

Stretch Factor

Standard Form of Rational Functions

The standard form of $f(x) = \frac{a}{x-h} + k$ is

$$f(x) = \text{_____} +$$


The standard form of $f(x) = \frac{a}{(x-h)^2} + k$ is

$$f(x) = \text{_____} +$$


Intercepts of Rational Functions

If a rational function does not have a vertical asymptote at $x = 0$ (that is, if our function is defined at $x = 0$), then the y -intercept of the function occurs when _____. If a rational function does not have a horizontal asymptote at $y = 0$ (that is if our function is defined for all x -values where $f(x) = 0$), then the x -intercept of the function occurs when _____.

Example 8. Write an equation for the graph of the rational function below. Assume $a = 1$ or $a = -1$:



Example 9. Write an equation for the graph of the rational function below:



7.3 Solving Rational Equations

How to Solve Rational Equations

1. Identify the _____ (ie find the values of x for which the _____ are equal to 0).
2. Find the least common denominator.
3. Multiply both sides by the _____
_____.
4. You should now have a linear or quadratic equation. Simplify your remaining equation and solve.
5. Check your solutions and check that your solutions are in the _____.

Example 10. Solve the rational equation below:

$$\frac{5}{x} - \frac{1}{3} = \frac{1}{x}$$

Example 11. Solve the rational equation below:

$$-\frac{8}{8x-5} + 1 = \frac{32}{-32x+20}$$

Example 12. Solve the rational equation below:

$$-\frac{2x^2}{3(2x^2 + x - 10)} - \frac{4x}{3(x - 2)} = -\frac{3}{2x + 5}$$

Example 13. Solve the rational equation below:

$$\frac{10}{x(x-2)} + \frac{4}{x} = \frac{5}{(x-2)}$$