

The graph isomorphism game

Samuel Harris

Texas A&M University

OTWIA

August 13, 2020

Joint work with Alexandru Chirvasitu, Michael Brannan, Kari Eifler,
Vern Paulsen, Xiaoyu Su, and Mateusz Wasilewski

Outline

- 1 Background
- 2 Quantum Isomorphisms
- 3 Main Results

Non-local games: a primer

- We look at two-player non-local games. The referee sends numbers $x, y \in I = \{1, \dots, m\}$ to Alice and Bob.

Non-local games: a primer

- We look at two-player non-local games. The referee sends numbers $x, y \in I = \{1, \dots, m\}$ to Alice and Bob.
- Alice and Bob respond with numbers $a, b \in O = \{1, \dots, k\}$.

Non-local games: a primer


- We look at two-player non-local games. The referee sends numbers $x, y \in I = \{1, \dots, m\}$ to Alice and Bob.
- Alice and Bob respond with numbers $a, b \in O = \{1, \dots, k\}$.
- Rule function $\Lambda : I \times I \times O \times O \rightarrow \{0, 1\}$. Alice and Bob win if $\Lambda(a, b, x, y) = 1$, and lose if $\Lambda(a, b, x, y) = 0$.

Non-local games: a primer

- We look at two-player non-local games. The referee sends numbers $x, y \in I = \{1, \dots, m\}$ to Alice and Bob.
- Alice and Bob respond with numbers $a, b \in O = \{1, \dots, k\}$.
- Rule function $\Lambda : I \times I \times O \times O \rightarrow \{0, 1\}$. Alice and Bob win if $\Lambda(a, b, x, y) = 1$, and lose if $\Lambda(a, b, x, y) = 0$.
- Catch: Alice and Bob can agree upon a strategy before the game, but *cannot* communicate once it begins.

Non-local games: a primer

- We look at two-player non-local games. The referee sends numbers $x, y \in I = \{1, \dots, m\}$ to Alice and Bob.
- Alice and Bob respond with numbers $a, b \in O = \{1, \dots, k\}$.
- Rule function $\Lambda : I \times I \times O \times O \rightarrow \{0, 1\}$. Alice and Bob win if $\Lambda(a, b, x, y) = 1$, and lose if $\Lambda(a, b, x, y) = 0$.
- Catch: Alice and Bob can agree upon a strategy before the game, but *cannot* communicate once it begins.



Example: $\text{Iso}(G, H)$

Let G and H be two finite, undirected graphs with $|V(G)| = |V(H)|$. (No multiple edges, no loops)

Example: $\text{Iso}(G, H)$

Let G and H be two finite, undirected graphs with $|V(G)| = |V(H)|$. (No multiple edges, no loops)

- Input set is $I = V(G) \sqcup V(H)$. The referee sends vertices x, y to Alice and Bob, respectively.

Example: $\text{Iso}(G, H)$

Let G and H be two finite, undirected graphs with $|V(G)| = |V(H)|$. (No multiple edges, no loops)

- Input set is $I = V(G) \sqcup V(H)$. The referee sends vertices x, y to Alice and Bob, respectively.
- Alice and Bob each respond with a vertex a and b , respectively, from $O = V(G) \sqcup V(H)$.

Example: $\text{Iso}(G, H)$

Let G and H be two finite, undirected graphs with $|V(G)| = |V(H)|$. (No multiple edges, no loops)

- Input set is $I = V(G) \sqcup V(H)$. The referee sends vertices x, y to Alice and Bob, respectively.
- Alice and Bob each respond with a vertex a and b , respectively, from $O = V(G) \sqcup V(H)$.
- Alice and Bob win the game if:

Example: $\text{Iso}(G, H)$

Let G and H be two finite, undirected graphs with $|V(G)| = |V(H)|$. (No multiple edges, no loops)

- Input set is $I = V(G) \sqcup V(H)$. The referee sends vertices x, y to Alice and Bob, respectively.
- Alice and Bob each respond with a vertex a and b , respectively, from $O = V(G) \sqcup V(H)$.
- Alice and Bob win the game if:
 - a and x are in different graphs, and b and y are in different graphs; *and*

Example: $\text{Iso}(G, H)$

Let G and H be two finite, undirected graphs with $|V(G)| = |V(H)|$. (No multiple edges, no loops)

- Input set is $I = V(G) \sqcup V(H)$. The referee sends vertices x, y to Alice and Bob, respectively.
- Alice and Bob each respond with a vertex a and b , respectively, from $O = V(G) \sqcup V(H)$.
- Alice and Bob win the game if:
 - a and x are in different graphs, and b and y are in different graphs; *and*
 - Writing $\{a, x\} = \{g_A, h_A\}$ and $\{b, y\} = \{g_B, h_B\}$ where $g_A, g_B \in V(G)$, $h_A, h_B \in V(H)$, we require $\text{rel}(g_A, g_B) = \text{rel}(h_A, h_B)$, where $\text{rel}(x, x') = 1$ if $x \sim x'$, 0 if $x = x'$, and -1 if $x \neq x'$ and $x \not\sim x'$.

Example: $\text{Iso}(G, H)$

Let G and H be two finite, undirected graphs with $|V(G)| = |V(H)|$. (No multiple edges, no loops)

- Input set is $I = V(G) \sqcup V(H)$. The referee sends vertices x, y to Alice and Bob, respectively.
- Alice and Bob each respond with a vertex a and b , respectively, from $O = V(G) \sqcup V(H)$.
- Alice and Bob win the game if:
 - a and x are in different graphs, and b and y are in different graphs; *and*
 - Writing $\{a, x\} = \{g_A, h_A\}$ and $\{b, y\} = \{g_B, h_B\}$ where $g_A, g_B \in V(G)$, $h_A, h_B \in V(H)$, we require $\text{rel}(g_A, g_B) = \text{rel}(h_A, h_B)$, where $\text{rel}(x, x') = 1$ if $x \sim x'$, 0 if $x = x'$, and -1 if $x \neq x'$ and $x \not\sim x'$.

Recall that Alice and Bob *cannot communicate* once the game begins. (But they can share an entanglement resource space!)

Probabilistic models

- There are several different possibilities for modelling the probability distributions $p(a, b|x, y)$ for an m -input, k -output system (eg. for the graph isomorphism game.)

Probabilistic models

- There are several different possibilities for modelling the probability distributions $p(a, b|x, y)$ for an m -input, k -output system (eg. for the graph isomorphism game.)
- We consider when $p(a, b|x, y) = \langle E_{a,x} F_{b,y} \psi, \psi \rangle$, where ψ is a unit vector in a Hilbert space \mathcal{H} , $\{E_{a,x}\}_{a=1}^m$ and $\{F_{b,y}\}_{b=1}^m$ are projection-valued measures for each x, y , and $[E_{a,x}, F_{b,y}] = 0$ for all a, b, x, y .

Probabilistic models

- There are several different possibilities for modelling the probability distributions $p(a, b|x, y)$ for an m -input, k -output system (eg. for the graph isomorphism game.)
- We consider when $p(a, b|x, y) = \langle E_{a,x} F_{b,y} \psi, \psi \rangle$, where ψ is a unit vector in a Hilbert space \mathcal{H} , $\{E_{a,x}\}_{a=1}^m$ and $\{F_{b,y}\}_{b=1}^m$ are projection-valued measures for each x, y , and $[E_{a,x}, F_{b,y}] = 0$ for all a, b, x, y .
- We could model (a subset of) these probability distributions using a finite-dimensional tensor product model (giving $C_q(m, k)$), a tensor product model (giving $C_{qs}(m, k)$), an approximate tensor product model (giving $C_{qa}(m, k)$), a commuting model (giving $C_{qc}(m, k)$), or a local model (giving $C_{loc}(m, k)$).

Probabilistic models

- There are several different possibilities for modelling the probability distributions $p(a, b|x, y)$ for an m -input, k -output system (eg. for the graph isomorphism game.)
- We consider when $p(a, b|x, y) = \langle E_{a,x} F_{b,y} \psi, \psi \rangle$, where ψ is a unit vector in a Hilbert space \mathcal{H} , $\{E_{a,x}\}_{a=1}^m$ and $\{F_{b,y}\}_{b=1}^m$ are projection-valued measures for each x, y , and $[E_{a,x}, F_{b,y}] = 0$ for all a, b, x, y .
- We could model (a subset of) these probability distributions using a finite-dimensional tensor product model (giving $C_q(m, k)$), a tensor product model (giving $C_{qs}(m, k)$), an approximate tensor product model (giving $C_{qa}(m, k)$), a commuting model (giving $C_{qc}(m, k)$), or a local model (giving $C_{loc}(m, k)$).
- **By a lot of theorems**, we know that

$$C_{loc}(m, k) \subseteq C_q(m, k) \subseteq C_{qs}(m, k) \subseteq C_{qa}(m, k) \subseteq C_{qc}(m, k),$$

and *all* of these containments are (in general) strict. ▶ ◀ ≡ ≡ ≡ ↺ ↻ ↻

The synchronicity condition

- Key property: $\text{Iso}(G, H)$ is a **synchronous game**: if Alice and Bob receive inputs x, y and $x = y$, then Alice and Bob's outputs a and b *must* be equal. That is, if they win with probability 1, then

$$p(a, b|x, x) = 0 \text{ if } a \neq b.$$

The synchronicity condition

- Key property: $\text{Iso}(G, H)$ is a **synchronous game**: if Alice and Bob receive inputs x, y and $x = y$, then Alice and Bob's outputs a and b *must* be equal. That is, if they win with probability 1, then

$$p(a, b|x, x) = 0 \text{ if } a \neq b.$$

- For $t \in \{loc, q, qs, qa, qc\}$, one defines

$$C_t^s(m, k) = \{(p(a, b|x, y)) \in C_t(m, k) : p \text{ is synchronous}\}.$$

The synchronicity condition

- Key property: $\text{Iso}(G, H)$ is a **synchronous game**: if Alice and Bob receive inputs x, y and $x = y$, then Alice and Bob's outputs a and b *must* be equal. That is, if they win with probability 1, then

$$p(a, b|x, x) = 0 \text{ if } a \neq b.$$

- For $t \in \{loc, q, qs, qa, qc\}$, one defines

$$C_t^s(m, k) = \{(p(a, b|x, y)) \in C_t(m, k) : p \text{ is synchronous}\}.$$

- Surprisingly, $C_q^s(m, k) = C_{qs}^s(m, k)$ (Kim-Paulsen-Schafhauser, '18), but all the other analogous containments are (in general) strict.

Synchronous realizations

Theorem (Paulsen, Severini, Stahlke, Todorov, Winter, '16)

$(p(a, b|x, y)) \in C_{qc}^s(m, k) \iff$ *there are projection-valued measures $\{E_{a,x}\}_{a=1}^k$ for each x in a tracial C^* -algebra (\mathcal{A}, τ) , such that*

$$p(a, b|x, y) = \tau(E_{a,x}E_{b,y}).$$

Moreover,

- $(p(a, b|x, y)) \in C_{loc}^s(m, k) \iff \mathcal{A}$ *can be taken to be abelian;*
- $(p(a, b|x, y)) \in C_q^s(m, k) \iff \mathcal{A}$ *can be taken to be finite-dimensional;*
- (Kim-Paulsen-Schafhauser, '18) $(p(a, b|x, y)) \in C_{qa}^s(m, k) \iff \mathcal{A}$ *can be taken to be \mathcal{R}^ω (ultrapower of hyperfinite II_1 -factor).*

Quantum permutations

- Let \mathcal{A} be a unital C^* -algebra. A matrix $U = (E_{g,h}) \in M_n(\mathcal{A})$ is a **quantum permutation** if $U^*U = UU^* = I_n$ and $E_{g,h} = E_{g,h}^* = E_{g,h}^2$ for all g, h .

Quantum permutations

- Let \mathcal{A} be a unital C^* -algebra. A matrix $U = (E_{g,h}) \in M_n(\mathcal{A})$ is a **quantum permutation** if $U^*U = UU^* = I_n$ and $E_{g,h} = E_{g,h}^* = E_{g,h}^2$ for all g, h .
- Equivalently, $E_{g,h} = E_{g,h}^* = E_{g,h}^2$ and $\sum_{g \in V(G)} E_{g,h} = \sum_{h \in V(G)} E_{g,h} = 1$ for all g, h .

Quantum permutations

- Let \mathcal{A} be a unital C^* -algebra. A matrix $U = (E_{g,h}) \in M_n(\mathcal{A})$ is a **quantum permutation** if $U^*U = UU^* = I_n$ and $E_{g,h} = E_{g,h}^* = E_{g,h}^2$ for all g, h .
- Equivalently, $E_{g,h} = E_{g,h}^* = E_{g,h}^2$ and $\sum_{g \in V(G)} E_{g,h} = \sum_{h \in V(G)} E_{g,h} = 1$ for all g, h .
- $p = (p(a, b|x, y)) = \tau(E_{a,x}E_{b,y}) \in C_t^s(|V(G) \sqcup V(H)|, |V(G) \sqcup V(H)|)$ wins the game $\text{Iso}(G, H)$ with probability 1 if and only if $U = (E_{g,h})_{g \in V(G), h \in V(H)}$ is a quantum permutation such that

$$(A_G \otimes 1)U = U(A_H \otimes 1),$$

where A_G (resp. A_H) is the adjacency matrix of G (resp. H).

- Write $G \simeq_t H$ if there's a winning t -strategy for $\text{Iso}(G, H)$ (where $t \in \{\text{loc}, q, qa, qc\}$).

- Write $G \simeq_t H$ if there's a winning t -strategy for $\text{Iso}(G, H)$ (where $t \in \{loc, q, qa, qc\}$).
- $G \simeq_{loc} H \iff G \simeq H \iff$ there's a permutation matrix $U \in M_n$ with $A_G U = U A_H$.

- Write $G \simeq_t H$ if there's a winning t -strategy for $\text{Iso}(G, H)$ (where $t \in \{loc, q, qa, qc\}$).
- $G \simeq_{loc} H \iff G \simeq H \iff$ there's a permutation matrix $U \in M_n$ with $A_G U = U A_H$.
- $G \simeq_q H \iff$ there's a quantum permutation $U \in M_n(M_d)$ for some d , such that $(A_G \otimes 1)U = U(A_H \otimes 1)$.

- Write $G \simeq_t H$ if there's a winning t -strategy for $\text{Iso}(G, H)$ (where $t \in \{loc, q, qa, qc\}$).
- $G \simeq_{loc} H \iff G \simeq H \iff$ there's a permutation matrix $U \in M_n$ with $A_G U = U A_H$.
- $G \simeq_q H \iff$ there's a quantum permutation $U \in M_n(M_d)$ for some d , such that $(A_G \otimes 1)U = U(A_H \otimes 1)$.
- $G \simeq_{qa} H \iff$ we can take U in $M_n(\mathcal{R}^\omega)$.

- Write $G \simeq_t H$ if there's a winning t -strategy for $\text{Iso}(G, H)$ (where $t \in \{loc, q, qa, qc\}$).
- $G \simeq_{loc} H \iff G \simeq H \iff$ there's a permutation matrix $U \in M_n$ with $A_G U = U A_H$.
- $G \simeq_q H \iff$ there's a quantum permutation $U \in M_n(M_d)$ for some d , such that $(A_G \otimes 1)U = U(A_H \otimes 1)$.
- $G \simeq_{qa} H \iff$ we can take U in $M_n(\mathcal{R}^\omega)$.
- $G \simeq_{qc} H \iff$ we can take $U \in M_n(\mathcal{A})$ for some tracial C^* -algebra (\mathcal{A}, τ) .

- Write $G \simeq_t H$ if there's a winning t -strategy for $\text{Iso}(G, H)$ (where $t \in \{loc, q, qa, qc\}$).
- $G \simeq_{loc} H \iff G \simeq H \iff$ there's a permutation matrix $U \in M_n$ with $A_G U = U A_H$.
- $G \simeq_q H \iff$ there's a quantum permutation $U \in M_n(M_d)$ for some d , such that $(A_G \otimes 1)U = U(A_H \otimes 1)$.
- $G \simeq_{qa} H \iff$ we can take U in $M_n(\mathcal{R}^\omega)$.
- $G \simeq_{qc} H \iff$ we can take $U \in M_n(\mathcal{A})$ for some tracial C^* -algebra (\mathcal{A}, τ) .
- All of these cases can be thought of as representations of a universal $*$ -algebra for $\text{Iso}(G, H)$.

Algebra of the Game

Definition (Kim-Paulsen-Schafhauser, '18)

We define $\mathcal{A}(\text{Iso}(G, H))$ to be the free (unital) $*$ -algebra generated by entries $e_{g,h}$ ($g \in V(G)$, $h \in V(H)$) such that

$$e_{g,h}^2 = e_{g,h} = e_{g,h}^*, \quad \sum_{g \in V(G)} e_{g,h} = \sum_{h \in V(H)} e_{g,h} = 1,$$

and

$$e_{g,h} e_{g',h'} = 0 \text{ if } \text{rel}(g, g') \neq \text{rel}(h, h').$$

Say that $\mathcal{A}(\text{Iso}(G, H))$ **exists** if $1 \neq 0$ in $\mathcal{A}(\text{Iso}(G, H))$.

Algebra of the Game

Definition (Kim-Paulsen-Schafhauser, '18)

We define $\mathcal{A}(\text{Iso}(G, H))$ to be the free (unital) $*$ -algebra generated by entries $e_{g,h}$ ($g \in V(G)$, $h \in V(H)$) such that

$$e_{g,h}^2 = e_{g,h} = e_{g,h}^*, \quad \sum_{g \in V(G)} e_{g,h} = \sum_{h \in V(H)} e_{g,h} = 1,$$

and

$$e_{g,h} e_{g',h'} = 0 \text{ if } \text{rel}(g, g') \neq \text{rel}(h, h').$$

Say that $\mathcal{A}(\text{Iso}(G, H))$ **exists** if $1 \neq 0$ in $\mathcal{A}(\text{Iso}(G, H))$.

- $G \simeq H \iff \exists \text{ unital } *-hom \mathcal{A}(\text{Iso}(G, H)) \rightarrow \mathbb{C}.$

Algebra of the Game

Definition (Kim-Paulsen-Schafhauser, '18)

We define $\mathcal{A}(\text{Iso}(G, H))$ to be the free (unital) $*$ -algebra generated by entries $e_{g,h}$ ($g \in V(G)$, $h \in V(H)$) such that

$$e_{g,h}^2 = e_{g,h} = e_{g,h}^*, \quad \sum_{g \in V(G)} e_{g,h} = \sum_{h \in V(H)} e_{g,h} = 1,$$

and

$$e_{g,h} e_{g',h'} = 0 \text{ if } \text{rel}(g, g') \neq \text{rel}(h, h').$$

Say that $\mathcal{A}(\text{Iso}(G, H))$ **exists** if $1 \neq 0$ in $\mathcal{A}(\text{Iso}(G, H))$.

- $G \simeq H \iff \exists \text{ unital } *-hom \mathcal{A}(\text{Iso}(G, H)) \rightarrow \mathbb{C}.$
- $G \simeq_q H \iff \exists \text{ unital } *-hom \mathcal{A}(\text{Iso}(G, H)) \rightarrow M_d.$

Algebra of the Game

Definition (Kim-Paulsen-Schafhauser, '18)

We define $\mathcal{A}(\text{Iso}(G, H))$ to be the free (unital) $*$ -algebra generated by entries $e_{g,h}$ ($g \in V(G)$, $h \in V(H)$) such that

$$e_{g,h}^2 = e_{g,h} = e_{g,h}^*, \quad \sum_{g \in V(G)} e_{g,h} = \sum_{h \in V(H)} e_{g,h} = 1,$$

and

$$e_{g,h} e_{g',h'} = 0 \text{ if } \text{rel}(g, g') \neq \text{rel}(h, h').$$

Say that $\mathcal{A}(\text{Iso}(G, H))$ **exists** if $1 \neq 0$ in $\mathcal{A}(\text{Iso}(G, H))$.

- $G \simeq H \iff \exists \text{ unital } *-hom \mathcal{A}(\text{Iso}(G, H)) \rightarrow \mathbb{C}.$
- $G \simeq_q H \iff \exists \text{ unital } *-hom \mathcal{A}(\text{Iso}(G, H)) \rightarrow M_d.$
- $G \simeq_{qa} H \iff \exists \text{ unital } *-hom \mathcal{A}(\text{Iso}(G, H)) \rightarrow \mathcal{R}^\omega.$

Algebra of the Game

Definition (Kim-Paulsen-Schafhauser, '18)

We define $\mathcal{A}(\text{Iso}(G, H))$ to be the free (unital) $*$ -algebra generated by entries $e_{g,h}$ ($g \in V(G)$, $h \in V(H)$) such that

$$e_{g,h}^2 = e_{g,h} = e_{g,h}^*, \quad \sum_{g \in V(G)} e_{g,h} = \sum_{h \in V(H)} e_{g,h} = 1,$$

and

$$e_{g,h} e_{g',h'} = 0 \text{ if } \text{rel}(g, g') \neq \text{rel}(h, h').$$

Say that $\mathcal{A}(\text{Iso}(G, H))$ **exists** if $1 \neq 0$ in $\mathcal{A}(\text{Iso}(G, H))$.

- $G \simeq H \iff \exists \text{ unital } *- \text{hom } \mathcal{A}(\text{Iso}(G, H)) \rightarrow \mathbb{C}.$
- $G \simeq_q H \iff \exists \text{ unital } *- \text{hom } \mathcal{A}(\text{Iso}(G, H)) \rightarrow M_d.$
- $G \simeq_{qa} H \iff \exists \text{ unital } *- \text{hom } \mathcal{A}(\text{Iso}(G, H)) \rightarrow \mathcal{R}^\omega.$
- $G \simeq_{qc} H \iff \exists \text{ unital } *- \text{hom from } \mathcal{A}(\text{Iso}(G, H)) \rightarrow (\mathcal{A}, \tau).$

Algebra of the Game

Definition (Kim-Paulsen-Schafhauser, '18)

We define $\mathcal{A}(\text{Iso}(G, H))$ to be the free (unital) $*$ -algebra generated by entries $e_{g,h}$ ($g \in V(G)$, $h \in V(H)$) such that

$$e_{g,h}^2 = e_{g,h} = e_{g,h}^*, \quad \sum_{g \in V(G)} e_{g,h} = \sum_{h \in V(H)} e_{g,h} = 1,$$

and

$$e_{g,h} e_{g',h'} = 0 \text{ if } \text{rel}(g, g') \neq \text{rel}(h, h').$$

Say that $\mathcal{A}(\text{Iso}(G, H))$ **exists** if $1 \neq 0$ in $\mathcal{A}(\text{Iso}(G, H))$.

- $G \simeq H \iff \exists \text{ unital } *- \text{hom } \mathcal{A}(\text{Iso}(G, H)) \rightarrow \mathbb{C}.$
- $G \simeq_q H \iff \exists \text{ unital } *- \text{hom } \mathcal{A}(\text{Iso}(G, H)) \rightarrow M_d.$
- $G \simeq_{qa} H \iff \exists \text{ unital } *- \text{hom } \mathcal{A}(\text{Iso}(G, H)) \rightarrow \mathcal{R}^\omega.$
- $G \simeq_{qc} H \iff \exists \text{ unital } *- \text{hom from } \mathcal{A}(\text{Iso}(G, H)) \rightarrow (\mathcal{A}, \tau).$
- We say that $G \simeq_{alg} H$ if $\mathcal{A}(\text{Iso}(G, H))$ exists.

Algebraic Isomorphisms

Definition

We define $\mathcal{A}(\text{Iso}(G, H))$ to be the free (unital) $*$ -algebra generated by entries $e_{g,h}$ ($g \in V(G)$, $h \in V(H)$) such that

$$e_{g,h}^2 = e_{g,h} = e_{g,h}^*, \quad \sum_{g \in V(G)} e_{g,h} = \sum_{h \in V(H)} e_{g,h} = 1,$$

and

$$e_{g,h} e_{g',h'} = 0 \text{ if } \text{rel}(g, g') \neq \text{rel}(h, h').$$

Say that $\mathcal{A}(\text{Iso}(G, H))$ **exists** if $1 \neq 0$ in $\mathcal{A}(\text{Iso}(G, H))$.

$$G \simeq H \Rightarrow G \simeq_q H \Rightarrow G \simeq_{qa} H \Rightarrow G \simeq_{qc} H \Rightarrow G \simeq_{alg} H$$

$$\neq \qquad \qquad \neq \qquad \qquad \Leftarrow ?$$

Algebraic Isomorphisms

Definition

We define $\mathcal{A}(\text{Iso}(G, H))$ to be the free (unital) $*$ -algebra generated by entries $e_{g,h}$ ($g \in V(G)$, $h \in V(H)$) such that

$$e_{g,h}^2 = e_{g,h} = e_{g,h}^*, \quad \sum_{g \in V(G)} e_{g,h} = \sum_{h \in V(H)} e_{g,h} = 1,$$

and

$$e_{g,h} e_{g',h'} = 0 \text{ if } \text{rel}(g, g') \neq \text{rel}(h, h').$$

Say that $\mathcal{A}(\text{Iso}(G, H))$ **exists** if $1 \neq 0$ in $\mathcal{A}(\text{Iso}(G, H))$.

- Write $G \simeq_{alg} H$ if $\mathcal{A}(\text{Iso}(G, H))$ exists.

$$G \simeq H \Rightarrow G \simeq_q H \Rightarrow G \simeq_{qa} H \Rightarrow G \simeq_{qc} H \Rightarrow G \simeq_{alg} H$$

$\neq \qquad \qquad \neq \qquad \qquad \Leftarrow ?$

Algebraic Isomorphisms, continued

- It's not known if there are any graphs G, H with $G \simeq_{qc} H$ but $G \not\simeq_{qa} H$. (It's possible that Connes' embedding has a negative answer *and* qa/qc isomorphisms are the same for all G, H !)

Algebraic Isomorphisms, continued

- It's not known if there are any graphs G, H with $G \simeq_{qc} H$ but $G \not\simeq_{qa} H$. (It's possible that Connes' embedding has a negative answer *and* qa/qc isomorphisms are the same for all G, H !)
- What about $\simeq_{alg} \implies \simeq_{qc}$? (What could go wrong?)

Algebraic Isomorphisms, continued

- It's not known if there are any graphs G, H with $G \simeq_{qc} H$ but $G \not\simeq_{qa} H$. (It's possible that Connes' embedding has a negative answer *and* qa/qc isomorphisms are the same for all G, H !)
- What about $\simeq_{alg} \implies \simeq_{qc}$? (What could go wrong?)
- Eg. There's a graph *homomorphism* game, with an associated $*$ -algebra $\mathcal{A}(\text{Hom}(G, H))$. But $\mathcal{A}(\text{Hom}(K_5, K_4)) \neq 0$ (Helton-Meyer-Paulsen-Satriano '19), so there's an (algebraic) 4-coloring of K_5 !

Algebraic Isomorphisms, continued

- It's not known if there are any graphs G, H with $G \simeq_{qc} H$ but $G \not\simeq_{qa} H$. (It's possible that Connes' embedding has a negative answer *and* qa/qc isomorphisms are the same for all G, H !)
- What about $\simeq_{alg} \implies \simeq_{qc}$? (What could go wrong?)
- Eg. There's a graph *homomorphism* game, with an associated $*$ -algebra $\mathcal{A}(\text{Hom}(G, H))$. But $\mathcal{A}(\text{Hom}(K_5, K_4)) \neq 0$ (Helton-Meyer-Paulsen-Satriano '19), so there's an (algebraic) 4-coloring of K_5 !
- What's the problem? In a C^* -algebra, if p_1, \dots, p_n are self-adjoint idempotents and $\sum_{i=1}^n p_i = 1$, then $p_i p_j = 0$ if $i \neq j$. In a unital $*$ -algebra, this only works if $n \leq 3$, and is *false* if $n \geq 4$. (In $\mathcal{A}(\text{Hom}(K_5, K_4))$, there are p_1, \dots, p_4 with $p_i^2 = p_i = p_i^*$ and $\sum p_i = -1$.)

$$\simeq_{alg} \implies \simeq_{qc}$$

Theorem (Chirvasitu-Brannan-Eifler-H.-Paulsen-Su-Wasilewski, '19)

If G and H are graphs with $G \simeq_{alg} H$, then $G \simeq_{qc} H$.

$$\simeq_{alg} \implies \simeq_{qc}$$

Theorem (Chirvasitu-Brannan-Eifler-H.-Paulsen-Su-Wasilewski, '19)

If G and H are graphs with $G \simeq_{alg} H$, then $G \simeq_{qc} H$.

- This also means that if $G \simeq_{C^*} H$ (i.e. there's a representation $\mathcal{A}(\text{Iso}(G, H)) \rightarrow \mathcal{B}(\mathcal{H})$ for some Hilbert space \mathcal{H}), then $G \simeq_{qc} H$.

Related: Lovasz's theorem on homomorphism counts

Theorem (Lovasz, 1967)

If G and H are graphs, then $G \simeq H$ if and only if $|\text{Hom}(Z, G)| = |\text{Hom}(Z, H)|$ for all graphs Z .

Related: Lovasz's theorem on homomorphism counts

Theorem (Lovasz, 1967)

If G and H are graphs, then $G \simeq H$ if and only if $|\text{Hom}(Z, G)| = |\text{Hom}(Z, H)|$ for all graphs Z .

- Question: can one relax the theorem to *planar* graphs Z ?

Related: Lovasz's theorem on homomorphism counts

Theorem (Lovasz, 1967)

If G and H are graphs, then $G \simeq H$ if and only if $|\text{Hom}(Z, G)| = |\text{Hom}(Z, H)|$ for all graphs Z .

- Question: can one relax the theorem to *planar* graphs Z ?

Theorem (Mancinska-Roberson, '19)

Let G and H be graphs. Then $G \simeq_{qc} H$ if and only if $|\text{Hom}(Z, G)| = |\text{Hom}(Z, H)|$ for all planar graphs Z .

Related: Lovasz's theorem on homomorphism counts

Theorem (Lovasz, 1967)

If G and H are graphs, then $G \simeq H$ if and only if $|\text{Hom}(Z, G)| = |\text{Hom}(Z, H)|$ for all graphs Z .

- Question: can one relax the theorem to *planar* graphs Z ?

Theorem (Mancinska-Roberson, '19)

Let G and H be graphs. Then $G \simeq_{qc} H$ if and only if $|\text{Hom}(Z, G)| = |\text{Hom}(Z, H)|$ for all planar graphs Z .

Thus, $G \simeq_{alg} H \iff G \simeq_{qc} H \iff$
 $|\text{Hom}(Z, G)| = |\text{Hom}(Z, H)|$ for all planar graphs Z .