The Spectral Radius of A Planar Graph

Dasong Cao and Andrew Vince Department of Mathematics University of Florida Gainesville, Florida 32611

Submitted by Richard A. Brualdi

ABSTRACT

A decomposition result for planar graphs is used to prove that the spectral radius of a planar graph on n vertices is less than $4 + \sqrt{3(n-3)}$. Moreover, the spectral radius of an outerplanar graph on n vertices is less than $1 + \sqrt{2 + \sqrt{2}} + \sqrt{n-5}$.

1. INTRODUCTION

All graphs are finite, undirected, without loops and multiple edges. Let G be a graph with vertices v_1, v_2, \ldots, v_n . The *complement* in G of a subgraph H is the subgraph of G obtained by deleting all edges in H. The *join* $G_1 \vee G_2$ of two graphs G_1 and G_2 is obtained by adding an edge from each vertex in G_1 to each vertex in G_2 . Let K_n be the complete graph and P_n the path with n vertices. Let $\delta(G)$ and $\Delta(G)$ be the minimum and the maximum degree of vertices in G. The spectral radius r(G) of G is the largest eigenvalue of its adjacency matrix A(G).

Spectra of graphs have been studied in recent years, but the results are often weak when applied to planar graphs. In 1978, A. J. Schwenk and R. J. Wilson [6] asked, in particular, what can be said about the eigenvalues of a planar graph. For ten years after this paper little work was done on this problem. Then in 1988 Hong Yuan [7] proved that the spectral radius of a planar graph on n vertices is less than or equal to $\sqrt{5n-11}$. In [3] Cvetković and Rowlinson conjectured that $K_1 \vee P_{n-1}$, with spectral radius very close to $1 + \sqrt{n}$, is the unique graph with the largest spectral radius among all

LINEAR ALGEBRA AND ITS APPLICATIONS 187: 251-257 (1993)

© Elsevier Science Publishing Co., Inc., 1993 655 Avenue of the Americas, New York, NY 10010

0024-3795/93/\$6.00

251

outerplanar graphs with *n* vertices. In this paper we improve some decomposition results for planar graphs to show that $r(G) < 4 + \sqrt{3(n-3)}$ for any planar graph G on *n* vertices. Moreover, $r(G) < 1 + \sqrt{2 + \sqrt{2}} + \sqrt{n-5}$ for any outerplanar graph on *n* vertices.

2. RESULTS

LEMMA 1 (Courant-Weyl inequality) [4]. Let A and B be two real symmetric matrices of order n. If C = A + B then $r(C) \leq r(A) + r(B)$.

LEMMA 2 (Hong Yuan) [7]. If G is a graph with n vertices, m edges, and no isolated vertices, then

$$r(G) \leqslant \sqrt{2m - n + 1}$$

with equality if and only if G is the disjoint union of either a star or a complete graph with copies of K_2 .

LEMMA 3 (Barnette) [1]. If G is planar and 3-connected, then G has a spanning tree with maximum degree at most 3.

The following lemma is easy to prove.

LEMMA 4. Let T be a tree with at least one vertex of degree 3. Color some vertices of degree 3 red. Then there exists a red vertex v with at least two neighbors that are not adjacent to any other red vertices.

LEMMA 5. A maximal planar graph with at least four vertices has a disjoint edge decomposition into a spanning tree with maximum degree at most 4 and a spanning subgraph with no isolated vertices.

Proof. For a spanning tree T of the graph G let \overline{T} denote the complement of T in G. Color the isolated vertices of \overline{T} red. Consider the set **S** of all spanning trees T of G with the following properties:

- (1) $\Delta(T) \leq 4$.
- (2) All red vertices have degree 3 in G.
- (3) For any vertex v, if $\deg_T(v) = 4$ then v is adjacent to no red vertex.

We first claim that **S** is not empty. Since G is maximal, G is 3-connected, and by Lemma 3, G has a spanning tree T with $\Delta(T) \leq 3$. This verifies conditions (1) and (3). Since G is 3-connected, a red vertex v must satisfy $\deg_T(v) = \deg_G(v) = 3$, verifying condition (2).

SPECTRAL RADIUS OF A PLANAR CRAPH

Let T be a spanning tree in **S** such that \overline{T} has the least number of isolated vertices. If \overline{T} has no isolated vertex, we are done. Otherwise, by Lemma 4 there exists a red vertex u with at least two neighbors that are not adjacent to other red vertices. Construct a new tree T' as follows by edge switching on a single K_4 . Let x, y, z be the three vertices adjacent to u where x and y are not adjacent to other red vertices. Then the edges $ux, uy, uz \in T$ and $xy, yz, zx \in T$. Now let T' = T - ux + xy. Clearly T' is a spanning tree of G. We claim that conditions (1)-(3) hold for T'. Only vertex y has increased its degree in the spanning tree. Since y was adjacent to red vertex u, by condition (3) we have $\deg_{T'}(y) = \deg_T(y) + 1 \leq 3 + 1$ = 4. Hence $\Delta(T') \leq 4$, verifying condition (1). Since $ux, zy \in \overline{T'}$, vertices u, x, y, and z are not isolated in $\overline{T'}$. In particular, vertex u is no longer red in $\overline{T'}$. Hence the set of isolated vertices of $\overline{T'}$ is a proper subset of the set of isolated vertices of \overline{T} . This verifies that condition (2) holds for T', and also condition (3), because vertex y, the only possible vertex of degree 4 in T' not of degree 4 in T, was adjacent to no red vertex besides u.

The fact that the complement of T' in G has one less isolated vertex than the complement of T contradicts the minimality of the number of isolated vertices.

THEOREM 1. If G is a planar graph with $n \ge 3$ vertices, then

$$r(G) < 4 + \sqrt{3(n-3)}.$$

Proof. Suppose $n \ge 4$; otherwise the result is obvious. Let G' be a maximal planar graph containing G as a spanning subgraph. By Lemma 5, G' can be decomposed into a spanning tree T with maximum degree at most 4 and a spanning subgraph H with no isolated vertex. Then A(G') = A(T) + A(H) implies, by Lemma 1, that $r(G') \le r(T) + r(H)$. Since the largest eigenvalue of a graph is less than or equal to the maximum degree with equality if and only if the graph is regular [2], then $r(T) < \Delta(T) \le 4$. Since H has (3n - 6) - (n - 1) = 2n - 5 edges and no isolated vertices, by Lemma 2, $r(H) \le \sqrt{2(2n - 5) - n + 1} = \sqrt{3(n - 3)}$.

It is not difficult to show that $r(\overline{K_2} \vee C_{n-2}) = 1 + \sqrt{2n-3}$, and that $r(P_2 \vee P_{n-2})$ is between $1 + \sqrt{2n-3}$ and $2 + \sqrt{2n-3}$. After examining the spectral radius of several other families of graphs and some small graphs we conjecture that $P_2 \vee P_{n-2}$ and $\overline{K_2} \vee C_{n-2}$ are optimal in the following

sense:

CONJECTURE 1. If G is a planar graph with n vertices, then

 $r(G) \leq r(P_2 \vee P_{n-2}) < 2 + \sqrt{2n-3}.$ Moreover, if $\delta(G) = 4$ then $r(G) \leq r(\overline{K_2} \vee C_{n-2}) = 1 + \sqrt{2n-3}.$

REMARK. To prove the conjecture it is tempting to try to find a spanning subgraph of G with small maximum degree, such that its complement in G has about 3n/2 edges and no isolated vertices. The same argument used in Theorem 1 would then improve the bound of Theorem 1 from order of magnitude $\sqrt{3n}$ to order $\sqrt{2n}$. However, the graph $P_2 \vee P_{n-2}$ allows no such decomposition. Another approach is required if the conjecture is true.

3. OUTERPLANAR GRAPHS

An outerplanar graph G is a graph which can be embedded in the plane so that all vertices are on one face, say the outer face. An *internal triangle* is a triangle with no edge on the outer face. Let U_n $(n \ge 4)$ be the set of all such maximal outerplanar graphs which have n vertices and no internal triangles. Rowlinson [5] proved that $K_1 \lor P_{n-1}$ is the unique graph in U_n with maximal spectral radius. Moreover, he and Cvetković [3] conjectured that $K_1 \lor P_{n-1}$ is the unique graph with maximal spectral radius among all outerplanar graphs with n vertices. It is not difficult to prove that the largest eigenvalue of $K_1 \lor P_{n-1}$ is between $1 + \sqrt{n} - 2/(2 + n - 2\sqrt{n})$ and $1 + \sqrt{n}$. The theorem in this section comes close to confirming the conjecture of Rowlinson and Cvetković.

Clearly, a maximal outerplanar graph can be decomposed into a spanning 2-regular subgraph, the outer face, and its complement in G with exactly n-3 edges. Furthermore, we have the following improvement.

LEMMA 6. A maximal outerplanar graph G has a spanning subgraph H with the following properties:

(1) $\Delta(H) \leq 4$.

(2) The complement of H in G has no isolated vertices.

- (3) H consists of a single cycle together with some pendant edges.
- (4) No two vertices of degree 4 in H are adjacent.
- (5) If H has a vertex with degree 4, then H also has a vertex with degree
- 2.

SPECTRAL RADIUS OF A PLANAR GRAPH

Proof. If the order n of G is less than 7, then Lemma 6 can be routinely checked. Suppose that $n \ge 7$. Let $C = (v_1v_2 \cdots v_n)$ be the Hamiltonian cycle of G on the outer face, and $D = \{v_{k_1}, v_{k_2}, \ldots, v_{k_i}\}$ the set of all vertices of degree 2 in G. It is easy to prove that $|D| \ge 2$ and no two vertices of D are adjacent in G. Let $v_i \in D$. By the maximality of G, v_{i-1} must be adjacent to v_{i+1} , with the convention $v_{n+1} = v_1$. Further, either v_{i+1} or v_{i-1} has degree at least 4. Let F be the subgraph of G obtained from C by adding the set of edges $\{v_{k_1-1}v_{k_1+1}, v_{k_2-1}v_{k_2+1}, \ldots, v_{k_i-1}v_{k_i+1}\}$. Further, let H be the subgraph of G obtained from F by the following procedure. For each $v_{k_i} \in D$ let u and y be the two neighbors of v_{k_i} . There are three cases:

Case 1. If one of the neighbors of v_{k_i} , say u, has degree 3, then delete the edge uv_{k_i} from F.

Case 2. If one of the neighbors of v_{k_i} , say u, has degree 4 and is adjacent to another vertex v in D, then delete the edges uv_{k_i} and uv from F. Since G is maximal, the second neighbor x of v is adjacent to y. Since $n \ge 7$, the degrees of x and y are at least 4; moreover, neither x nor y can assume the role of u, and so neither of the edges xv or yv_{k_i} is deleted from F. (In Figure 1 the dark arcs are in F and the dotted arcs are those deleted from F.)

Case 3. Otherwise delete $v_{k_i}u$ from F, where u is either neighbor of v_{k_i} .

Now the remaining graph H is clearly a spanning subgraph satisfying properties (1), (2), and (3). Concerning property (4), assume vertices x and z are adjacent in H and both have degree 4 in H. Then x and z also have degree 4 in F, which implies that x and z are both adjacent to a vertex y



FIG. 1. Case (2) of Lemma 6.

with degree 2 in *F*. But by the construction, either *xy* or *zy* was deleted from *F* in forming *H*, making it impossible for both *x* and *y* to have degree 4. This is a contradiction. Concerning property (5), note that the pendant vertices of *H* are the vertices of *D*. Let *Q* denote the set of vertices on the cycle of *H* (nonpendant vertices). Since $|D| \leq \frac{1}{2}n$, therefore $|Q| \geq \frac{1}{2}n$. If there are no vertices of degree 2 and at least one vertex of degree 4 in *Q*, then the total number of vertices in *H* is greater than 2(n/2) + 1 = n + 1, a contradiction.

REMARK. Note that if $G = K_1 \vee P_{n-1}$, then G does not have a spanning, unicyclic, connected subgraph with maximum degree 3 such that its complement in G has no isolated vertices. In this sense Lemma 6 cannot be improved.

LEMMA 7. If H is the graph in Lemma 6, then

$$r(H) \leq 1 + \sqrt{2 + \sqrt{2}} < 3.$$

Proof. Let L_5 be the graph in Figure 2. We claim that H can be decomposed into an edge disjoint union of subgraphs I and J where each component of I is isomorphic to K_2 and each component of J is isomorphic to a subgraph of P_5 or L_5 .

To see this let $C = (u_1, u_2, ..., u_s)$ be the cycle in H. If s is even, let I consist of every other edge in C and let J consist of all edges in H not in I. Properties (3) and (4) in Lemma 6 guarantee that the claim is true.

If s is odd, let I' consist of every other edge in C beginning at a vertex of degree 2, if such a vertex exists. Let J' consist of all edges in H not in I'. Then all components of $C \\ I'$ are isomorphic to K_2 except one that is isomorphic to P_3 . Let the three consecutive vertices of the P_3 component mentioned above be x, y, z. If y is adjacent to a vertex v other than x or z in H, then let I = I' + yv and J = J' - yv. Otherwise let I = I' and J = J'. If H has no vertices of degree 4, then each component of J is



FIG. 2. The graph L_5 .

isomorphic to a subgraph of P_5 . If there is a vertex of degree 4 in H, then by property (5) of Lemma 6, the graph H has a vertex with degree 2, which without loss of generality can be chosen to be vertex x. Then each component of J is isomorphic to a subgraph of L_5 . This verifies the claim in the odd case.

Since r(I) = 1 and $r(J) = \max\{r(P_5), r(L_5)\} = r(L_5) = \sqrt{2 + \sqrt{2}}$, by Lemma 1 we have $r(H) \leq r(I) + r(J) \leq 1 + \sqrt{2 + \sqrt{2}} < 3$.

THEOREM 2. If G is an outerplanar graph with n vertices, then

$$r(G) \leq 1 + \sqrt{2 + \sqrt{2}} + \sqrt{n - 5}.$$

Proof. Let G' be a maximal outerplanar graph containing G as a spanning graph. Let H be the unicyclic subgraph of Lemma 6, and F its complement in G'. Since F has no isolated vertices and (2n-3) - n = n - 3 edges, Lemma 2 yields $r(F) \leq \sqrt{2(n-3) - n + 1} = \sqrt{n-5}$. By Lemmas 1 and 7, $r(G) \leq 1 + \sqrt{2 + \sqrt{2}} + \sqrt{n-5}$.

REFERENCES

- 1 D. Barnette, Trees in polyhedral graphs, Canad. J. Math. 18:731-736 (1966).
- 2 D. Cvetković, M. Doob, and H. Sachs, *Spectra of Graphs*, Academic, New York, 1980.
- 3 D. Cvetković and P. Rowlinson, The largest eigenvalue of a graph: A survey, Linear and Multilinear Algebra 28:3-33 (1990).
- 4 A. J. Hoffman, On eigenvalues and colourings of graphs, in *Graph Theory and its* Applications (B. Harris, Ed.), Academic, 1970, pp. 78–91.
- 5 P. Rowlinson, On the index of certain outerplanar graphs, Ars Combin. 29C:221–225 (1990).
- 6 A. J. Schwenk and R. J. Wilson, On the eigenvalues of a graph, in *Selected Topics in Graph Theory* (L. W. Beineke and R. J. Wilson, Eds.), Academic, 1978, pp. 307–336.
- 7 Hong Yuan, A bound on the spectral radius of graphs, *Linear Algebra Appl*. 108:135-139 (1988).

Received 10 March 1992; final manuscript accepted 24 August 1992