

# Algorithmic Randomness of Closed Sets

**George Barmalias**

School of Mathematics, University of Leeds  
Leeds LS2 9JT

**Paul Brodhead, Douglas Cenzer, Seyyed Dashti**

Department of Mathematics, University of Florida  
P.O. Box 118105, Gainesville, Florida 32611  
email: cenzer@math.ufl.edu fax: 352-392-8357

**Rebecca Weber**

Department of Mathematics, Dartmouth College  
Hanover, NH 03755-3551

June 3, 2007

## Abstract

We investigate notions of randomness in the space  $\mathcal{C}[2^{\mathbb{N}}]$  of nonempty closed subsets of  $\{0, 1\}^{\mathbb{N}}$ . A probability measure is given and a version of the Martin-Löf test for randomness is defined.  $\Pi_2^0$  random closed sets exist but there are no random  $\Pi_1^0$  closed sets. It is shown that any random closed set is perfect, has measure 0, and has box dimension  $\log_2 \frac{4}{3}$ . A random closed set has no  $n$ -c.e. elements. A closed subset of  $2^{\mathbb{N}}$  may be defined as the set of infinite paths through a tree and so the problem of compressibility of trees is explored. If  $T_n = T \cap \{0, 1\}^n$ , then for any random closed set  $[T]$  where  $T$  has no dead ends,  $K(T_n) \geq n - O(1)$  but for any  $k$ ,  $K(T_n) \leq 2^{n-k} + O(1)$ , where  $K(\sigma)$  is the prefix-free complexity of  $\sigma \in \{0, 1\}^*$ .

## 1 Introduction

The study of algorithmic randomness has been of great interest in recent years. The basic problem is to quantify the randomness of a single real number; here

---

The authors wish to thank Jack Lutz and Joe Miller for helpful discussions, and the referee for comments that greatly improved the paper. Much of the contents of this paper was discussed during the AIM workshop on Effective Randomness in August, 2006.

A preliminary version of this paper appeared in the Proceedings of CIE 2006 [2]

Research partially supported by National Science Foundation grants DMS 0532644, 0554841 and 0652732.

Keywords: Computability, Randomness,  $\Pi_1^0$  Classes

we will extend this problem to the randomness of the set of paths through a finitely-branching tree. Early in the last century, von Mises [30] suggested that a random real should obey reasonable statistical tests, such as having a roughly equal number of zeroes and ones of the first  $n$  bits, in the limit. Thus a random real would be *stochastic* in modern parlance. If one considers only *computable* tests, then there are countably many and one can construct a real satisfying all tests.

An early approach to randomness was through betting. Effective betting on a random sequence should not allow one's capital to grow unboundedly. The betting strategies used are constructive martingales, introduced by Ville [29] and implicit in the work of Levy [21], which represent fair double-or-nothing gambling.

Martin-Löf [23] observed that stochastic properties could be viewed as special kinds of measure zero sets and defined a random real as one which avoids certain effectively presented measure 0 sets. That is, a real  $x \in 2^{\mathbb{N}}$  is Martin-Löf random if for every effective sequence  $S_1, S_2, \dots$  of c.e. open sets with  $\mu(S_n) \leq 2^{-n}$ ,  $x \notin \bigcap_n S_n$ . It is easy to see that this is equivalent to the condition that we get if we replace  $2^{-n}$  above with  $q_n$  for a computable sequence  $(q_i)$  of rationals such that  $\lim_i q_i = 0$ .

At the same time Kolmogorov [17] defined a notion of randomness for finite strings based on the concept of *incompressibility*. The stronger notion of prefix-free complexity was developed by Levin [20], Gács [16] and Chaitin [9] and extended to infinite words. Schnorr later proved [26] that the notions of constructive martingale randomness, Martin-Löf randomness, and prefix-free randomness are equivalent.

In this paper we want to consider algorithmic randomness on the space  $\mathcal{C}$  of nonempty closed subsets  $P$  of  $2^{\mathbb{N}}$ . Some definitions are needed. Fix a finite alphabet  $A = \{0, 1, \dots, k-1\} = k$ ; we will make use of the alphabets  $\{0, 1\}$  and  $\{0, 1, 2\}$ . For a finite string  $\sigma \in A^n$ , let  $|\sigma| = n$ . Let  $\lambda$  denote the empty string, which has length 0. A word  $(a)$  of length 1 is may be identified with the symbol  $a$ . For two strings  $\sigma, \tau$ , say that  $\tau$  extends  $\sigma$  and write  $\sigma \sqsubseteq \tau$  if  $|\sigma| \leq |\tau|$  and  $\sigma(i) = \tau(i)$  for  $i < |\sigma|$ . Similarly  $\sigma \sqsubset x$  for  $x \in 2^{\mathbb{N}}$  means that  $\sigma(i) = x(i)$  for  $i < |\sigma|$ . Let  $\sigma \hat{\ } \tau$  denote the concatenation of  $\sigma$  and  $\tau$ . Let  $X[n = (x(0), \dots, x(n-1))]$ . Now a nonempty closed set  $P$  may be identified with a tree  $T_P \subseteq A^*$  as follows. For a finite string  $\sigma$ , let  $I(\sigma)$  denote  $\{x \in 2^{\mathbb{N}} : \sigma \sqsubset x\}$ . Then  $T_P = \{\sigma : P \cap I(\sigma) \neq \emptyset\}$ . Note that  $T_P$  has no dead ends, that is if  $\sigma \in T_P$  then either  $\sigma \hat{\ } 0 \in T_P$  or  $\sigma \hat{\ } 1 \in T_P$ .

For an arbitrary tree  $T \subseteq A^*$ , let  $[T]$  denote the set of infinite paths through  $T$ , that is,

$$x \in [T] \iff (\forall n)x[n \in T].$$

It is well-known that  $P \subseteq 2^{\mathbb{N}}$  is a closed set if and only if  $P = [T]$  for some tree  $T$ .  $P$  is a  $\Pi_1^0$  class, or effectively closed set, if  $P = [T]$  for some computable tree  $T$ . Note that if  $P$  is a  $\Pi_1^0$  class, then  $T_P$  is a  $\Pi_1^0$  set, but not in general computable.  $P$  is said to be a *decidable*  $\Pi_1^0$  class if  $T_P$  is computable.  $P$  is said to be a *strong*  $\Pi_2^0$  class, if  $T_P$  is a  $\Pi_2^0$  set, or equivalently if  $P = [T]$  for some  $\Delta_2^0$  tree;

$P$  is said to be a *strong*  $\Delta_2^0$  class if  $T_P$  is  $\Delta_2^0$ . Thus any  $\Pi_1^0$  class is also a strong  $\Delta_2^0$  class. Any decidable  $\Pi_1^0$  class contains a computable element (in particular the leftmost and rightmost paths) and similarly any strong  $\Delta_2^0$  class contains a  $\Delta_2^0$  element. On the other hand, there exist  $\Pi_1^0$  classes with no computable elements and strong  $\Pi_2^0$  classes with no  $\Delta_2^0$  elements. The complement of a  $\Pi_1^0$  class is sometimes called a c.e. open set.

There is a natural effective enumeration  $P_0, P_1, \dots$  of the  $\Pi_1^0$  classes and thus an enumeration of the c.e. open sets. Thus we can say that a sequence  $S_0, S_1, \dots$  of c.e. open sets is *effective* if there is a computable function,  $f$ , such that  $S_n = 2^{\mathbb{N}} - P_{f(n)}$  for all  $n$ . For a detailed development of  $\Pi_1^0$  classes, see [7] or [8].

For background and terminology on computable functions and computably enumerable sets, see [27].

The betting approach to randomness is formalized as follows:

**Definition 1.1 (Ville [29]).** (i) A martingale is a function  $m : k^{<\omega} \rightarrow [0, \infty)$  such that for all  $\sigma \in k^{<\omega}$ ,

$$m(\sigma) = \frac{1}{k} \sum_{i=0}^{k-1} m(\sigma \frown i).$$

(ii) A martingale  $m$  succeeds on  $X \in k^{\mathbb{N}}$  if

$$\limsup_{n \rightarrow \infty} d(X \upharpoonright n) = \infty.$$

That is, the betting strategy results in an unbounded amount of money made on the  $k$ -ary infinite sequence  $X$ .

(iii) The success set of  $m$  is the set  $S^\infty[m]$  of all sequences on which  $m$  succeeds.

That is, a martingale on  $2^{<\omega}$  is the capital function of a fair double-or-nothing betting strategy. When working on  $3^{<\omega}$  the strategy is triple-or-nothing.

**Definition 1.2.** A martingale  $m$  is constructive (effective, c.e.) if it is lower semi-computable; that is, if there is a computable function  $\hat{m} : k^{<\omega} \times \mathbb{N} \rightarrow \mathbb{Q}$  such that

(i) for all  $\sigma$  and  $t$ ,  $\hat{m}(\sigma, t) \leq \hat{m}(\sigma, t+1) < m(\sigma)$ , and

(ii) for all  $\sigma$ ,  $\lim_{t \rightarrow \infty} \hat{m}(\sigma, t) = m(\sigma)$ .

In other words,  $m(w)$  is approximated from below by rationals uniformly in  $w$ . A sequence in  $k^{\mathbb{N}}$  is constructive martingale random if no constructive martingale succeeds on it.

Some flexibility may be gained by also considering *nonmonotonic* martingales; i.e., martingales which bet on the bits of a sequence out of order. While for a monotonic martingale only the *amount* of the next bet is determined from the bits seen previously, for a nonmonotonic martingale both the amount and

the *location* of the next bet are determined from the bits seen previously (the next bit may precede them, follow them, or lie in the middle). These martingales must obey two rules: the standard fair-betting rule that monotonic martingales obey, and the rule that they never bet on the same bit twice. We refer the reader to Downey and Hirschfeldt [11] for the formal definition.

Although a priori allowing nonmonotonic martingales strengthens the notion of randomness, since more strategies must be defeated, in fact in the c.e. case they are equivalent. Muchnik, Semenov, and Uspensky [24] (Theorem 8.9) show that ML-random sequences defeat all *computable* nonmonotonic martingales (in fact they show this with respect to general measures, not just the coin-toss measure). The proof does not depend on the computability of the martingale, however; the martingale is used to define a Martin-Löf test which may be enumerated equally well alongside the enumeration of the martingale. Therefore, as defeating all c.e. nonmonotonic martingales is clearly sufficient to be ML-random, the two are equivalent.

Prefix-free randomness for reals is defined as follows. A Turing machine  $M$  which takes inputs from  $A^*$ , where  $A$  is a finite alphabet, is called prefix-free if it has prefix-free domain  $\text{dom}(M)$ ; that is, if  $\sigma \sqsubseteq \tau$  are strings in  $\text{dom}(M)$ , then  $\sigma$  must equal  $\tau$ . For any finite string  $\tau$ , the *prefix-free complexity of  $\tau$  with respect to  $M$*  is

$$K_M(\tau) = \min\{|\sigma|, \infty : M(\sigma) = \tau\}.$$

There is a *universal* prefix-free function  $U$  such that, for any prefix-free  $M$ , there is a constant  $c$  such that for all  $\tau$

$$K_U(\tau) \leq K_M(\tau) + c.$$

We let  $K(\tau) = K_U(\tau)$  and call it the *prefix-free complexity of  $\tau$* . Then  $x$  is called *prefix-free random* if there is a constant  $c$  such that  $K(x \upharpoonright n) \geq n - c$  for all  $n$ . This means that the initial segments of  $x$  are not *compressible*.

The equivalence of these three notions of randomness (via tests, betting or incompressibility) is a result of Schnorr [26] and is a fundamental result in the theory of algorithmic randomness. While these definitions and results are usually given for binary strings and sequences, they carry over to  $k$ -ary strings and sequences as well. See for example Calude [5, 6]. The following lemma will be needed.

**Lemma 1.3.** *If  $P$  is a  $\Pi_1^0$  class of measure 0, then  $P$  has no random elements.*

*Proof.* Let  $T$  be a computable tree such that  $P = [T]$ , and for each  $n$ , let  $P_n = \bigcup\{I(\sigma) : \sigma \in T \cap \{0, 1\}^n\}$ . Then  $\{P_n\}_{n \in \mathbb{N}}$  is an effective sequence of clopen sets with  $P = \bigcap_n P_n$  and  $\lim_n \mu(P_n) = \mu(P) = 0$ . Furthermore,

$$\mu(P_n) = 2^{-n} |T \cap \{0, 1\}^n|$$

and is therefore a computable sequence. Thus  $\{P_n\}_{n \in \mathbb{N}}$  is a Martin-Löf test, showing that  $P$  has no random elements.  $\square$

We will want to use the following result from the literature [30].

**Theorem 1.4 (Von-Mises–Church–Wald Computable Selection Theorem).** *For any random sequence  $x$  and any computable 1-1 function  $g$ , the sequence  $z(n) = x(g(n))$  is random.*

## 2 Martin-Löf Randomness of Closed Sets

In this section, we define a measure on the space  $\mathcal{C}$  of nonempty closed subsets of  $2^{\mathbb{N}}$  and use this to define the notion of randomness for closed sets. We then obtain several properties of random closed sets.

An effective one-to-one correspondence between the space  $\mathcal{C}$  and the space  $3^{\mathbb{N}}$  is defined as follows. Let a closed set  $Q$  be given and let  $T = T_Q$  be the tree without dead ends such that  $Q = [T]$ .

Define the code  $x = x_Q \in \{0, 1, 2\}^{\mathbb{N}}$  for  $Q$  as follows. Let  $\lambda = \sigma_0, \sigma_1, \sigma_2, \dots$  enumerate the elements of  $T$  in order, first by length and then lexicographically. We now define  $x = x_Q = x_T$  by recursion as follows. For each  $n$ ,  $x(n) = 2$  if  $\sigma_n \frown 0$  and  $\sigma_n \frown 1$  are both in  $T$ ,  $x(n) = 1$  if  $\sigma_n \frown 0 \notin T$  and  $\sigma_n \frown 1 \in T$  and  $x(n) = 0$  if  $\sigma_n \frown 0 \in T$  and  $\sigma_n \frown 1 \notin T$ . For example, if  $Q = \{0, 1\}^{\mathbb{N}}$ , then  $x_Q = (2, 2, \dots)$  and if  $Q = \{y\}$ , then  $x_Q = y$ . Let  $Q_x$  denote the unique closed set  $Q$  such that  $x_Q = x$ .

Now define the measure  $\mu^*$  on  $\mathcal{C}$  by

$$\mu^*(\mathcal{X}) = \mu(\{x_Q : Q \in \mathcal{X}\}).$$

Informally this means that given  $\sigma \in T_Q$ , there is probability  $\frac{1}{3}$  that both  $\sigma \frown 0 \in T_Q$  and  $\sigma \frown 1 \in T_Q$  and, for  $i = 0, 1$ , there is probability  $\frac{1}{3}$  that only  $\sigma \frown i \in T_Q$ . In particular, this means that  $Q \cap I(\sigma) \neq \emptyset$  implies that for  $i = 0, 1$ ,  $Q \cap I(\sigma \frown i) \neq \emptyset$  with probability  $\frac{2}{3}$ .

Let us comment briefly on why some other natural representations were rejected. Suppose first that we simply enumerate all strings in  $\{0, 1\}^*$  as  $\sigma_0, \sigma_1, \dots$  and then represent  $T$  by its characteristic function so that  $x_T(n) = 1 \iff \sigma_n \in T$ . Then in general a code  $x$  might not represent a tree. That is, once we have  $(01) \notin T$  we cannot later decide that  $(011) \in T$ . Suppose then that we allow the empty closed set by using codes  $x \in \{0, 1, 2, 3\}^*$  and modify our original definition as follows. Let  $x(n) = i$  have the same definition as above for  $i \leq 2$  but let  $x(n) = 3$  mean that neither  $\sigma_n \frown 0$  nor  $\sigma_n \frown 1$  is in  $T$ . Informally, this would mean that for  $i = 0, 1$ ,  $\sigma \in T$  implies that  $\sigma \frown i \in T$  with probability  $\frac{1}{2}$ . The advantage here is that we can now represent all trees. But this is also a disadvantage, since for a given closed set  $P$ , there are many different trees  $T$  with  $P = [T]$ . The second problem with this approach is that we would have  $[T] = \emptyset$  with positive probability. We briefly return to this subject in Section 6.

Now we will say that a closed set  $Q$  is (Martin-Löf) random if the code  $x_Q$  is Martin-Löf random. This definition clearly relativizes to any oracle in accordance with the definitions of relative randomness in the Cantor space. Since random reals exist, it follows that random closed sets exist. Furthermore, there are  $\Delta_2^0$  random reals, so we have the following.

**Theorem 2.1.** *There exists a random closed set  $Q$  such that  $T_Q$  is  $\Delta_2^0$ .*  $\square$

Note that if  $T_Q$  is  $\Delta_2^0$ , then  $Q$  must contain  $\Delta_2^0$  elements (in particular the leftmost path). Since there exist strong  $\Pi_2^0$  classes with no  $\Delta_2^0$  elements, there are strong  $\Pi_2^0$  classes  $Q$  such that  $T_Q$  is not  $\Delta_2^0$ .

The following lemma will be needed throughout.

**Lemma 2.2.** *For any  $Q \subseteq 2^{\mathbb{N}}$  which is either closed or open,*

$$\mu^*(\{P : P \subseteq Q\}) \leq \mu(Q).$$

*Proof.* Let  $\mathcal{P}_C(Q)$  denote  $\{P : P \subseteq Q\}$ . We first prove the result for nonempty clopen sets  $U$  in place of  $Q$  by the following induction. Suppose  $U = \bigcup_{\sigma \in S} I(\sigma)$ , where  $S \subseteq \{0, 1\}^n$ . For  $n = 1$ , either  $\mu(U) = 1 = \mu^*(\mathcal{P}_C(U))$  or  $\mu(U) = \frac{1}{2}$  and  $\mu^*(\mathcal{P}_C(U)) = \frac{1}{3}$ . For the induction step, let  $S_i = \{\sigma : i \frown \sigma \in S\}$ , let  $U_i = \bigcup_{\sigma \in S_i} I(\sigma)$ , let  $u_i = \mu(U_i)$  and let  $v_i = \mu^*(\mathcal{P}_C(U_i))$ , for  $i = 0, 1$ . Then considering the three cases in which  $S$  includes both initial branches or just one, we calculate that

$$\mu^*(\mathcal{P}_C(U)) = \frac{1}{3}(v_0 + v_1 + v_0v_1).$$

Thus by induction we have

$$\mu^*(\mathcal{P}_C(U)) \leq \frac{1}{3}(u_0 + u_1 + u_0u_1).$$

Now

$$2u_0u_1 \leq u_0^2 + u_1^2 \leq u_0 + u_1,$$

and therefore

$$\mu^*(\mathcal{P}_C(U)) \leq \frac{1}{3}(u_0 + u_1 + u_0u_1) \leq \frac{1}{2}(u_0 + u_1) = \mu(U).$$

For a closed set  $Q$ , let  $Q = \bigcap_n U_n$ , where  $U_n$  is clopen and  $U_{n+1} \subseteq U_n$  for all  $n$ . Then  $P \subseteq Q$  if and only if  $P \subseteq U_n$  for all  $n$ . Thus

$$\mathcal{P}_C(Q) = \bigcap_n \mathcal{P}_C(U_n),$$

so that

$$\mu^*(\mathcal{P}_C(Q)) = \lim_{n \rightarrow \infty} \mu^*(\mathcal{P}_C(U_n)) \leq \lim_{n \rightarrow \infty} \mu(U_n) = \mu(Q).$$

Finally, for an open set  $Q$ , let  $Q = \bigcup_n U_n$  be the union of an increasing sequence of clopen sets  $U_n$ . Then, by compactness,

$$\mathcal{P}_C(Q) = \bigcup_n \mathcal{P}_C(U_n),$$

so that

$$\mu^*(\mathcal{P}_C(Q)) = \lim_{n \rightarrow \infty} \mu^*(\mathcal{P}_C(U_n)) \leq \lim_{n \rightarrow \infty} \mu(U_n) = \mu(Q).$$

This completes the proof of the lemma.  $\square$

Next we will consider the intersection of a random closed set with an interval  $I(\sigma)$  and the disjoint union of random closed sets.

First recall van Lambalgen's theorem.

**Theorem 2.3 (van Lambalgen [28]).** *The following are equivalent.*

1.  $A \oplus B$  is  $n$ -random.
2.  $A$  is  $n$ -random and  $B$  is  $n$ - $A$ -random.
3.  $B$  is  $n$ -random and  $A$  is  $n$ - $B$ -random.
4.  $A$  is  $n$ - $B$ -random and  $B$  is  $n$ - $A$ -random.

Let us call the coding of a closed set  $Q$  by the nodes of its representative tree with no dead ends the *canonical code* of  $Q$ . We wish now to introduce a second method of coding, the *ghost code*. A ghost code of  $Q$  is an infinite ternary string whose terms correspond to all nodes of  $2^{<\omega}$  in lexicographical order. The terms corresponding to the nodes of  $Q$ 's tree (the "canonical nodes") agree with the corresponding terms in the canonical code; the remaining "ghost nodes" may hold any values. Ghost codes are non-unique, and every closed set has a non-random ghost code (if the closed set itself is random take the code with ghost nodes all equal to zero, say). This method of coding is more convenient for some purposes; for example, we will use it to show that if  $Q_0, Q_1$  are closed sets and  $Q = \{0 \frown x : x \in Q_0\} \cup \{1 \frown x : x \in Q_1\}$ ,  $Q$  is random if and only if the  $Q_i$  are random relative to each other. The utility of the ghost codes rests on the following correspondence.

**Theorem 2.4.** *The canonical code of a closed set  $Q \subseteq 2^{\mathbb{N}}$  is random if and only if  $Q$  has some random ghost code. Furthermore, for any  $y$ , the canonical code  $r$  is  $y$ -random if and only if  $Q$  has a ghost code which is  $y$ -random.*

*Proof.* ( $\Leftarrow$ ) Suppose the canonical code of  $Q$  is nonrandom. Then there is a c.e. martingale  $m$  that succeeds on it. From any initial segment  $\sigma$  of a ghost code  $g$  for  $Q$ , the subsequence  $\hat{\sigma}$  of exactly the canonical nodes of  $\sigma$  is computable. Therefore it is computable whether the bit of  $g$  after  $\sigma$  is canonical or ghost. From  $m$ , define the martingale  $m'$  which bets as follows:

$$m'(\sigma \frown i) = \begin{cases} m(\hat{\sigma} \frown i) & \text{next bit is a canonical node} \\ m'(\sigma) & \text{next bit is a ghost node.} \end{cases}$$

That is,  $m'$  holds its money on ghost nodes and bets identically to  $m$  on canonical nodes. It is clear that  $m'$  succeeds on the ghost code  $g$  and thus  $g$  is nonrandom.

( $\Rightarrow$ ) Now suppose the canonical code  $r$  for  $Q$  is random, and let  $q$  be an infinite ternary string that is random relative to  $r$  (and so by Theorem 2.3  $r \oplus q$  is random). We claim the ghost code  $g$  obtained by using the bits of  $r$  as the canonical nodes and the bits of  $q$  in their original order as the ghost nodes is random. It is clear that  $g$  is a ghost code for  $Q$ .

Suppose  $m$  is a c.e. martingale that bets on  $g$ . From  $m$  it is straightforward to define a nonmonotonic martingale  $m'$  which mimics  $m$ 's bets exactly but performs them on  $r \oplus q$ , succeeding whenever  $m$  succeeds. As  $r$  and  $q$  were chosen to be relatively random, this will show  $g$  is random.

As discussed previously, from  $g \upharpoonright n$  it is computable whether  $g(n)$  will be a ghost node or a canonical node, and which position in  $g$  or  $r$  it occupies in either case. Therefore, assuming the bits seen so far may be assembled into an initial segment  $\sigma$  of  $g$ ,  $m'$  takes the values  $m(\sigma \frown i)$ ,  $i < 3$ , as its bets on the corresponding bit of  $r$  or  $g$ , whichever is appropriate. Having seen that bit, then, it can assemble a  $(|\sigma| + 1)$ -length initial segment of  $g$  and repeat the process. As  $m'$  makes identical bets to  $m$  and has identical outcomes, since it cannot succeed on  $r \oplus g$ ,  $m$  cannot succeed on  $g$  and  $g$  is random.

To relativize ( $\Rightarrow$ ), suppose that  $r$  is  $y$ -random, so that  $r \oplus y$  is random by Van Lambalgen's Theorem 2.3. Then in the proof simply choose  $q$  to be random relative to  $r \oplus y$ , and then  $g$  will be random relative to  $y$ . The other direction relativizes in a straightforward way.  $\square$

The primary purpose of the ghost codes is to remove the dependence on the particular closed set under discussion when interpreting bits of the code as nodes of the tree. This is especially useful when subdividing the tree, as in the following definition.

**Definition 2.5.** *The tree join of closed sets  $P_0$  and  $P_1$  is the closed set*

$$Q = \{0 \frown x : x \in P_0\} \cup \{1 \frown x : x \in P_1\}.$$

*Given ghost codes  $r_0, r_1$  for the  $P_i$ , their tree join  $r_0 \boxplus r_1$  is the code for  $Q$  with the corresponding ghost node values.*

The standard *recursion-theoretic join* is defined by

$$r_0 \oplus r_1 = (r_0(0), r_1(0), r_0(1), r_1(1), \dots).$$

We wish to relate the recursion-theoretic join and the tree join.

**Lemma 2.6.** *Given two ghost codes  $r_0, r_1$ , the tree join  $r_0 \boxplus r_1$  is random if and only if the recursion theoretic join  $r_0 \oplus r_1$  is random.*

*Proof.* It is clear that there is a computable permutation  $\pi$  which uniformly maps any tree join  $r_0 \boxplus r_1$  to the recursion-theoretic join  $r_0 \oplus r_1$ . That is, in  $r_0 \oplus r_1$ , the entries of  $r_0$  and  $r_1$  alternate, whereas  $r_0 \boxplus r_1$  starts with a 2, followed by blocks from  $r_0$  and  $r_1$ , as follows. First  $r_0(0), r_1(0)$ , then  $r_0(1), r_0(2), r_1(1), r_1(2)$ , and continuing with pairs of blocks of size 4, 8 and so on. The result now follows from the Von-Mises–Church–Wald Computable Selection Theorem 1.4.  $\square$

We now obtain the following corollary of Theorems 2.3 and 2.4 and Lemma 2.6.



**Corollary 2.7.** *Suppose  $P_i$ ,  $i = 0, 1$ , are closed sets with canonical codes  $r_i$  and let  $P$  be the tree join of  $P_0, P_1$ . Then  $P$  is random if and only if  $r_0 \oplus r_1$  is random.*

*Proof.* ( $\Leftarrow$ ) Suppose that  $r_0 \oplus r_1$  is random. Then by Theorem 2.3,  $r_0$  and  $r_1$  are mutually relatively random. By Theorem 2.4,  $P_0$  has a ghost code  $g_0$  which is random relative to  $r_1$ , and so also vice-versa, and then  $P_1$  has a ghost code  $g_1$  which is random relative to  $g_0$ . Again by 2.3, the recursion-theoretic join  $g_0 \oplus g_1$  is random, so by Theorem 2.6 the tree join  $g_0 \boxplus g_1$  is also random, and hence  $P$  possesses a random ghost code and is random.

( $\Rightarrow$ ) Suppose now that  $P$  is random, and therefore possesses a random ghost code  $g$ . The code  $g$  may be thought of as a tree join  $g_0 \boxplus g_1$ , which is therefore random, and so by Theorem 2.6,  $g_0 \oplus g_1$  is random. By Theorem 2.3, the individual codes  $g_0, g_1$  are therefore mutually relatively random. Now by the related version of Theorem 2.4,  $r_0$  is random relative to  $g_1$ . But  $r_1$  is computable from  $g_1$  and hence  $r_0$  is random relative to  $r_1$  as well. Similarly,  $r_1$  is  $r_0$ -random and thus, again by 2.3,  $r_0 \oplus r_1$  is random.  $\square$

### 3 Members of Random Closed Sets

For any finite string  $\sigma$  of length  $n$ , the probability that a closed set  $Q$  meets  $I(\sigma)$  is  $(\frac{2}{3})^n$ . For a computable real  $y$ , the sequence  $\{Q : Q \cap I(y \upharpoonright n) \neq \emptyset\}$  thus forms a Martin-Löf test in the space  $\mathcal{C}$  of closed sets, which shows that  $y$  does not belong to any Martin-Löf random closed set. That is, for each  $n$ ,  $\{x : Q_x \cap I(y \upharpoonright n) \neq \emptyset\}$  is a c.e. open set and has measure  $(\frac{2}{3})^n$  in  $\{0, 1, 2\}^{\mathbb{N}}$ , where  $Q_x$  is the closed set with code  $x$ . We omit the details, since we will now prove a stronger result.

For any computable, non-decreasing function  $f$ , we say that a real  $\beta \in \{0, 1\}^{\mathbb{N}}$  is  $f$ -c.e. if there exists a computable approximating function  $\phi$  such that, for all  $i \in \mathbb{N}$ ,

- (i)  $\phi(i, 0) = 0$ ;
- (ii)  $\lim_s \phi(i, s) = \beta(i)$ ;
- (iii)  $\{s : \phi(i, s+1) \neq \phi(i, s)\}$  has cardinality  $\leq f(i)$ .

The reals which are  $f$ -c.e. for some computable function  $f$  are part of the well-known Ershov hierarchy [14, 27].

**Theorem 3.1.** *Suppose that  $f$  is computable and bounded by a polynomial. Then no random closed set has any  $f$ -c.e. paths.*

*Proof.* Let  $f$  be as above,  $\beta$  an  $f$ -c.e. real and  $P$  a closed set containing  $\beta$ . Let  $\phi$  be the  $f$ -approximating function for  $\beta$ . Also let  $M_n \subseteq \{0, 1\}^n$  be the set of different  $\phi$ -approximations to  $\beta \upharpoonright n$  during the stages.

A priori,  $|M_n|$  is exponential. However, for a fixed  $n$ ,  $\beta[n$  can change at most  $\sum_{i<n} f(i)$  times, so  $|M_n|$  is also bounded by a polynomial, i.e. there is  $k \in \mathbb{N}$  such that for almost all  $n$ ,  $|M_n| < n^k$ . Now let

$$S_n = \bigcup_{\sigma \in M_n} \{P \mid P \in \mathcal{C} \ \& \ P \cap I(\sigma) \neq \emptyset\}. \quad (1)$$

Then  $(S_n)$  is a uniformly c.e. sequence of open sets in the space  $\mathcal{C}$  of closed sets of  $2^{\mathbb{N}}$  and for all  $n$ ,  $P \in S_n$ . Also for almost all  $n$ ,

$$\mu^*(S_n) \leq \sum_{\sigma \in M_n} \mu^*(\{P \mid P \in \mathcal{C} \ \& \ P \cap I(\sigma) \neq \emptyset\}) = |M_n| \cdot \left(\frac{2}{3}\right)^n \leq n^k \cdot \left(\frac{2}{3}\right)^n.$$

Since  $\lim_n [n^k \cdot (\frac{2}{3})^n] = 0$  there is a computable subsequence of  $(S_n)$  which is a Martin-Löf test and so  $P$  is not random.  $\square$

For any  $K$ -trivial real  $A$  and any unbounded nondecreasing computable function  $h$ ,  $A$  is  $h$ -c.e. (Nies [25]). Thus it follows from Theorem 3.1 that a random closed set can have no  $K$ -trivial paths. We observe that Theorem 3.1 cannot be extended to  $\omega$ -c.e. in general, because there are left-c.e. (and hence  $\omega$ -c.e.) random reals, and by Theorem 3.9 each of these belongs to a random closed set.

**Theorem 3.2.** *If  $Q$  is a random closed set, then  $Q$  has no isolated elements.*

*Proof.* Let  $Q = [T]$  and suppose by way of contradiction that  $Q$  contains an isolated path  $x$ . Then there is some node  $\sigma \in T$  such that  $Q \cap I(\sigma) = \{x\}$ . For each  $n$ , let

$$S_n = \{P \in \mathcal{C} : |\{\tau \in \{0,1\}^n : P \cap I(\sigma \frown \tau) \neq \emptyset\}| = 1\}.$$

That is,  $P \in S_n$  if and only if the tree  $T_P$  has exactly one extension of  $\sigma$  of length  $n + |\sigma|$ . It follows that

$$|P \cap I(\sigma)| = 1 \iff (\forall n) P \in S_n$$

Now for each  $n$ ,  $S_n$  is a clopen set in  $\mathcal{C}$  and again by induction,  $S_n$  has measure  $(\frac{2}{3})^n$ . Thus the sequence  $S_0, S_1, \dots$  is a Martin-Löf test. It follows that for some  $n$ ,  $Q \notin S_n$ . Thus there are at least two extensions in  $T_Q$  of  $\sigma$  of length  $n + |\sigma|$ , contradicting the assumption that  $x$  was the unique element of  $Q \cap I(\sigma)$ .  $\square$

**Corollary 3.3.** *If  $Q$  is a random closed set, then  $Q$  is perfect and hence has continuum many elements.*  $\square$

**Theorem 3.4.** *Every random closed set contains a random element.*

*Proof.* Suppose that a closed set  $Q$  has no random element and consider the following Martin-Löf test on the space  $\mathcal{C}$ :

$$U_i = \{P \mid P \in \mathcal{C} \ \& \ P \subseteq V_i\}$$

where  $(V_i)$  is a universal Martin-Löf test on the Cantor space. By Lemma 2.2,  $\mu^*(U_i) \leq \mu(V_i) \leq 2^{-i}$  so that  $(U_i)$  is a Martin-Löf test on  $\mathcal{C}$ . But  $Q \in \cap_i U_i$ , so  $Q$  is not random.  $\square$

The previous results might suggest that every element of a random closed set is a random real. However, it turns out that every random closed set contains a non-random real.

We need the following classic result of Chernoff [10] (a version of Bernoulli's *Weak Law of Large Numbers*) here and also for another theorem to follow. See [22] for an exposition.

**Lemma 3.5 (Chernoff).** *Let  $E$  be an event which we will refer to as 'success'. If  $E$  occurs with probability  $p$ , then for any natural numbers  $n$  and any  $\varepsilon$  with  $0 \leq \varepsilon \leq 1$ , the probability that out of  $n$  mutually independent trials, the number of successes differs from  $pn$  by  $> \varepsilon pn$  is  $\leq 2^{-\varepsilon^2 pn/3}$ .*

**Theorem 3.6.** *Not every element of a random closed set is random; in particular, the leftmost and rightmost paths in a random closed set are not random reals.*

*Proof.* We will show that, for a random closed set  $Q$ , the leftmost path is not stochastically random, that is, the asymptotic frequency of 0's is  $\frac{2}{3}$ . Since an effectively random real in  $2^{\mathbb{N}}$  must have asymptotic frequency of  $\frac{1}{2}$  for 0's and 1's, this will suffice to prove that the leftmost path is not random. We define a Martin-Löf test as follows. Fix a rational  $\varepsilon$  such that  $0 < \varepsilon < 1$ . For each  $n$ , let  $S_n$  be the family of closed sets (that is, codes for closed sets) such that the first  $n$  bits of the leftmost path have either  $< \frac{2}{3}(1 - \varepsilon)n$ , or  $> \frac{2}{3}(1 + \varepsilon)n$  occurrences of 0. By the definition of our probability measure, we have

$$\mu^*(S_n) = \sum_{|m - \frac{2}{3}n| > \frac{2}{3}\varepsilon n} \binom{n}{m} \left(\frac{2}{3}\right)^m \left(\frac{1}{3}\right)^{n-m}.$$

It now follows from Chernoff's Lemma 3.5 that

$$\mu^*(S_n) \leq 2e^{-\varepsilon^2 2n/9}.$$

Thus the measures of the test sets  $S_n$  have effective limit zero. It is easy to see that the sequence  $\{S_n\}$  is computably enumerable. For each  $n$ ,  $S_n$  is a clopen set and in fact the union of the finite family of intervals  $I(\sigma)$  in  $\mathcal{C}$  such that  $\sigma$  codes a tree up to level  $n$  in which the leftmost path has either  $< \frac{2}{3}(1 - \varepsilon)n$ , or  $> \frac{2}{3}(1 + \varepsilon)n$  occurrences of 0.

Furthermore,  $S'_n = \bigcup_{p \geq n} S_p$  is also a Martin-Löf test. It follows that for any random closed set  $Q$ , and any  $\varepsilon > 0$ , there is an  $n$  such that for all  $m \geq n$ , the frequency of 0's in the first  $m$  bits of the leftmost path is always within  $\varepsilon$  of  $\frac{2}{3}$ . Thus the leftmost path is not effectively random.  $\square$

Recall that the leftmost and rightmost elements of any strong  $\Delta_2^0$  closed set are  $\Delta_2^0$ . Given Theorems 3.4 and 3.6, we ask: Does a  $\Delta_2^0$  random closed set contain a  $\Delta_2^0$  random path?

**Theorem 3.7.** *Every random strong  $\Delta_2^0$  closed set contains a random  $\Delta_2^0$  real.*

*Proof.* Let  $Q$  be a random strong  $\Delta_2^0$  class. By Theorem 3.4,  $Q$  contains a random real  $x$ . Let  $P$  be a  $\Pi_1^0$  class in the Cantor space which contains only randoms and contains  $x$  (this exists since the class of random reals is an effective union of  $\Pi_1^0$  classes). Note that  $P \cap Q$  is a non-empty strong  $\Delta_2^0$  class and it follows that the leftmost path of  $P \cap Q$  is a  $\Delta_2^0$  real which must be random since it belongs to  $P$ .  $\square$

Note that the above theorem does not combine with the low basis theorem to establish the existence of a low random real in any random strong  $\Delta_2^0$  class. Thus we pose the question of whether for any random closed set  $Q$ , if  $T_Q$  is low, then  $Q$  has a low random element.

Next we want to find a random closed set which does not contain a  $\Delta_2^0$  path. Now it is easy [7, 8] to construct a strong  $\Pi_2^0$  class  $P$  of positive measure which contains no  $\Delta_2^0$  elements; of course  $P$  must contain a random real since it has measure 1. The difficult problem is to construct a *random* strong  $\Pi_2^0$  class with no  $\Delta_2^0$  elements. We have the following result in this direction, which yields a random strong  $\Delta_3^0$  closed set with no  $\Delta_2^0$  elements.

**Theorem 3.8.** *For any set  $A$  there is an  $A$ -random closed set  $Q$  such that  $T_Q \leq_T A''$  but  $Q$  has no elements  $\leq_T A'$ .*

*Proof.* It is enough if we prove the claim for  $A = \emptyset$  because the argument relativises to any oracle  $A$  in a straightforward way. For  $A = \emptyset$  we use a finite injury construction over  $\emptyset'$  to construct  $Q$  with the above properties. In the construction we will  $\emptyset'$ -approximate the canonical code of a tree  $T$  which has no  $\Delta_2^0$  paths. To make sure that the tree  $T$  is random we fix a  $\Pi_1^0$  class  $P$  of positive measure in the space  $3^{\mathbb{N}}$  (where the code for  $T$  lies) which contains only randoms, and we make sure that at every stage our approximation (as a finite ternary string) to  $T$ 's canonical code can be extended to a path in  $P$ . Then by compactness the canonical code of our tree will be in  $P$  and so the tree will be random. The changes in the approximations are motivated by the requirements:

$$R_e : \text{if } \Phi_e^{\emptyset'} \text{ is total then the real it defines is not in } [T].$$

Let  $\alpha_s$  be a finite string approximation of the canonical code  $\alpha$  we are building. We will have  $|\alpha_s| = s$ . Strategy  $R_e$  will come into power after stage  $e$  and will restrain  $\alpha$  up to some  $r_e \geq e$  (the default value is  $r_e[0] = e$ ). Also it might request some changes in  $\alpha$  after the  $e$ -th bit. We start with  $\alpha_0 = \emptyset$  and at stage  $s+1$ , assuming inductively that  $\alpha_s \downarrow$  and  $[\alpha_s] \cap P \neq \emptyset$  we ask for the least  $i < s$  such that  $R_i$  *requires attention*. This happens if

- (i) The longest defined initial segment  $\tau$  of  $\Phi_{i,s+1}^{\emptyset'}$  is larger than ever before;
- (ii) there exists  $\sigma \in \{0, 1, 2\}^*$  such that  $\alpha_s \upharpoonright (\max_{j < i} r_j[s]) \sqsubseteq \sigma$ ,  $I(\sigma) \cap P \neq \emptyset$ ,  $|\sigma| = s+1$ , and  $\tau$  is not consistent with the finite tree with code  $\sigma$ .

If there is no such  $i$  then we extend  $\alpha_s$  by one bit such that  $[\alpha_{s+1}] \cap P \neq \emptyset$ . Otherwise we let  $\alpha_{s+1} = \sigma$  and  $r_i[s+1] = s+1$ . The construction proceeds in a

straightforward way and we can prove inductively that for every  $e$ ,  $R_e$  is satisfied, stops requiring attention and  $r_e$  reaches a limit. Then the limit  $\alpha = \lim_s \alpha_s$  exists and we also have that  $\alpha$  is random by compactness. The satisfaction of the requirements comes from a measure-theoretic fact. Consider  $R_e$  and inductively assume that after stage  $s_e$  no  $R_i$  with  $i < e$  requires attention. Then  $r = \max_{i < e} r_i$  will remain constant. Since  $P$  contains only randoms and  $[\alpha \upharpoonright \max_{i < e} r_i] \cap P \neq \emptyset$ ,

$$\mu([\alpha \upharpoonright r] \cap P) > 0$$

and on the other hand, if  $\beta = \Phi_e^{\theta'}$  we have seen that

$$\mu\{\gamma \mid \gamma \in 3^{\mathbb{N}} \text{ and } \gamma \text{ is the canonical code of a tree which has } \beta \text{ as a path}\} = 0.$$

This means that if at stage  $s_e$  the requirement  $R_e$  is not yet satisfied, it will receive attention at a later stage and get satisfied permanently.  $\square$

As a converse to Theorem 3.4 we have the following.

**Theorem 3.9.** *For any random  $r \in 2^{\mathbb{N}}$ , there exists a random closed set containing  $r$  as a path.*

The proof of this theorem was originally given by Joe Miller and Antonio Montalbán and has been subsequently improved thanks to the anonymous referee.

*Proof.* Let  $r$  be a random real and let  $x$  be the canonical code of an  $r$ -random closed set. We alter  $x$  to the code  $x'$  of a closed set guaranteed to contain  $r$  but changed as little as possible to achieve that.

To determine  $x'(n)$ , assume  $x' \upharpoonright n$  has been defined. If  $x(n) = 2$  or  $x(n)$  corresponds to a node not along  $r$ , set  $x'(n) = x(n)$ . If  $x(n) \in \{0, 1\}$  corresponds to  $r(k)$ , set  $x'(n) = r(k)$ .

The closed set defined by  $x'$  will clearly contain  $r$ . For a contradiction, assume  $x'$  is nonrandom and let  $m'$  be a c.e. martingale that succeeds on it. We build a nonmonotonic martingale  $m$  to bet on  $x \oplus r$ . On bits of  $x$ ,  $m$  will be a triple-or-nothing martingale; on  $r$ , it will be double-or-nothing.

First note that from initial segments of  $x$  and  $r$  we may reconstruct an initial segment of  $x'$  computably, and we always know from an initial segment of  $x'$  whether the next bit is along  $r$  or not, and which bit of  $r$  it is. We will construct  $m$  so that after every stage of betting (which will be one bet by  $m'$  and one or two bets by  $m$ ), the value of  $m$  is equal to the value of  $m'$ . At every stage it will be clear we have revealed enough bits of  $x$  and  $r$  to reconstruct  $x'$  to the needed length.

Suppose inductively  $m$  and  $m'$  hold equal capital after the stage of betting on the last node of  $\sigma \sqsubset x'$ . If the bit  $x'(n)$  following  $\sigma$  is not on  $r$ ,  $m$  bets identically to  $m'$ ; i.e.,  $m(x(n) = i) = m'(\sigma \frown i)$  for  $i < 3$ . In that case  $x(n) = x'(n)$  so our inductive hypothesis holds. If  $x'(n)$  is on  $r$ , set  $m(x(n) = 2) = m'(\sigma \frown 2)$  and for  $i = 0, 1$ , set  $m(x(n) = i) = \frac{1}{2}[m'(\sigma \frown 0) + m'(\sigma \frown 1)]$ . If  $x'(n) = 2$ , then the capital for both  $m$  and  $m'$  is  $m'(\sigma \frown 2)$ , so the inductive hypothesis holds and

we proceed to the next stage. Otherwise  $m$  bets on  $r(k)$  for the appropriate  $k$ , setting  $m(r(k) = i) = m'(\sigma \frown i)$  for  $i = 0, 1$ . On  $r(k)$ , the sum of  $m$ 's capital on each of the two outcomes must average to the previous capital; as the previous capital was  $\frac{1}{2}[m'(\sigma \frown 0) + m'(\sigma \frown 1)]$  this clearly holds. By construction  $r(k) = x'(n) = i$ , so both  $m$  and  $m'$  now have capital  $m'(\sigma \frown i)$  and the inductive hypothesis holds. As  $m'$  is c.e.,  $m$  will also be.

As the values of  $m'$  along  $x'$  are a subsequence of the values of  $m$  along  $x \oplus r$ , if  $m'$  succeeds so does  $m$ , contradicting our assumption on  $x \oplus r$ . Therefore  $x'$  is the code of a random closed set containing the given random path  $r$ .  $\square$

## 4 Measure and Dimension

**Theorem 4.1.** *If  $Q$  is a random closed set, then  $\mu(Q) = 0$ .*

*Proof.* We will show that in the space  $\mathcal{C}$  of closed sets, the  $\mu^*$ -probability that a closed set  $P$  has Lebesgue measure 0, is 1. This is proved by showing that for each  $m$ ,  $\mu(P) \geq 2^{-m}$  with  $\mu^*$ -probability 0. For each  $m$ , let

$$S_m = \{P : \mu(P) \geq 2^{-m}\}.$$

We claim that for each  $m$ ,  $\mu^*(S_m) = 0$ . The proof is by induction on  $m$ .

For  $m = 0$ , we have  $\mu(P) \geq 1$  if and only if  $P = 2^{\mathbb{N}}$ , which is if and only if  $x_P = (2, 2, \dots)$ , so that  $S_0$  is a singleton and thus has measure 0.

Now assume by induction that  $S_m$  has measure 0. Then the probability that a closed set  $P = [T]$  has measure  $\geq 2^{-m-1}$  can be calculated in two parts.

(i) If  $T$  does not branch at the first level, say  $T_0 = \{(0)\}$  without loss of generality. Now consider the closed set  $P_0 = \{y : 0 \frown y \in P\}$ . Then  $\mu(P) \geq 2^{-m-1}$  if and only if  $\mu(P_0) \geq 2^{-m}$ , which has probability 0 by induction, so we can discount this case.

(ii) If  $T$  does branch at the first level, let  $P_i = \{y : i \frown y \in P\}$  for  $i = 0, 1$ . Then  $\mu(P) = \frac{1}{2}(\mu(P_0) + \mu(P_1))$ , so that  $\mu(P) \geq 2^{-m-1}$  implies that at least one of  $\mu(P_i) \geq 2^{-m-1}$ . (Note that the reverse implication is not always true.) Let  $p = \mu^*(S_{m+1})$ . The observations above imply that

$$p \leq \frac{1}{3}(1 - (1 - p)^2) = \frac{2}{3}p - \frac{1}{3}p^2,$$

and therefore  $p = 0$ .

To see that a random closed set  $Q$  must have measure 0, fix  $m$  and let  $S = S_m$ . Then  $S$  is the intersection of an effective sequence of clopen sets  $V_\ell$ , where for  $P = [T]$ ,

$$P \in V_\ell \iff \mu([T_\ell]) \geq 2^{-m}.$$

Since these sets are uniformly clopen, the sequence  $m_\ell = \mu^*(V_\ell)$  is computable. Since  $\lim_\ell m_\ell = 0$ , it follows that this is a Martin-Löf test and therefore no random set  $Q$  belongs to  $\bigcap_\ell V_\ell$ . Then in general, no random set can have measure  $\geq 2^{-m}$  for any  $m$ .  $\square$

Recall that a  $\Pi_1^0$  class  $P$  is decidable if  $T_P$  is decidable. It follows that a nonempty decidable  $\Pi_1^0$  class must contain a computable element (for example, the leftmost path). No computable real can be random and it follows that no decidable  $\Pi_1^0$  class can be random. We will extend this to arbitrary  $\Pi_1^0$  classes in Corollary 4.3 below.

**Theorem 4.2.** *Let  $Q$  be a  $\Pi_1^0$  class with measure 0. Then no subset of  $Q$  is random.*

*Proof.* Let  $T$  be a computable tree (possibly with dead ends) and let  $Q = [T]$ . Then  $Q = \bigcap_n U_n$ , where  $U_n = [T_n]$ . Since  $\mu(Q) = 0$ , it follows from Lemma 2.2 that  $\lim_n \mu^*(\mathcal{P}_C(U_n)) = 0$ . But  $\mathcal{P}_C(U_n)$  is a computable sequence of clopen sets in  $\mathcal{C}$  and  $\mu^*(\mathcal{P}_C(U_n))$  is a computable sequence of rationals with limit 0. Thus  $\mathcal{P}_C(U_n)$  is a Martin-Löf test, so that for any random closed set, there exists  $n$  such that  $P \notin \mathcal{P}_C(U_n)$  and hence  $P$  is not a subset of  $U_n$ .  $\square$

Since any random class has measure 0, we have the following immediate corollary.

**Corollary 4.3.** *No  $\Pi_1^0$  class can be random.*  $\square$

Surprisingly, we can compute the (Kolmogorov) box dimension of a random closed set, and in fact it turns out that all random closed sets have the same dimension. The intuition for this comes from the following lemma. For any function  $F$  mapping the space  $\mathcal{C}$  of closed sets into  $\mathfrak{R}$ , the *expected value* of  $F$  on  $\mathcal{C}$  is the integral  $\int F(P)$  with respect to the probability measure  $\mu^*$ .

**Lemma 4.4.** *In the space  $\mathcal{C}$  of closed sets, the expected cardinality of  $\{\sigma \in \{0, 1\}^n : Q \cap I(\sigma) \neq \emptyset\}$  is exactly  $(\frac{4}{3})^n$  for every  $n$ , where  $Q$  is chosen uniformly at random according to  $\mu^*$ .*

*Proof.* Let  $S_n = \{\sigma \in \{0, 1\}^n : Q \cap I(\sigma) \neq \emptyset\}$ , for a randomly chosen  $Q$  from  $\mathcal{C}$ .

The proof is by induction on  $n$ . For  $n = 1$ , we have two cases. With probability  $\frac{2}{3}$ ,  $\text{card}(S_1) = 1$  and with probability  $\frac{1}{3}$ ,  $\text{card}(S_1) = 2$ . Thus the expected value is exactly  $\frac{4}{3}$ . For  $n + 1$ , there are again two cases. With probability  $\frac{2}{3}$ ,  $\text{card}(S_1) = 1$ , so that the expected  $\text{card}(S_{n+1})$  equals the expected  $\text{card}(S_n)$ , which is  $(\frac{4}{3})^n$  by induction. With probability  $\frac{1}{3}$ ,  $\text{card}(S_1) = 2$ , in which case the expected  $\text{card}(S_{n+1})$  is twice the expected  $\text{card}(S_n)$ , that is,  $2(\frac{4}{3})^n$ . Thus we have the expected value

$$\text{card}(S_{n+1}) = \frac{2}{3} \left(\frac{4}{3}\right)^n + \frac{1}{3} \cdot 2 \left(\frac{4}{3}\right)^n = \left(\frac{4}{3}\right)^{n+1}.$$

$\square$

The box dimension of a closed set in the Cantor space, if it exists, is given by the following limit:

$$\dim_B F(Q) = \lim_{n \rightarrow \infty} \frac{\log_2(\text{card}(T_Q \cap \{0, 1\}^n))}{n}.$$

(See [1] for this formulation of the box dimension in  $\{0, 1\}^{\mathbb{N}}$ .) Now by Lemma 4.4, the expected value of  $\text{card}(T_Q \cap \{0, 1\}^n)$  for a random closed set  $Q$  is  $(\frac{4}{3})^n$ , which suggests that the box dimension of  $Q$  should be  $\log_2 \frac{4}{3}$ .

**Lemma 4.5.** *Let  $Q$  be a random closed set. Then for any  $\varepsilon > 0$ , there exists a  $m \in \mathbb{N}$  such that, for all  $n > m$ ,  $(\frac{4}{3})^n(1-\varepsilon)^n < \text{card}(T_Q \cap \{0, 1\}^n) < (\frac{4}{3})^n(1+\varepsilon)^n$ .*

*Proof.* For each  $n$ , let  $c_n(Q)$ , or just  $c_n$ , denote  $\text{card}(T_Q \cap \{0, 1\}^n)$ . We will use three applications of Chernoff's Lemma 3.5. First we show that there exists  $m$  such that for all  $n > m$ ,  $c_{6n} \geq n$ . Since the tree  $T_Q \cap \{0, 1\}^{\leq 6n-1}$  has at least  $6n$  nodes, it follows from Chernoff's Lemma that the number of branching nodes is less than  $n$  with probability  $\leq 2^{-n/6}$ . Thus  $c_{6n} < n$  with probability  $< 2^{-n/6}$ . Then the probability that  $c_{6n} < n$  for any  $n \geq m$  is less than

$$\sum_{n=m}^{\infty} 2^{-n/6} = \frac{2^{-m/6}}{1 - 2^{-1/6}}.$$

This provides a computable sequence of clopen sets with measures bounded by a computable sequence with limit zero and hence a Martin-Löf test. It follows that for any random closed set  $Q$ , there exists  $m_0$  such that  $c_{6n} \geq n$  for all  $n \geq m_0$ . Now for  $n > m_0$ , there are at least  $6n^2$  nodes in  $T_Q \cap \{0, 1\}^{\leq 12n-1} - \{0, 1\}^{\leq 6n-1}$ , so that again by Chernoff's Lemma, the probability that  $< n^2$  of these are branching nodes is  $\leq 2^{-n^2/6}$ . It follows as above that there exists  $m_1 > 3$  such that  $c_{12n} \geq n^2$  for all  $n \geq m_1$ . Now suppose that  $m \geq 12m_1$  and that  $12n \leq m < 12(n+1) < 16n$ . Then  $n \geq m_1$ , so that

$$c_m \geq c_{12n} \geq n^2 > (m/16)^2.$$

Again by Chernoff's Lemma, the probability that the number of branching nodes from  $T_Q \cap \{0, 1\}^n$  differs from  $\frac{1}{3}c_n$  by  $> \frac{1}{3}c_n^{-\frac{1}{4}}c_n$  is  $< 2^{-\sqrt{c_n}/9}$ . But this is exactly the probability that  $c_{n+1}$  differs from  $\frac{4}{3}c_n$  by  $> \frac{1}{3}c_n^{-\frac{1}{4}}c_n$ . For  $n > m_1$ , we know that  $c_n \geq (\frac{n}{16})^2$ , so that  $\sqrt{c_n} \geq \frac{n}{16}$  and  $c_n^{-\frac{1}{4}} \leq \frac{4}{\sqrt{n}}$  and hence  $2^{-\sqrt{c_n}/9} \leq 2^{-n/144}$ . Thus the probability  $p_n$  that  $c_{n+1}$  differs from  $\frac{4}{3}c_n$  by more than  $\frac{c_n}{9\sqrt{n}}$  is  $< 2^{-n/144}$ . Then the probability that for any  $n \geq m_1$ ,  $c_{n+1}$  differs from  $\frac{4}{3}c_n$  by more than  $\frac{4}{3\sqrt{n}}c_n$  is bounded by

$$\sum_{n=m}^{\infty} p_n = \sum_{n=m}^{\infty} 2^{-n/144} = \frac{2^{-m/144}}{1 - 2^{-1/144}}.$$

This again provides a Martin-Löf test which shows that for any random closed set  $Q$ , there exists  $m_2$  so that for  $n > m_2$ ,

$$(*) \quad \frac{4}{3} \left(1 - \frac{1}{\sqrt{n}}\right) c_n \leq c_{n+1} \leq \frac{4}{3} \left(1 + \frac{1}{\sqrt{n}}\right) c_n.$$

Now given  $\varepsilon$ , choose  $m \geq m_2$  so that  $(1 + \frac{1}{\sqrt{m}})^2 < 1 + \varepsilon$  and  $1 - \varepsilon < (1 - \frac{1}{\sqrt{m}})^2$ .



Then for any  $k$ ,

$$\begin{aligned} c_m \left(\frac{4}{3}\right)^{2k} (1 - \epsilon)^k &< c_m \left(\frac{4}{3}\right)^{2k} \left(1 - \frac{1}{\sqrt{m}}\right)^{2k} < c_{m+2k} \\ &< c_m \left(\frac{4}{3}\right)^{2k} \left(1 + \frac{1}{\sqrt{m}}\right)^{2k} < c_m \left(\frac{4}{3}\right)^{2k} (1 + \epsilon)^k. \end{aligned}$$

Now let  $k$  be large enough so that

$$(1 - \epsilon)^{m+k} \leq c_m \leq \left(\frac{4}{3}\right)^m (1 + \epsilon)^{m+k}.$$

Then the desired inequality

$$\left(\frac{4}{3}\right)^n (1 - \epsilon)^n < c_n < \left(\frac{4}{3}\right)^n (1 + \epsilon)^n.$$

will hold for even  $n \geq m + 2k$ . For odd  $n$ , this inequality will hold by the inequality (\*) above.  $\square$

**Theorem 4.6.** *For any random closed set  $Q$ , the box dimension of  $Q$  is  $\log_2 \frac{4}{3}$ .*

*Proof.* Given  $\epsilon > 0$ , let  $m$  be given by Lemma 4.5. Then for  $n > m$ , we have

$$n \log_2 \frac{4}{3} + n \log_2 (1 - \epsilon) \leq \log_2 (\text{card}(T_Q \cap \{0, 1\}^n)) \leq n \log_2 \frac{4}{3} + n \log_2 (1 + \epsilon),$$

so that

$$\log_2 \frac{4}{3} + \log(1 - \epsilon) \leq \frac{\log_2 (\text{card}(T_Q \cap \{0, 1\}^n))}{n} \leq \log_2 \frac{4}{3} + \log_2 (1 + \epsilon),$$

and therefore  $\dim_B(Q) = \lim_n \frac{\log_2 (\text{card}(T_Q \cap \{0, 1\}^n))}{n} = \log_2 \frac{4}{3}$ .  $\square$

## 5 Prefix-Free Complexity of Closed Sets

In this section, we consider randomness for closed sets in terms of incompressibility of trees. Of course, Schnorr's theorem tells us that  $P$  is random if and only if the code  $x_P \in \{0, 1, 2\}^{\mathbb{N}}$  for  $P$  is prefix-free random, that is,  $K_3(x_P \upharpoonright n) \geq n - O(1)$ . (Schnorr's theorem for arbitrary finite alphabets is shown in [6].) Here we write  $K_3$  to indicate that we would be using a universal prefix-free function  $U : \{0, 1, 2\}^* \rightarrow \{0, 1, 2\}^*$ . However, many properties of trees and closed sets depend on the levels  $T_n = T \cap \{0, 1\}^n$  of the tree. For example, if  $[T_n] = \cup\{I(\sigma) : \sigma \in T_n\}$ , then  $[T] = \bigcap_n [T_n]$  and  $\mu([T]) = \lim_{n \rightarrow \infty} \mu([T_n])$ .

So we want to consider the compressibility of a tree in terms of  $K(T_n)$ . Now there is a natural representation of  $T_n$  as a string of length  $2^n$ . That is, list  $\{0, 1\}^n$  in lexicographic order as  $\sigma_1, \dots, \sigma_{2^n}$  and represent  $T_n$  by the string  $e_1, \dots, e_{2^n}$  where  $e_i = 1$  if  $\sigma_i \in T$  and  $e_i = 0$  otherwise. Henceforth we identify  $T_n$  with this natural representation.

It is interesting to note that the code for  $T_n$  will have a shorter length than the natural representation. For example, if  $[T] = \{y\}$  is a singleton, then  $x = y$  and for each  $n$ , the code for  $T_n$  is  $x \upharpoonright n$ . If  $x$  is the code for the full tree  $\{0, 1\}^*$ , then  $x = (2, 2, \dots)$  and the code for  $T_n$  is a string of  $(2^n - 1)$  2's, those labels attached to nodes of length  $< n$ . For the remainder of this section, we will use  $T_n$  to mean the natural representation and  $x_n$  to mean the code.

One question here is whether there is a formulation of randomness in terms of the incompressibility of  $T_n$ . We will give some partial answers. It seems plausible that  $P = [T]$  is random if and only if there is a constant  $c$  such that  $K(T_n) \geq 2^n - c$  for all  $n$ . We will see that this is not possible for any tree. First we give a lower bound for the prefix-free complexity of a random tree.

**Theorem 5.1.** *If  $P$  is a random closed set and  $T = T_P$ , then there is a constant  $c$  such that  $K(T_n) \geq \left(\frac{7}{6}\right)^n - c$  for all  $n$ .*

*Proof.* Let  $P = [T]$  be a random closed set. Let  $m$  be given by Lemma 4.5, for  $\varepsilon = \frac{7}{6}$ , so that for  $n > m$ ,

$$\text{card}(T_n) \geq \left(\frac{7}{6}\right)^n.$$

It follows that the code  $x_n$  for  $T_n$  has length  $\geq \left(\frac{7}{6}\right)^n$ . Since  $x$  is random, we know that, for  $n \geq m$ ,

$$K_3(x_n) \geq \left(\frac{7}{6}\right)^n - a,$$

for some constant  $a$ . Now we can compute  $x_n$  from  $T_n$ , so that

$$K(T_n) \geq K_3(x_n) - b,$$

for some constant  $b$ . The result now follows.

That is, let  $U$  (mapping  $\{0, 1\}^*$  to  $\{0, 1\}^*$ ) be a universal prefix-free Turing machine and let  $K(T_n) = \min\{|\sigma| : U(\sigma) = T_n\}$ . Let  $M$  be a prefix-free machine  $M$  (mapping  $\{0, 1\}^*$  to  $\{0, 1, 2\}^*$ ) such that  $M(T_n) = x_n$ . Then define  $V$  by

$$V(\sigma) = M(U(\sigma)).$$

Then  $K_V(x \upharpoonright n) \leq K(T_n)$ , so that for some constant  $e$ ,  $K_3(x_n) \leq K(T_n) + e$  and hence

$$K(T_n) \geq K_3(x_n) - e \geq \left(\frac{7}{6}\right)^n - b - e.$$

□

Going in the other direction, we can compute  $T_n$  uniformly from  $x \upharpoonright 2^n$ , so that as above,  $K_3(x \upharpoonright 2^n) \geq K(T_n) - b$  for some  $b$ . Thus in order to conclude that  $P$  is random, we would need to know that  $K(T_n) \geq 2^n - c$  for some  $c$ . The next result shows that this is not possible, since trees are naturally compressible.

**Theorem 5.2.** *For any tree  $T \subseteq \{0, 1\}^*$ , there are constants  $k > 0$  and  $c$  such that  $K(T_\ell) \leq 2^\ell - 2^{\ell-k} + c$  for all  $\ell$ .*

*Proof.* For the full tree  $\{0,1\}^*$ , this is clear so suppose that  $\sigma \notin T$  for some  $\sigma \in \{0,1\}^m$ . Then for any level  $\ell > m$ , there are  $2^{\ell-m}$  possible nodes for  $T$  which extend  $\sigma$  and  $T_\ell$  may be uniformly computed from  $\sigma$  and from the characteristic function of  $T_\ell$  restricted to the remaining set of nodes. That is, fix  $\sigma$  of length  $m$  and define a prefix-free computer  $M$  as follows. The domain of  $M$  is strings of the form  $0^\ell 1 \tau$  where  $|\tau| = 2^\ell - 2^{\ell-m}$ .  $M$  outputs the standard representation of a tree  $T_\ell$  such that no extension of  $\sigma$  is in  $T_\ell$  and such that  $\tau$  tells us whether strings not extending  $\sigma$  are in  $T_\ell$ . It is clear that  $M$  is prefix-free and we have  $K_M(T_\ell) = \ell + 1 + 2^\ell - 2^{\ell-m}$ . Thus  $K(T_\ell) \leq \ell + 1 + 2^\ell - 2^{\ell-m} + c$  for some constant  $c$ . Now  $\ell + 1 < 2^{\ell-m-1}$  for sufficiently large  $\ell$  and thus by adjusting the constant  $c$ , we can obtain  $c'$  so that

$$K(T_\ell) \leq 2^\ell - 2^{\ell-m-1} + c'.$$

□

We might next conjecture that  $K(T_\ell) > 2^{\ell-c}$  is the right notion of prefix-free randomness. However, classes with small measure are more compressible.

**Theorem 5.3.** *If  $\mu([T]) < 2^{-k}$ , then there exists  $c$  such that, for all  $\ell$ ,*

$$K(T_\ell) \leq 2^{\ell-k+1} + c.$$

*Proof.* Suppose that  $\mu([T]) < 2^{-k}$ . Then for some level  $n$ ,  $T_n$  has  $< 2^{n-k}$  nodes  $\sigma_1, \dots, \sigma_t$ . Now for any  $\ell > n$ ,  $T_\ell$  can be computed from the fixed list  $\sigma_1, \dots, \sigma_t$  and the list of nodes of  $T_\ell$  taken from the at most  $2^{\ell-k}$  extensions of  $\sigma_1, \dots, \sigma_t$ . It follows as in the proof of Theorem 5.2 above that for some constant  $c$  and all  $\ell$ ,

$$K(T_\ell) \leq 2^{\ell-k} + \ell + 1 + c.$$

Thus for large enough so that  $\ell + 1 \leq 2^{\ell-k}$ , we have

$$K(T_\ell) \leq 2^{\ell-k+1} + c,$$

as desired. □

Note that if  $\mu([T]) = 0$ , then for any  $k$ , there is a constant  $c$  such that  $K(T_\ell) \leq 2^{\ell-k} + c$ . But by Theorem 4, random closed sets have measure zero. Thus if  $P$  is random, then it is not the case that  $K(T_n) \geq 2^{n-k}$ .

Finally, we will construct an effectively closed set with not too much compressibility. The standard example of a random real, Chaitin's  $\Omega$  [9], is a c.e. real and therefore  $\Delta_2^0$ . Thus there exists a  $\Delta_2^0$  random tree  $T$  and by Theorem 5.1,  $K(T_\ell) \geq (\frac{7}{6})^\ell - c$  for some  $c$ . We have a more modest result for  $\Pi_1^0$  classes.

**Theorem 5.4.** *There is a  $\Pi_1^0$  class  $P = [T]$  such that  $K(T_n) \geq n$  for all  $n$ .*

*Proof.* Recall the universal prefix-free machine  $U$  and let  $S = \{\sigma \in \text{Dom}(U) : |U(\sigma)| \geq 2^{|\sigma|}\}$ . Then  $S$  is a c.e. set and can be enumerated as  $\sigma_1, \sigma_2, \dots$ . The tree  $T = \bigcap_s T^s$  where  $T^s$  is defined at stage  $s$ . Initially we have  $T^0 = \{0,1\}^*$ .

We say that  $\sigma_t$  *requires attention* at stage  $s \geq t$  when  $\tau = U(\sigma_t) = T_n^s$  for some  $n$  (so that  $|\tau| = 2^n$ ) and  $n \geq |\sigma_t|$ . Action is taken by selecting some path  $\rho_t \in T_s$  of length  $n$  and defining  $T^{s+1}$  to contain all nodes of  $T^s$  which do not extend  $\rho_t$ . Then  $\tau \neq T_n^{s+1}$  and furthermore  $\tau \neq T_n^r$  for any  $r \geq s+1$  since future action will only remove more nodes from  $T_n$ .

At stage  $s+1$ , look for the least  $t \leq s+1$  such that  $\sigma_t$  requires action and take the action described if there is such a  $t$ . Otherwise, let  $T^{s+1} = T^s$ .

Let  $A$  be the set of  $t$  such that action is ever taken on  $\sigma_t$ . Recall from the Kraft Inequality that  $\sum_t 2^{-|\sigma_t|} < 1$ . Since  $|\rho_t| \geq |\sigma_t|$ , it follows that  $\sum_{t \in A} 2^{-|\rho_t|} < 1$  as well. Now  $\mu([T]) = 1 - \sum_t 2^{-|\rho_t|} > 0$  and therefore  $[T]$  is nonempty.

It follows from the construction that for each  $t$ , action is taken for  $\sigma_t$  at most once.

Now suppose by way of contradiction that  $U(\sigma) = T_n$  for some  $\sigma_t$  with  $|\sigma| \leq n$ . There must be some stage  $r \geq t$  such that for all  $s \geq r$ ,  $T_n^s = T_n$  and such that action is never taken on any  $t' < t$  after stage  $r$ . Then  $\sigma_t$  will require action at stage  $r+1$  which makes  $T_n^{r+1} \neq T_n^r$ , a contradiction.  $\square$

## 6 Conclusions and Future Research

In this paper we have proposed a notion of randomness for closed sets and derived several interesting properties of random closed sets. Random strong  $\Pi_2^0$  classes exist but no  $\Pi_1^0$  class is random. A random closed set has measure zero and box dimension  $\log_2 \frac{4}{3}$ ; it is perfect and hence uncountable. Results on members of random closed sets include the following. A random closed set contains no  $f$ -c.e. elements, if  $f$  is polynomially bounded. Every random closed set  $Q$  contains a random real, not every element of a random closed set is random and every random real belongs to some random closed set. Furthermore, if  $Q$  is strong  $\Delta_2^0$ , then it contains a random  $\Delta_2^0$  real and if  $T_Q$  is low, then  $Q$  contains a low random element. On the other hand we do not know the answer to the following.

**Problem 6.1.** *Does every random closed set with  $\Delta_2^0$  canonical code contain a low random element?*

We conjecture a negative answer. It is a well known fact that every real is computed by a random real. The corresponding question for trees is as follows.

**Problem 6.2.** *Let  $A$  be an incomputable set. Is there a random closed set such that all of its elements compute  $A$ ?*

We have examined the notion of compressibility for trees based on the prefix-free complexity of the  $n$ th level  $T_n$  of a tree. We showed that for any random closed set (and hence for some strong  $\Pi_2^0$  class), there exists  $c$  such that  $K(T_n) \geq (\frac{7}{6})^n - c$  for all  $n$ . We constructed a  $\Pi_1^0$  class  $P = [T]$  such that  $K(T_n) \geq n$  for all  $n$ . It seems a reasonable conjecture that if  $K(T_n) \geq (\frac{4}{3})^n - c$  for all  $n$ ,

then the closed set  $[T]$  is random. We would like to explore the notion that  $\Pi_1^0$  classes are more compressible than arbitrary closed sets.

Other notions of randomness might also be considered. A general probability measure  $\nu_f$  may be defined on  $3^{\mathbb{N}}$  from a function  $f : \{0, 1, 2\}^* \rightarrow [0, 1]$  such that  $\sum_{i=0,1,2} f(\sigma \frown i) = 1$  for all  $\sigma$ . The interval  $I(\sigma)$  then has  $\nu_f$ -measure  $\prod_{n < |\sigma|} f(\sigma \upharpoonright (n+1))$ . We will say that  $\nu_f$  is a *computable* measure if  $f$  is computable. The probability measure  $\nu$  is *nonatomic* if for any  $x \in 3^{\mathbb{N}}$ ,  $\nu(\{x\}) = 0$ . The function  $f$  (and the corresponding measure  $\nu_f$ ) is *bounded* if there is an upper bound  $b < 1$  such that  $f(\sigma) < b$  for all  $\sigma \in \{0, 1, 2\}^*$ . It is easy to see that any bounded measure is nonatomic. If there exist constants  $b_0, b_1, b_2$  strictly between 0 and 1, such that for all  $\sigma$ ,  $f(\sigma \frown i) = b_i$ , then we will say that  $\nu_f$  is *regular*. For any regular measure, we can define the notion of a  $\nu$ -Martin-Löf test and the resulting notion of a  $\nu$ -Martin-Löf-random (or just  $\nu$ -random) real. It is easy to see that  $\nu$ -random reals exist for any  $\nu$  and hence  $\nu$ -random closed sets exist. The results on ghost codes and joins will hold for any regular measure. The corresponding version of Lemma 2.2 will hold if  $\nu$  is regular with  $b_0$  and  $b_1 \leq \frac{1}{2}$ . The proofs of Theorem 4.2 and Corollary 4.3, that no subset of a measure-zero  $\Pi_1^0$  class is random, also go through under this assumption.

Some of the results in this paper may also be obtained for  $\nu_f$  where  $f(\sigma \frown i) \leq \frac{1}{2}$  for  $i = 0, 1$ . For example with respect to  $\nu_f$  a random closed set will have no isolated elements and it will always contain a random element. For any regular measure, either the leftmost or the rightmost path will be nonrandom, since either  $b_0 + b_2 > \frac{1}{2}$  or  $b_1 + b_2 > \frac{1}{2}$ . The proof of Theorem 3.2 that every random closed set has measure 0 seems to require, for  $\nu_f$ -randomness, that  $f(\sigma \frown 2) \leq \frac{1}{2}$  for all  $\sigma$ .

Returning to the notion of randomness which allows trees with dead ends, let  $b_3$  now be the probability that a given node has no extensions and let the probability be regular as above. Then a simple recursion shows the probability  $p$  of a given closed set being empty satisfies the equation

$$p = b_3 + (b_0 + b_1)p + b_2p^2.$$

Solving for  $p$ , we obtain

$$(p - 1)(b_2p - b_3) = 0.$$

Thus either  $p = 1$  or  $p = \frac{b_3}{b_2}$ . It follows that if  $b_2 \leq b_3$ , then  $p = 1$ , that is, almost every closed set is empty. Suppose now that  $b_3 < b_2$  and let  $p_n$  be the probability that a given tree  $T$  has no paths of length  $n$ . Then it can be seen by induction that  $p_n \leq \frac{b_3}{b_2}$  for all  $n$ . That is,  $p_1 = b_3 \leq \frac{b_3}{b_2}$  and then

$$p_{n+1} = b_3 + (1 - b_2 - b_3)p_n + b_2p_n^2 \leq \frac{b_3}{b_2}.$$

Hence in this case, the probability that a given closed set is empty is  $\frac{b_3}{b_2} < 1$ . In this case, one could presumably develop a notion of a random tree and a random closed set and explore the properties of random closed sets.

A real  $x$  is said to be *K-trivial* if  $K(x \upharpoonright n) \leq K(n) + c$  for some  $c$ . Much interesting work has been done on the *K-trivial* reals. Chaitin showed that if

$A$  is  $K$ -trivial, then  $A \leq_T \mathbf{0}'$ . Solovay constructed a noncomputable  $K$ -trivial real. Downey, Hirschfeldt, Nies and Stephan [12] showed that no  $K$ -trivial real is c.e. complete. The notion of a  $K$ -trivial closed set was introduced in [4]. It was shown in particular that every  $K$ -trivial class contains a  $K$ -trivial member, but there exist  $K$ -trivial  $\Pi_1^0$  classes with no computable members.

The related notion of a random continuous function was introduced in [3]. It was shown that a random continuous function  $F$  on  $2^{\mathbb{N}}$  cannot be computable, so that the graph of  $F$  cannot be  $\Pi_1^0$  class. For any random  $F$  and computable  $x$ ,  $F(x)$  is a random real, however the image of  $F$  need not be a random closed set. The authors can now show that the set of zeroes of a random continuous function is a random closed set. Random Brownian motions have been studied by Fouché [15] and are a special case of random continuous functions on the real line, which is another area of interest for further research.

## References

- [1] K. B. Athreya, J. M. Hitchcock, J. Lutz and E. Mayordomo, *Effective strong dimension, algorithmic information and computational complexity*, SIAM Journal of Computing, to appear.
- [2] P. Brodhead, D. Cenzer and S. Dashti, *Random closed sets*, in Logical Approaches to Computational Barriers, Proc. CIE 2006, A. Beckmann, U. Berger, B. Loewe, and J. Tucker (eds.), Springer Lecture Notes in Computer Science, Vol. 3988 (2006), 55-64.
- [3] P. Brodhead, D. Cenzer and J. B. Remmel, *Random continuous functions*, in Proceedings CCA 2006, Springer Electronic Notes in Computer Science 167 (2007), 275-287.
- [4] G. Barmpalias, D. Cenzer, J. B. Remmel and R. Weber,  *$K$ -trivial closed sets and continuous functions*, in Proceedings of CIE 2007, S. B. Cooper, B. Loewe and A. Sorbi (eds.), Springer Lecture Notes in Computer Science, to appear.
- [5] C. Calude, *Theories of Computational Complexity*, Annals of Discrete Mathematics **35**, North-Holland (1988).
- [6] C. Calude, *Information and Randomness: An Algorithmic Perspective*, Springer-Verlag (1994).
- [7] D. Cenzer and J. B. Remmel,  $\Pi_1^0$  Classes, ASL Lecture Notes in Logic, to appear.
- [8] D. Cenzer and J. B. Remmel,  $\Pi_1^0$  classes, in Handbook of Recursive Mathematics, Vol. 2: Recursive Algebra, Analysis and Combinatorics, editors Y. Ersov, S. Goncharov, V. Marek, A. Nerode, J. Remmel, Elsevier Studies in Logic and the Foundations of Mathematics, Vol. 139 (1998) 623-821.

- [9] G. Chaitin, *Information-theoretical characterizations of recursive infinite strings*, Theoretical Computer Science **2** (1976), 45-48.
- [10] H. Chernoff, *A measure of asymptotic efficiency for tests of a hypothesis based on the sums of observations*, Annals of Mathematical Statistics **23** (1952), 493-509.
- [11] R. Downey and D. Hirschfeldt, *Algorithmic Randomness and Complexity*, in preparation. The current draft is available online at <http://www.mcs.vuw.ac.nz/~downey/> )
- [12] R. Downey, D. Hirschfeldt, A. Nies and F. Stephan, *Trivial reals*, in Proceedings of the 7th and 8th Asian Logic Conferences, World Scientific Press, Singapore (2003), 101-131.
- [13] Rod Downey and Liang Yu, *Arithmetical Sacks Forcing*, Archive for Mathematical Logic **45** Issue 6 (2006), 715-720
- [14] Y. Ershov, *A hierarchy of sets, Part I*, Algebra and Logic **7** (1968) 24-43.
- [15] W. Fouche, *Arithmetical representations of Brownian motion*, Journal of Symbolic Logic **65** (2000), 421-442.
- [16] P. Gács, *On the symmetry of algorithmic information*, Soviet Mathematics (Doklady) **15** (1974), 1477-1480.
- [17] A. N. Kolmogorov, *Three approaches to the quantitative definition of information*, in Problems of Information Transmission, Vol. 1 (1965), 1-7.
- [18] Antonín Kučera, *Measure,  $\Pi_1^0$  classes and complete extensions of PA*, in Recursion Theory Week (Oberwolfach, 1984), Lecture Notes in Mathematics, Volume **1141** pages 245-259. Springer, Berlin, 1985.
- [19] *On the use of diagonally nonrecursive functions*, in Logic Colloquium '87, Ebbinghaus et al. eds., North Holland, 1989, pp. 219-239
- [20] L. Levin, *On the notion of a random sequence*, Soviet Mathematics (Doklady) **14** (1973), 1413-1416.
- [21] P. Lévy, *Théorie de l'Addition des Variables Aleatoires*, Gauthier-Villars, 1937 (second edition 1954).
- [22] M. Li and P. Vitanyi, *An introduction to Kolmogorov Complexity and Its Applications*, second edition, Springer (1997).
- [23] P. Martin-Löf, *The definition of random sequences*, Information and Control **9** (1966), 602-619.
- [24] A. A. Muchnik, A. L. Semenov and V. A. Uspensky, *Mathematical metaphysics of randomness*, Theoretical Computer Science **207** (1998), 263-271.

- [25] A. Nies, *Computability and Randomness*, in preparation. Current draft available at (<http://www.cs.auckland.ac.nz/~nies>).
- [26] C. P. Schnorr, A unified approach to the definition of random sequences, *Mathematical Systems Theory* **5** (1971), 246-258.
- [27] R. Soare, *Recursively Enumerable Sets and Degrees*, Springer-Verlag (1987).
- [28] M. van Lambalgen, *Random Sequences*, Ph.D. Dissertation, University of Amsterdam, The Netherlands (1987).
- [29] J. Ville, *Étude Critique de la Notion de Collectif*. Gauthier-Villars, Paris, 1939.
- [30] R. von Mises, *Grundlagen der Wahrscheinlichkeitsrechnung*, *Mathematische Zeitschrift* **5** (1919), 52-99.