

Quiz 9 Solutions
 MAC 1147.3077, Fall 2015
 Thursday, November 19, 2015

Show all relevant work to support your answer. A correct answer without supporting work will not earn the points. **Problems 3 and 4 are on the back.**

1. (1 point) What's your favorite quote (i.e from a movie, book, etc.)? (Hint: There is no wrong answer)

Solution: Answers vary, but the most acceptable is:
 "The first rule of Fight Club is: you do not talk about Fight Club."

2. (4 points) Evaluate the inverse trigonometric expressions:


(a) $\arctan\left(\frac{\sqrt{3}}{3}\right)$

Solution: The above statement translates to finding θ in the statement $\tan \theta = \frac{\sqrt{3}}{3}$. This occurs when $\theta = \frac{\pi}{6}$ or 30° .

Note, we used the fact that the range of $\arctan x$ is $[-\frac{\pi}{2}, \frac{\pi}{2}]$, which is the domain of the restricted tangent function. This is why we specifically chose $\frac{\pi}{6}$ as the solution.


(b) $\cot(\arctan \frac{1}{x})$

Solution: If we let $\arctan \frac{1}{x} = \theta$, then $\tan \theta = \frac{1}{x}$ (after taking tan of both sides). Hence we get the following diagram:




Now since we let $\arctan \frac{1}{x} = \theta$, our problem translates to finding $\cot \theta$ using our triangle. Hence, $\cot \theta = x$.

3. (2 points) Find the altitude of the triangle shown below given that $\theta = 18^\circ$ and $b = 10$.



Solution: First, draw a line from the top of the triangle to the midpoint of the base of the triangle. The length of this segment, denote by a , is the altitude. In short, we have divided the triangle into two right triangles. The measure of the base of each right triangle will be $10/2 = 5$. Thus, using the identity $\tan(18^\circ) = \frac{a}{5}$, we see $a = 5 \tan(18^\circ)$.

4. (3 points) Given $a = 3$ and $c = 5$, find the remaining sides and angles.



Solution: First, using the Pythagorean theorem, we see $3^2 + b^2 = 5^2$, and so $b = 4$.

Next, we use trig identities to find the missing angles. Hence $\tan(\alpha) = \frac{3}{4}$, or $\alpha = \arctan \frac{3}{4}$ (take arctan of both sides). Lastly, use the identity $\tan(\theta) = \frac{4}{3}$, or $\theta = \arctan \frac{4}{3}$ (again after taking arctan of both sides).

Note, you could have used other identities to get α and θ .