

1. (2 points) Find the point on the graph  $y = \frac{1}{2}x^2$  closest to the point  $(0, 1)$ .

**Solution.** The distance between a point  $(x, \frac{1}{2}x^2)$  on the curve  $y = \frac{1}{2}x^2$  and the point  $(0, 1)$  is given by

$$\begin{aligned}d(x) &= \sqrt{(x-0)^2 + \left(\frac{1}{2}x^2 - 1\right)^2} \\&= \sqrt{x^2 + \frac{1}{4}x^4 - x^2 + 1} \\&= \sqrt{\frac{1}{4}(x^4 + 4)} \\&= \frac{1}{2}\sqrt{x^4 + 4}.\end{aligned}$$

Now, we find the absolute minimum of  $d$  on its domain  $(-\infty, \infty)$ .

$$\begin{aligned}d'(x) &= \frac{d}{dx} \left( \frac{1}{2}\sqrt{x^4 + 4} \right) \\&= \frac{1}{4} \frac{4x^3}{\sqrt{x^4 + 4}} \\&= \frac{x^3}{\sqrt{x^4 + 4}}.\end{aligned}$$

Since the denominator  $\sqrt{x^4 + 4}$  is always positive, our only critical point is where the numerator is 0, meaning  $x = 0$ , and this point must represent an absolute minimum of  $d$ . Thus the closest point to  $(0, 1)$  on  $y = \frac{1}{2}x^2$  is  $(0, 0)$ .

$$\boxed{(0, 0)}$$

2. (4 points) Sketch the graph of  $f(x)$ , given

$$f(x) = \frac{4x - 3}{2x + 4} \quad f'(x) = \frac{22}{(2x + 4)^2} \quad f''(x) = \frac{-44}{(2x + 4)^3}$$

Be sure to include asymptotes,  $x$ - and  $y$ -intercepts, where  $f$  is increasing/decreasing and concave up/down.

**Solution.** First, the horizontal asymptote(s):

$$\lim_{x \rightarrow -\infty} \frac{4x - 3}{2x + 4} = \frac{4}{2} = 2 \quad \text{and} \quad \lim_{x \rightarrow +\infty} \frac{4x - 3}{2x + 4} = \frac{4}{2} = 2,$$

using either L'Hospital's Rule or the ratio of the leading coefficients. Now, the only candidate for a vertical asymptote is when  $2x + 4 = 0$ , meaning  $x = -2$ . Indeed,

$$\lim_{x \rightarrow -2^-} \frac{4x - 3}{2x + 4} = \frac{-11}{0^-} = +\infty \quad \text{and} \quad \lim_{x \rightarrow -2^+} \frac{4x - 3}{2x + 4} = \frac{-11}{0^+} = -\infty,$$

so  $x = -2$  does represent a vertical asymptote. For our  $x$ -intercept, we have  $f(x) = 0$  when  $4x - 3 = 0$ , so  $(3/4, 0)$  is our  $x$ -intercept. For our  $y$ -intercept, we have  $f(0) = -3/4$ , so  $(0, -3/4)$  is our  $y$ -intercept.

Both  $f'$  and  $f''$  are undefined at  $x = -2$ , so that point should go on both the  $f'$  and  $f''$  sign lines. Checking the sign of both  $f'$  and  $f''$  on either side of this point lets us finish our sign lines. Together with our intercepts and asymptotes, we can sketch our graph.

