

MAA6617 COURSE NOTES SPRING 2020

CONTENTS

20. Normed vector spaces	1
20.1. Examples	3
20.2. Linear transformations between normed spaces	6
20.3. Examples	8
20.4. Problems	9
21. Linear functionals and the Hahn-Banach theorem	11
21.1. Examples	12
21.2. The Hahn-Banach Extension Theorem	14
21.3. Dual spaces and adjoint operators	18
21.4. Duality for Sub and Quotient Spaces	18
21.5. Problems	20
22. The Baire Category Theorem and applications	23
22.1. Problems	28

20. NORMED VECTOR SPACES

Let $\mathcal{X}$ be a vector space over a field $\mathbb{K}$ (in this course we always have either $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$).

Definition 20.1. A *norm* on $\mathcal{X}$ is a function $\|\cdot\| : \mathcal{X} \rightarrow \mathbb{R}$ satisfying:

- (i) (positivity) $\|x\| \geq 0$ for all $x \in \mathcal{X}$, and $\|x\| = 0$ if and only if $x = 0$;
- (ii) (homogeneity) $\|kx\| = |k|\|x\|$ for all $x \in \mathcal{X}$ and $k \in \mathbb{K}$, and
- (iii) (triangle inequality) $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in \mathcal{X}$.

◁

Using these three properties it is straightforward to check that the quantity

$$d(x, y) := \|x - y\|$$

defines a metric on $\mathcal{X}$. The resulting topology is the *norm topology*. The next proposition is simple but fundamental; it says that the norm and the vector space operations are continuous in the norm topology.

Proposition 20.2 (Continuity of vector space operations). *Let $\mathcal{X}$ be a normed vector space over $\mathbb{K}$.*

- a) *If (x_n) converges to x in $\mathcal{X}$, then $(\|x_n\|)$ converges to $\|x\|$ in $\mathbb{R}$.*
- b) *If (k_n) converges to k in $\mathbb{K}$ and (x_n) converges to x in $\mathcal{X}$, then $(k_n x_n)$ converges to kx in $\mathcal{X}$.*
- c) *If (x_n) converges to x and (y_n) converges to y in $\mathcal{X}$, then $(x_n + y_n)$ converges to $x + y$ in $\mathcal{X}$.*

Proof. The proofs follow readily from the properties of the norm, and are left as exercises. $\square$

Two norms $\|\cdot\|_1, \|\cdot\|_2$ on $\mathcal{X}$ are *equivalent* if there exist absolute constants $C, c > 0$ such that

$$c\|x\|_1 \leq \|x\|_2 \leq C\|x\|_1 \quad \text{for all } x \in \mathcal{X}.$$

Equivalent norms determine the same topology on $\mathcal{X}$ and the same Cauchy sequences (Problem 20.2). A normed space is a *Banach space* if it is complete in the norm topology. It follows that if $\mathcal{X}$ is equipped with two equivalent norms $\|\cdot\|_1, \|\cdot\|_2$ then it is complete (a Banach space) in one norm if and only if it is complete in the other.

The following proposition gives a convenient criterion for a normed vector space to be complete. A series $\sum_{n=1}^{\infty} x_n$ in $\mathcal{X}$ is *absolutely convergent* if $\sum_{n=1}^{\infty} \|x_n\| < \infty$. The series *converges* in $\mathcal{X}$ if the limit $\lim_{N \rightarrow \infty} \sum_{n=1}^N x_n$ exists in $\mathcal{X}$ (in the norm topology). (Quite explicitly, the series $\sum_{n=1}^{\infty} x_n$ converges to $x \in \mathcal{X}$ if $\lim_{N \rightarrow \infty} \left\| x - \sum_{n=1}^N x_n \right\| = 0$.)

Proposition 20.3. *A normed space $(\mathcal{X}, \|\cdot\|)$ is complete if and only if every absolutely convergent series in $\mathcal{X}$ is convergent.*

Proof. First suppose $\mathcal{X}$ is complete and $\sum_{n=1}^{\infty} x_n$ is absolutely convergent. Write $s_N = \sum_{n=1}^N x_n$ for the N^{th} partial sum and let $\epsilon > 0$ be given. Since $\sum_{n=1}^{\infty} \|x_n\|$ is convergent, there exists an L such that $\sum_{n=L}^{\infty} \|x_n\| < \epsilon$. If $N > M \geq L$, then

$$\|s_N - s_M\| = \left\| \sum_{n=M+1}^N x_n \right\| \leq \sum_{n=M+1}^N \|x_n\| < \epsilon.$$

Thus the sequence (s_N) is Cauchy in $\mathcal{X}$, hence convergent by hypothesis.

Conversely, suppose every absolutely convergent series in $\mathcal{X}$ is convergent. Given a Cauchy sequence (x_n) from $\mathcal{X}$, choose a super-Cauchy subsequence (y_k) ; i.e., $(y_k = x_{n_k})_k$ and

$$\sum_{k=1}^{\infty} \|y_{k+1} - y_k\| < \infty.$$

(To do this, first choose N_1 such that $\|x_n - x_m\| < 2^{-1}$ for all $n, m \geq N_1$. Next choose $N_2 > N_1$ such that $\|x_n - x_m\| < 2^{-2}$ for all $n, m \geq N_2$. Continuing in this way recursively defines an increasing sequence of integers $(N_k)_{k=1}^{\infty}$ such that $\|x_n - x_m\| < 2^{-k}$ for all $n, m \geq N_k$. Set $y_k = x_{N_k}$.) The series $\sum_{k=1}^{\infty} (y_{k+1} - y_k)$ is absolutely convergent and

hence, by hypothesis, convergent in $\mathcal{X}$. In other words, the sequence $(y_k - y_1)_k$ of partial sums converges in $\mathcal{X}$ which means that (x_n) has a convergent subsequence. The proof is finished by invoking a standard fact about convergence in metric spaces: if (x_n) is a Cauchy sequence which has a convergent subsequence, then the full sequence converges. $\square$

20.1. Examples.

- (a) Of course, $\mathbb{K}^n$ with the usual Euclidean norm $\|(x_1, \dots, x_n)\| = (\sum_{k=1}^n |x_k|^2)^{1/2}$ is a Banach space. The vector space $\mathbb{K}^n$ can also be equipped with the ℓ^p -norms

$$\|(x_1, \dots, x_n)\|_p := \left(\sum_{k=1}^n |x_k|^p \right)^{1/p}$$

for $1 \leq p < \infty$, and the ℓ^∞ -norm

$$\|(x_1, \dots, x_n)\|_\infty := \max(|x_1|, \dots, |x_n|).$$

For $1 \leq p < \infty$ and $p \neq 2$, it is not immediately obvious that $\|\cdot\|_p$ defines a norm. We will prove this assertion later. It is not too hard to show that all of the ℓ^p norms ($1 \leq p \leq \infty$) are equivalent on $\mathbb{K}^n$ (though the constants c, C depend on the dimension n). It turns out that *any* two norms on a finite-dimensional vector space are equivalent. As a corollary, every finite-dimensional normed space is a Banach space. See Problem 20.3.

- (b) (Sequence spaces) Define

$$c_0 := \{f : \mathbb{N} \rightarrow \mathbb{K} \mid \lim_{m \rightarrow \infty} |f(m)| = 0\}$$

$$\ell^\infty := \{f : \mathbb{N} \rightarrow \mathbb{K} \mid \sup_{m \in \mathbb{N}} |f(m)| < \infty\}$$

$$\ell^1 := \{f : \mathbb{N} \rightarrow \mathbb{K} \mid \sum_{m=0}^{\infty} |f(m)| < \infty\}.$$

It is a simple exercise to check that each of these is a vector space (a subspace of the vector space of all functions $f : \mathbb{N} \rightarrow \mathbb{K}$). Define, for functions $f : \mathbb{N} \rightarrow \mathbb{K}$,

$$\|f\|_\infty := \sup_m |f(m)|$$

$$\|f\|_1 := \sum_{m=1}^{\infty} |f(m)|.$$

Then $\|f\|_\infty$ defines a norm on both c_0 and ℓ^∞ , and $\|f\|_1$ is a norm on ℓ^1 . Equipped with these respective norms, each is a Banach space. We sketch the proof for c_0 . Verification of the other two assertions is left as exercises (Problem 20.4).

The key observation is that (f_n) converges to f in the $\|\cdot\|_\infty$ norm if and only if (f_n) converges to f uniformly as functions on $\mathbb{N}$. Suppose (f_n) is a Cauchy sequence in c_0 . Then the sequence of functions f_n is uniformly Cauchy on $\mathbb{N}$, and in particular converges pointwise to a function f .

We must show that this f belongs to c_0 and that $f_n \rightarrow f$ uniformly on $\mathbb{N}$. For this, let $\epsilon > 0$ be given. There is an N so that $\|f_m - f_n\| < \epsilon$ for $m, n \geq N$. Thus, for each $n \geq N$, all $\ell \in \mathbb{N}$ and all $m \geq n \geq N$, $|f_m(\ell) - f_n(\ell)| < \epsilon$. Thus, fixing $n = N$ and letting m tend to infinity, we have $|f(\ell) - f_N(\ell)| \leq \epsilon$ for all ℓ . There is an M so that $|f_N(\ell)| < \epsilon$ for $\ell \geq M$. Hence, for such ℓ ,

$$|f(\ell)| \leq |f_N(\ell)| + \epsilon < 2\epsilon.$$

Thus $f \in c_0$ and (f_n) converges to f in c_0 .

Along with these spaces it is also helpful to consider the vector space

$$c_{00} := \{f : \mathbb{N} \rightarrow \mathbb{K} \mid f(n) = 0 \text{ for all but finitely many } n\}$$

Notice that c_{00} is a vector subspace of each of c_0 , ℓ^1 and ℓ^∞ . Thus it can be equipped with either the $\|\cdot\|_\infty$ or $\|\cdot\|_1$ norms. It is not complete in either of these norms, however. What is true is that c_{00} is *dense* in c_0 and ℓ^1 (but not in ℓ^∞). (See Problem 20.9).

(c) (L^1 spaces) Let $(X, \mathcal{M}, m)$ be a measure space. The quantity

$$\|f\|_1 := \int_X |f| dm$$

defines a norm on $L^1(m)$, provided we agree to identify f and g when $f = g$ a.e. (Indeed the chief motivation for making this identification is that it makes $\|\cdot\|_1$ into a norm. Note that ℓ^1 from the last example is a special case (what is the measure space?))

Proposition 20.4. $L^1(m)$ is a Banach space.

Proof. It suffices to verify the hypotheses of Proposition 20.3. If $\sum_{n=1}^\infty f_n$ is absolutely convergent (so that $\sum_{n=1}^\infty \|f_n\|_1 < \infty$), then

$$\sum_{n=1}^\infty |f_n| dm < \infty.$$

Thus the function $g := \sum_{n=1}^\infty |f_n|$ belongs to L^1 and is thus finite m -a.e. by Tonelli's theorem. In particular the sequence of partial sums $s_N = \sum_{n=1}^N f_n$ is a sequence of measurable functions with $|s_N| \leq g$ that converges pointwise a.e. to a measurable function f . Hence by the DCT and its corollary, $f \in L^1$ and the partial sums $(s_N)_N$ converge to f in L^1 . $\square$

(d) (L^p spaces) Again let $(X, \mathcal{M}, m)$ be a measure space. For $1 \leq p < \infty$ let $L^p(m)$ denote the set of measurable functions f for which

$$\|f\|_p := \left(\int_X |f|^p dm \right)^{1/p} < \infty$$

(again we identify f and g when $f = g$ a.e.). It turns out that this quantity is a norm on $L^p(m)$, and $L^p(m)$ is complete, though we will not prove this yet (it is not

immediately obvious that the triangle inequality holds when $p > 1$). The sequence space ℓ^p is defined analogously: it is the set of $f : \mathbb{N} \rightarrow \mathbb{K}$ for which

$$\|f\|_p := \left(\sum_{n=1}^{\infty} |f(n)|^p \right)^{1/p} < \infty$$

and this quantity is a norm making ℓ^p into a Banach space.

When $p = \infty$, and the measure m is σ -finite, we define $L^\infty(m)$ to be the set of all functions $f : X \rightarrow \mathbb{K}$ with the following property: there exists $M > 0$ such that

$$|f(x)| \leq M \quad \text{for } m - \text{a.e. } x \in X; \quad (1)$$

as for the other L^p spaces we identify f and g when there are equal a.e. When $f \in L^\infty$, let $\|f\|_\infty$ be the smallest M for which (1) holds. Then $\|\cdot\|_\infty$ is a norm making $L^\infty(m)$ into a Banach space.

- (e) ($C(X)$ spaces) Let X be a compact metric space and let $C(X)$ denote the set of continuous functions $f : X \rightarrow \mathbb{K}$. It is a standard fact from advanced calculus that the quantity $\|f\|_\infty := \sup_{x \in X} |f(x)|$ is a norm on $C(X)$. A sequence is Cauchy in this norm if and only if it is uniformly Cauchy. It is thus also a standard fact that $C(X)$ is complete in this norm—completeness just means that a uniformly Cauchy sequence of continuous functions on X converges uniformly to a continuous function.

This example can be generalized somewhat: let X be a locally compact metric space. Say a function $f : X \rightarrow \mathbb{K}$ *vanishes at infinity* if for every $\epsilon > 0$, there exists a compact set $K \subset X$ such that $\sup_{x \notin K} |f(x)| < \epsilon$. Let $C_0(X)$ denote the set of continuous functions $f : X \rightarrow \mathbb{K}$ that vanish at infinity. Then $C_0(X)$ is a vector space, the quantity $\|f\|_\infty := \sup_{x \in X} |f(x)|$ is a norm on $C_0(X)$, and $C_0(X)$ is complete in this norm. (Note that c_0 from above is a special case.)

- (f) (Subspaces and direct sums) If $(\mathcal{X}, \|\cdot\|)$ is a normed vector space and $\mathcal{Y} \subset \mathcal{X}$ is a vector subspace, then the restriction of $\|\cdot\|$ to $\mathcal{Y}$ is clearly a norm on $\mathcal{Y}$. If $\mathcal{X}$ is a Banach space, then $(\mathcal{Y}, \|\cdot\|)$ is a Banach space if and only if $\mathcal{Y}$ is *closed* in the norm topology of $\mathcal{X}$. (This is just a standard fact about metric spaces—a subspace of a complete metric space is complete in the restricted metric if and only if it is closed.)

If $\mathcal{X}, \mathcal{Y}$ are vector spaces then the *algebraic direct sum* is the vector space of ordered pairs

$$\mathcal{X} \oplus \mathcal{Y} := \{(x, y) : x \in \mathcal{X}, y \in \mathcal{Y}\}$$

with entrywise operations. If $\mathcal{X}, \mathcal{Y}$ are equipped with norms $\|\cdot\|_{\mathcal{X}}, \|\cdot\|_{\mathcal{Y}}$, then each of the quantities

$$\begin{aligned} \|(x, y)\|_\infty &:= \max(\|x\|_{\mathcal{X}}, \|y\|_{\mathcal{Y}}), \\ \|(x, y)\|_1 &:= \|x\|_{\mathcal{X}} + \|y\|_{\mathcal{Y}} \end{aligned}$$

is a norm on $\mathcal{X} \oplus \mathcal{Y}$. These two norms are equivalent; indeed it follows from the definitions that

$$\|(x, y)\|_\infty \leq \|(x, y)\|_1 \leq 2\|(x, y)\|_\infty.$$

If $\mathcal{X}$ and $\mathcal{Y}$ are both complete, then $\mathcal{X} \oplus \mathcal{Y}$ is complete in both of these norms. The resulting Banach spaces are denoted $\mathcal{X} \oplus_\infty \mathcal{Y}$, $\mathcal{X} \oplus_1 \mathcal{Y}$ respectively.

- (g) (Quotient spaces) If $\mathcal{X}$ is a normed vector space and $\mathcal{M}$ is a proper subspace, then one can form the *algebraic quotient* $\mathcal{X}/\mathcal{M}$, defined as the collection of distinct cosets $\{x + \mathcal{M} : x \in \mathcal{X}\}$. From linear algebra, $\mathcal{X}/\mathcal{M}$ is a vector space under the standard operations. If $\mathcal{M}$ is a *closed* subspace of $\mathcal{X}$, then the quantity

$$\|x + \mathcal{M}\| := \inf_{y \in \mathcal{M}} \|x - y\|$$

is a norm on $\mathcal{X}/\mathcal{M}$, called the *quotient norm*. (Geometrically, $\|x + \mathcal{M}\|$ is the distance in $\mathcal{X}$ from x to the closed set $\mathcal{M}$.) It turns out that if $\mathcal{X}$ is complete, so is $\mathcal{X}/\mathcal{M}$. See Problem 20.20.

More examples are given in the exercises and further examples will appear after the development of some theory.

20.2. Linear transformations between normed spaces.

Definition 20.5. Let $\mathcal{X}, \mathcal{Y}$ be normed vector spaces. A linear transformation $T : \mathcal{X} \rightarrow \mathcal{Y}$ is *bounded* if there exists a constant $C > 0$ such that $\|Tx\|_{\mathcal{Y}} \leq C\|x\|_{\mathcal{X}}$ for all $x \in \mathcal{X}$. $\triangleleft$

Remark 20.6. Note that in this definition it would suffice to require that $\|Tx\|_{\mathcal{Y}} \leq C\|x\|_{\mathcal{X}}$ just for all $x \neq 0$, or for all x with $\|x\|_{\mathcal{X}} = 1$ (why?) $\diamond$

The importance of boundedness and the following simple proposition is hard to overstate. Recall, a mapping $f : X \rightarrow Y$ between metric spaces is *Lipschitz continuous* if there is a constant $C > 0$ such that $d(f(x), f(y)) \leq Cd(x, y)$ for all $x, y \in X$. A simple exercise shows Lipschitz continuity implies (uniform) continuity.

Proposition 20.7. If $T : \mathcal{X} \rightarrow \mathcal{Y}$ is a linear transformation between normed spaces, then the following are equivalent:

- (i) T is bounded.
- (ii) T is Lipschitz continuous.
- (iii) T is uniformly continuous.
- (iv) T is continuous.
- (v) T is continuous at 0.

Moreover, in this case,

$$\begin{aligned} \|T\| &:= \sup\{\|Tx\| : \|x\| = 1\} \\ &= \sup\left\{\frac{\|Tx\|}{\|x\|} : x \neq 0\right\} \\ &= \inf\{C : \|Tx\| \leq C\|x\| \text{ for all } x \in \mathcal{X}\} \end{aligned}$$

and $\|T\|$ is the smallest number (the infimum is attained) such that

$$\|Tx\| \leq \|T\| \|x\| \tag{2}$$

for all $x \in \mathcal{X}$.

Proof. Suppose T is bounded so that there exists a $C > 0$ such that $\|Tx\| \leq C\|x\|$ for all $x \in \mathcal{X}$. Thus, if $x, y \in \mathcal{X}$, then, $\|Tx - Ty\| = \|T(x - y)\| \leq C\|x - y\|$ by linearity of T . Hence (i) implies (ii). The implications (ii) implies (iii) implies (iv) implies (v) are evident. The proof of (v) implies (i) exploits the homogeneity of the norm and the linearity of T . By hypothesis, with $\epsilon = 1$ there exists $\delta > 0$ such that if $\|x\| < \delta$, then $\|Tx\| < 1$. Fix a nonzero vector $x \in \mathcal{X}$ and a real number $0 < \lambda < \delta$. The vector $\lambda x / \|x\|$ has norm less than δ , so

$$\left\| T \left(\frac{\lambda x}{\|x\|} \right) \right\| = \lambda \frac{\|Tx\|}{\|x\|} < 1.$$

Rearranging this we find $\|Tx\| \leq (1/\lambda)\|x\|$ for all $x \neq 0$, which shows T is bounded; in fact we can take $C = \frac{1}{\delta}$.

The rest of the proof is left as an exercise. $\square$

The set of all bounded linear operators from $\mathcal{X}$ to $\mathcal{Y}$ is denoted $B(\mathcal{X}, \mathcal{Y})$. It is a vector space under the operations of pointwise addition and scalar multiplication. The quantity $\|T\|$ is called the *operator norm* of T .

Proposition 20.8. *For normed vector spaces $\mathcal{X}$ and $\mathcal{Y}$, the operator norm makes $B(\mathcal{X}, \mathcal{Y})$ into a normed vector space that is complete if $\mathcal{Y}$ is complete.*

Proof. That $B(\mathcal{X}, \mathcal{Y})$ is a normed vector space follows readily from the definitions and is left as an exercise. Suppose now $\mathcal{Y}$ is complete, and let T_n be a Cauchy sequence in $B(\mathcal{X}, \mathcal{Y})$. For each $x \in \mathcal{X}$, we have

$$\|T_n x - T_m x\| = \|(T_n - T_m)x\| \leq \|T_n - T_m\| \|x\| \quad (3)$$

which shows that $(T_n x)$ is a Cauchy sequence in $\mathcal{Y}$. By hypothesis, $T_n x$ converges in $\mathcal{Y}$. Define $T : \mathcal{X} \rightarrow \mathcal{Y}$ by setting $Tx := y$. It is straightforward to check that T is linear.

Let B denote the closed unit ball in $\mathcal{X}$. The sequence $(T_n|_B)$ is uniformly Cauchy by equation (3) and converges pointwise to $T|_B$. Hence $T|_B$ is continuous and $(T_n|_B)$ converges uniformly to $T|_B$. It follows that T is continuous at 0 and therefore T is continuous. Since $\|T_n - T\| = \sup\{\|(T_n - T)x\| : x \in B\}$ and since $(T_n|_B)$ converges to $T|_B$ uniformly, it follows that (T_n) converges to T in $B(\mathcal{X}, \mathcal{Y})$. $\square$

If $T \in B(\mathcal{X}, \mathcal{Y})$ and $S \in B(\mathcal{Y}, \mathcal{Z})$, then two applications of the inequality (2) gives, for $x \in \mathcal{X}$,

$$\|STx\| \leq \|S\| \|Tx\| \leq \|S\| \|T\| \|x\|$$

and it follows that $ST \in B(\mathcal{X}, \mathcal{Z})$ and $\|ST\| \leq \|S\| \|T\|$. In the special case that $\mathcal{Y} = \mathcal{X}$ is complete, $B(\mathcal{X}) := B(\mathcal{X}, \mathcal{X})$ is an example of a *Banach algebra*.

The following proposition is very useful in constructing bounded operators—at least when the codomain is complete. Namely, it suffices to define the operator (and show that it is bounded) on a dense subspace.

Proposition 20.9 (Extending bounded operators). *Let $\mathcal{X}, \mathcal{Y}$ be normed vector spaces with $\mathcal{Y}$ complete, and $\mathcal{E} \subset \mathcal{X}$ a dense linear subspace. If $T : \mathcal{E} \rightarrow \mathcal{Y}$ is a bounded linear operator, then there exists a unique bounded linear operator $\tilde{T} : \mathcal{X} \rightarrow \mathcal{Y}$ extending T (so $\tilde{T}|_{\mathcal{E}} = T$). Further $\|\tilde{T}\| = \|T\|$.*

Proof. Recall, if X, Y are metric spaces, Y is complete, $D \subset X$ is dense and $f : D \rightarrow Y$ is uniformly continuous, then f has a unique continuous extension $\tilde{f} : X \rightarrow Y$. Moreover, this extension can be defined as follows. Given $x \in X$, choose a sequence (x_n) from D converging to x and let $\tilde{f}(x) = \lim f(x_n)$ (that the sequence $f(x_n)$ is Cauchy follows from uniform continuity; that it converges from the assumption that $\mathcal{Y}$ is complete and finally it is an exercise to show $\tilde{f}(x)$ is well defined independent of the choice of (x_n)). Thus, it only remains to verify that the extension $\tilde{T}$ of T is in fact linear and $\|T\| = \|\tilde{T}\|$. Both are routine exercises. $\square$

Remark: The completeness of $\mathcal{Y}$ is essential in the above proposition; Problem 20.11 suggests a counterexample.

A bounded linear transformation $T \in B(\mathcal{X}, \mathcal{Y})$ is said to be *invertible* if it is bijective (being bijective, automatically T^{-1} exists and is a linear transformation) and T^{-1} is bounded from $\mathcal{Y}$ to $\mathcal{X}$. Two normed spaces $\mathcal{X}, \mathcal{Y}$ are said to be (*boundedly*) *isomorphic* if there exists an invertible linear transformation $T : \mathcal{X} \rightarrow \mathcal{Y}$. As an example, given equivalent norms $\|\cdot\|_1$ and $\|\cdot\|_2$ on a vector space $\mathcal{X}$, the identity mapping $\iota : (\mathcal{X}, \|\cdot\|_1) \rightarrow (\mathcal{X}, \|\cdot\|_2)$ is (*boundedly*) invertible and witnesses the fact that these two normed vector spaces are boundedly isomorphic.

An operator $T : \mathcal{X} \rightarrow \mathcal{Y}$ such that $\|Tx\| = \|x\|$ for all $x \in \mathcal{X}$ is an *isometry*. Note that an isometry is automatically injective and if it is also surjective then it is automatically invertible and T^{-1} is also an isometry. An isometry need not be surjective, however. The normed vector spaces are *isometrically isomorphic* if there is an invertible isometry $T : \mathcal{X} \rightarrow \mathcal{Y}$.

20.3. Examples.

- (a) If $\mathcal{X}$ is a finite-dimensional normed space and $\mathcal{Y}$ is any normed space, then every linear transformation $T : \mathcal{X} \rightarrow \mathcal{Y}$ is bounded. See Problem 20.14
- (b) Let $\mathcal{X}$ denote c_{00} equipped with the $\|\cdot\|_1$ norm, and $\mathcal{Y}$ denote c_{00} equipped with the $\|\cdot\|_{\infty}$ norm. Then the identity map $id_{\mathcal{X}, \mathcal{Y}} : \mathcal{X} \rightarrow \mathcal{Y}$ is bounded as an operator (in fact its norm is equal to 1), but its inverse, the identity map $\iota_{\mathcal{Y}, \mathcal{X}} : \mathcal{Y} \rightarrow \mathcal{X}$ is unbounded.
- (c) Consider c_{00} with the $\|\cdot\|_{\infty}$ norm. Let $a : \mathbb{N} \rightarrow \mathbb{K}$ be any function and define a linear transformation $T_a : c_{00} \rightarrow c_{00}$ by

$$T_a f(n) = a(n)f(n). \quad (4)$$

The mapping T_a is bounded if and only if $M = \sup_{n \in \mathbb{N}} |a(n)| < \infty$, in which case $\|T_a\| = M$. In this case, T_a extends uniquely to a bounded operator from c_0 to c_0 ,

and one may check that the formula (4) defines the extension. All of these claims remain true if we use the $\|\cdot\|_1$ norm instead of the $\|\cdot\|_\infty$ norm. In this case, we get a bounded operator from ℓ^1 to itself.

- (d) Define $S : \ell^1 \rightarrow \ell^1$ as follows given the sequence $(f(n))_n$ from ℓ^1 let $Sf(1) = 0$ and $Sf(n) = f(n-1)$ for $n > 1$. (Viewing f as a sequence, S shifts the sequence one place to the right and fills in a 0 in the first position). This S is an isometry, but is not surjective. In contrast, if $\mathcal{X}$ is finite-dimensional, then the rank-nullity theorem from linear algebra guarantees that every injective linear map $T : \mathcal{X} \rightarrow \mathcal{X}$ is also surjective.
- (e) Let $C^\infty([0, 1])$ denote the vector space of functions on $[0, 1]$ with continuous derivatives of all orders. The differentiation map $D : C^\infty([0, 1]) \rightarrow C^\infty([0, 1])$ defined by $Df = \frac{df}{dx}$ is a linear transformation. Since, for $t \in \mathbb{R}$, we have $De^{tx} = te^{tx}$, it follows that there is *no* norm on $C^\infty([0, 1])$ such that $\frac{d}{dx}$ is bounded.

20.4. Problems.

Problem 20.1. Prove Proposition 20.2.

Problem 20.2. Prove equivalent norms define the same topology and the same Cauchy sequences.

Problem 20.3. (a) Prove all norms on a finite dimensional vector space $\mathcal{X}$ are equivalent. Suggestion: Fix a basis $e_1, \dots, e_n$ for $\mathcal{X}$ and define $\|\sum a_k e_k\|_1 := \sum |a_k|$. It is routine to check that $\|\cdot\|_1$ is a norm on $\mathcal{X}$. Now complete the following outline.

- (i) Let $\|\cdot\|$ be the given norm on $\mathcal{X}$. Show there is an M such that $\|x\| \leq M\|x\|_1$. Conclude that the mapping $\iota : (\mathcal{X}, \|\cdot\|_1) \rightarrow (\mathcal{X}, \|\cdot\|)$ defined by $\iota(x) = x$ is continuous;
 - (ii) Show that the *unit sphere* $S = \{x \in \mathcal{X} : \|x\|_1 = 1\}$ in $(\mathcal{X}, \|\cdot\|_1)$ is compact in the $\|\cdot\|_1$ topology;
 - (iii) Show that the mapping $f : S \rightarrow (\mathcal{X}, \|\cdot\|)$ given by $f(x) = \|x\|$ is continuous and hence attains its infimum. Show this infimum is not 0 and finish the proof.
- (b) Combine the result of part (a) with the result of Problem 20.2 to conclude that every finite-dimensional normed vector space is complete.
- (c) Let $\mathcal{X}$ be a normed vector space and $\mathcal{M} \subset \mathcal{X}$ a finite-dimensional subspace. Prove $\mathcal{M}$ is closed in $\mathcal{X}$.

Problem 20.4. Finish the proofs from Example 20.1(b).

Problem 20.5. A function $f : [0, 1] \rightarrow \mathbb{K}$ is called *Lipschitz continuous* if there exists a constant C such that

$$|f(x) - f(y)| \leq C|x - y|$$

for all $x, y \in [0, 1]$. Define $\|f\|_{Lip}$ to be the best possible constant in this inequality. That is,

$$\|f\|_{Lip} := \sup_{x \neq y} \frac{|f(x) - f(y)|}{|x - y|}$$

Let $\text{Lip}[0, 1]$ denote the set of all Lipschitz continuous functions on $[0, 1]$. Prove $\|f\| := |f(0)| + \|f\|_{\text{Lip}}$ is a norm on $\text{Lip}[0, 1]$, and that $\text{Lip}[0, 1]$ is complete in this norm.

Problem 20.6. Let $C^1([0, 1])$ denote the space of all functions $f : [0, 1] \rightarrow \mathbb{R}$ such that f is differentiable in $(0, 1)$ and f' extends continuously to $[0, 1]$. Prove

$$\|f\| := \|f\|_\infty + \|f'\|_\infty$$

is a norm on $C^1([0, 1])$ and that C^1 is complete in this norm. Do the same for the norm $\|f\| := |f(0)| + \|f'\|_\infty$. (Is $\|f'\|_\infty$ a norm on C^1 ?)

Problem 20.7. Let $(X, \mathcal{M})$ be a measurable space. Let $M(X)$ denote the (real) vector space of all signed measures on $(X, \mathcal{M})$. Prove the *total variation norm* $\|\mu\| := |\mu|(X)$ is a norm on $M(X)$, and $M(X)$ is complete in this norm.

Problem 20.8. Prove, if $\mathcal{X}, \mathcal{Y}$ are normed spaces, then the operator norm is a norm on $B(\mathcal{X}, \mathcal{Y})$.

Problem 20.9. Prove c_{00} is dense in c_0 and ℓ^1 . (That is, given $f \in c_0$ there is a sequence f_n in c_{00} such that $\|f_n - f\|_\infty \rightarrow 0$, and the analogous statement for ℓ^1 .) Using these facts, or otherwise, prove that c_{00} is *not* dense in ℓ^∞ . (In fact there exists $f \in \ell^\infty$ with $\|f\|_\infty = 1$ such that $\|f - g\|_\infty \geq 1$ for all $g \in c_{00}$.)

Problem 20.10. Prove c_{00} is not complete in the $\|\cdot\|_1$ or $\|\cdot\|_\infty$ norms. (After we have studied the Baire Category theorem, you will be asked to prove that there is *no* norm on c_{00} making it complete.)

Problem 20.11. Consider c_0 and c_{00} equipped with the $\|\cdot\|_\infty$ norm. Prove there is no bounded operator $T : c_0 \rightarrow c_{00}$ such that $T|_{c_{00}}$ is the identity map. (Thus the conclusion of Proposition 20.9 can fail if $\mathcal{Y}$ is not complete.)

Problem 20.12. Prove the $\|\cdot\|_1$ and $\|\cdot\|_\infty$ norms on c_{00} are not equivalent. Conclude from your proof that the identity map on c_{00} is bounded from the $\|\cdot\|_1$ norm to the $\|\cdot\|_\infty$ norm, but not the other way around.

Problem 20.13. a) Prove $f \in C_0(\mathbb{R}^n)$ if and only if f is continuous and $\lim_{|x| \rightarrow \infty} |f(x)| = 0$. b) Let $C_c(\mathbb{R}^n)$ denote the set of continuous, compactly supported functions on $\mathbb{R}^n$. Prove $C_c(\mathbb{R}^n)$ is dense in $C_0(\mathbb{R}^n)$ (where $C_0(\mathbb{R}^n)$ is equipped with sup norm).

Problem 20.14. Prove, if $\mathcal{X}, \mathcal{Y}$ are normed spaces and $\mathcal{X}$ is finite dimensional, then every linear transformation $T : \mathcal{X} \rightarrow \mathcal{Y}$ is bounded. Suggestion: Let d denote the dimension of X and let $\{e_1, \dots, e_d\}$ denote a basis. The function $\|\cdot\|_1$ on $\mathcal{X}$ defined by $\|\sum x_j e_j\|_1 = \sum |x_j|$ is a norm. Apply Problem 20.3.

Problem 20.15. Prove the claims in Example 20.3(c).

Problem 20.16. Let $g : \mathbb{R} \rightarrow \mathbb{K}$ be a (Lebesgue) measurable function. The map $M_g : f \rightarrow gf$ is a linear transformation on the space of measurable functions. Prove, if $g \notin L^\infty(\mathbb{R})$, then there is an $f \in L^1(\mathbb{R})$ such that $gf \notin L^1(\mathbb{R})$. Conversely, show if $g \in L^\infty(\mathbb{R})$, then M_g is bounded from $L^1(\mathbb{R})$ to itself and $\|M_g\| = \|g\|_\infty$.

Problem 20.17. Prove the claims about direct sums in Example 20.1(f).

Problem 20.18. Let $\mathcal{X}$ be a normed vector space and $\mathcal{M}$ a proper *closed* subspace. Prove for every $\epsilon > 0$, there exists $x \in \mathcal{X}$ such that $\|x\| = 1$ and $\inf_{y \in \mathcal{M}} \|x - y\| > 1 - \epsilon$. (Hint: take any $u \in \mathcal{X} \setminus \mathcal{M}$ and let $a = \inf_{y \in \mathcal{M}} \|u - y\|$. Choose $\delta > 0$ small enough so that $\frac{a}{a+\delta} > 1 - \epsilon$, and then choose $v \in \mathcal{M}$ so that $\|u - v\| < a + \delta$. Finally let $x = \frac{u-v}{\|u-v\|}$.)

Note that the distance to a (closed) subspace need not be attained. Here is an example. Consider the Banach space $C([0, 1])$ (with the sup norm of course and either real or complex valued functions) and the closed subspace

$$T = \{f \in C([0, 1]) : f(0) = 0 = \int_0^1 f dt\}.$$

Using machinery in the next section it will be evident that T is a closed subspace of $C([0, 1])$. For now, it can be easily verified directly. Let g denote the function $g(t) = t$. Verify that, for $f \in T$, that

$$\frac{1}{2} = \int_0^1 g dt = \int_0^1 (g - f) dt \leq \|g - f\|_\infty.$$

In particular, the distance from g to T is at least $\frac{1}{2}$.

Note that the function $h = x - \frac{1}{2}$, while not in T , satisfies $\|g - h\|_\infty = \frac{1}{2}$.

On the other hand, for any $\epsilon > 0$ there is an $f \in T$ so that $\|g - f\|_\infty \leq \frac{1}{2} + \epsilon$ (simply modify h appropriately). Thus, the distance from g to T is $\frac{1}{2}$. Now verify, using the inequality above, that h is the only element of $C([0, 1])$ such that $\int h dt = 0$ and $\|g - h\|_\infty = \frac{1}{2}$.

Problem 20.19. Prove, if $\mathcal{X}$ is an infinite-dimensional normed space, then the unit ball $ball(\mathcal{X}) := \{x \in \mathcal{X} : \|x\| \leq 1\}$ is not compact in the norm topology. (Hint: use the result of Problem 20.18 to construct inductively a sequence of vectors $x_n \in \mathcal{X}$ such that $\|x_n\| = 1$ for all n and $\|x_n - x_m\| \geq \frac{1}{2}$ for all $m < n$.)

Problem 20.20. (The quotient norm) Let $\mathcal{X}$ be a normed space and $\mathcal{M}$ a proper closed subspace.

- Prove the quotient norm is a norm (see Example 20.1(g)).
- Show that the quotient map $x \rightarrow x + \mathcal{M}$ has norm 1. (Use Problem 20.18.)
- Prove, if $\mathcal{X}$ is complete, so is $\mathcal{X}/\mathcal{M}$.

Problem 20.21. A normed vector space $\mathcal{X}$ is called *separable* if it is separable as a metric space (that is, there is a countable subset of $\mathcal{X}$ which is dense in the norm topology). Prove c_0 and ℓ^1 are separable, but ℓ^∞ is not. (Hint: for ℓ^∞ , show that there is an uncountable collection of elements $\{f_\alpha\}$ such that $\|f_\alpha - f_\beta\| = 1$ for $\alpha \neq \beta$.)

21. LINEAR FUNCTIONALS AND THE HAHN-BANACH THEOREM

If there is a “fundamental theorem of functional analysis,” it is the Hahn-Banach theorem. The theorem is somewhat abstract-looking at first, but its importance will be clear after studying some of its corollaries.

Let $\mathcal{X}$ be a normed vector space over the field $\mathbb{K}$. A *linear functional* on $\mathcal{X}$ is a linear map $L : \mathcal{X} \rightarrow \mathbb{K}$. As one might expect, we are especially interested in bounded linear functionals. Since $\mathbb{K} = \mathbb{R}$ or $\mathbb{C}$ is complete, the vector space of bounded linear functionals $B(\mathcal{X}, \mathbb{K})$ is itself a Banach space (complete normed vector space). This space is called the *dual space* of $\mathcal{X}$ and is denoted $\mathcal{X}^*$. It is not yet obvious that $\mathcal{X}^*$ need be non-trivial (that is, that there are any bounded linear functionals on $\mathcal{X}$ besides 0). One corollary of the Hahn-Banach theorem is there exist enough bounded linear functionals on $\mathcal{X}$ to separate points.

21.1. Examples. For each of the sequence spaces c_0, ℓ^1, ℓ^∞ , for each n the map $f \rightarrow f(n)$ is a bounded linear functional. If we fix $g \in \ell^1$, then the functional $L_g : c_0 \rightarrow \mathbb{K}$ defined by

$$L_g(f) := \sum_{n=0}^{\infty} f(n)g(n)$$

is bounded, since

$$|L_g(f)| \leq \sum_{n=0}^{\infty} |f(n)g(n)| \leq \|f\|_{\infty} \sum_{n=0}^{\infty} |g(n)| = \|g\|_1 \|f\|_{\infty}.$$

This inequality shows that $\|L_g\| \leq \|g\|_1$. In fact, equality holds, and every bounded linear functional on c_0 is of this form:

Proposition 21.1. *The map $\Phi : \ell^1 \rightarrow c_0^*$ defined by $\Phi(g) = L_g$ is an isometric isomorphism from ℓ^1 onto the dual space c_0^* .*

Proof. We have already seen that each $g \in \ell^1$ gives rise to a bounded linear functional $L_g \in c_0^*$ via

$$L_g(f) := \sum_{n=0}^{\infty} g(n)f(n)$$

and that $\|L_g\| \leq \|g\|_1$. We will prove simultaneously that this map is onto and that $\|L_g\| \geq \|g\|_1$.

Let $L \in c_0^*$. We will first show that there is unique $g \in \ell^1$ so that $L = L_g$. Let $e_n \in c_0$ be the indicator function of n , that is

$$e_n(m) = \delta_{nm}.$$

Define a function $g : \mathbb{N} \rightarrow \mathbb{K}$ by

$$g(n) = L(e_n).$$

We claim that $g \in \ell^1$ and $L = L_g$. To see this, fix an integer N and let $h \in c_{00}$ be the function

$$h(n) = \begin{cases} \overline{g(n)}/|g(n)| & \text{if } n \leq N \text{ and } g(n) \neq 0 \\ 0 & \text{otherwise.} \end{cases}$$

By definition $h \in c_{00}$ and $\|h\|_\infty \leq 1$. Note that $h = \sum_{n=0}^N h(n)e_n$. Now

$$\sum_{n=0}^N |g(n)| = \sum_{n=0}^N h(n)g(n) = L(h) = |L(h)| \leq \|L\| \|h\| \leq \|L\|.$$

It follows that $g \in \ell^1$ and $\|g\|_1 \leq \|L\|$. Moreover, the same calculation shows that $L = L_g$ when restricted to c_{00} , so by the uniqueness of extensions of bounded operators, $L = L_g$. Thus the map $g \rightarrow L_g$ is onto and

$$\|g\|_1 \leq \|L\| = \|L_g\| \leq \|g\|_1.$$

□

Proposition 21.2. $(\ell^1)^*$ is isometrically isomorphic to ℓ^∞ .

Proof. The proof follows the same lines as the proof of the previous proposition; the details are left as an exercise. □

The same mapping $g \rightarrow L_g$ also shows that every $g \in \ell^1$ gives a bounded linear functional on ℓ^∞ , but it turns out these do not exhaust $(\ell^\infty)^*$ (see Problem 21.12).

If $f \in L^1(m)$ and g is a bounded measurable function with $\sup_{x \in X} |g(x)| = M$, then the map

$$L_g(f) := \int_X fg \, dm$$

is a bounded linear functional of norm at most M . We will prove in Section ?? that the norm is in fact equal to M , and every bounded linear functional on $L^1(m)$ is of this type (at least when m is σ -finite).

If X is a compact metric space and μ is a finite, signed Borel measure on X , then

$$L_\mu(f) := \int_X f \, d\mu$$

is a bounded linear functional on $C_{\mathbb{C}}(X)$ with norm $\|\mu\| = |\mu|(X)$ (see Problem 21.8). A version of the Riesz Markov Theorem says the converse is true too.

Theorem 21.3 (Riesz-Markov). *Suppose X is a compact Hausdorff space. If $\lambda \in C(X)^*$, then there exists a unique regular Borel measure σ such that, for $f \in C(X)$,*

$$\lambda(f) = \int_X f \, d\sigma.$$

The result is true with both real and complex scalars. Focusing on the case of real scalars, the strategy is to write λ as the difference of positive linear functionals on $C(X)$ and apply the Riesz-Markov Theorem (twice). (For complex scalars, write λ in terms of its real and imaginary parts and apply the result in the real case (twice)).

A function $f \in C(X)$ is *positive* (really nonnegative) if $f(x) \geq 0$ for all $x \in X$, written $f \geq 0$. Let $C(X)^+$ denote the positive elements of $C(X)$. Given linear functionals $\lambda, \rho \in C(X)^*$, the inequality $\lambda \leq \rho$ means that $\lambda(f) \leq \rho(f)$ for all $f \in C(X)^+$.

21.2. The Hahn-Banach Extension Theorem. To state and prove the Hahn-Banach Extension Theorem, we first work in the setting $\mathbb{K} = \mathbb{R}$, then extend the results to the complex case.

Definition 21.4. Let $\mathcal{X}$ be a real vector space. A *Minkowski functional* is a function $p : \mathcal{X} \rightarrow \mathbb{R}$ such that $p(x + y) \leq p(x) + p(y)$ and $p(\lambda x) = \lambda p(x)$ for all $x, y \in \mathcal{X}$ and nonnegative $\lambda \in \mathbb{R}$. $\triangleleft$

For example, if $L : \mathcal{X} \rightarrow \mathbb{R}$ is any linear functional, then the function $p : \mathcal{X} \rightarrow \mathbb{R}$ defined by $p(x) := |L(x)|$ is a Minkowski functional. More generally if $\|\cdot\|$ is a seminorm on $\mathcal{X}$, then $p : \mathcal{X} \rightarrow \mathbb{R}$ defined by $p(x) = \|x\|$ is a Minkowski functional.

Theorem 21.5 (The Hahn-Banach Theorem, real version). *Let $\mathcal{X}$ be a vector space over $\mathbb{R}$, p a Minkowski functional on $\mathcal{X}$, and $\mathcal{M}$ a subspace of $\mathcal{X}$. If L a linear functional on $\mathcal{M}$ such that $L(x) \leq p(x)$ for all $x \in \mathcal{M}$, then there exists a linear functional L' on $\mathcal{X}$ such that*

- (i) $L'|_{\mathcal{M}} = L$ (L' extends L)
- (ii) $L'(x) \leq p(x)$ for all $x \in \mathcal{X}$ (L' is dominated by p).

The proof will invoke *Zorn's Lemma*, a result that is equivalent to the axiom of choice (as well as the well ordering principle and the Hausdorff maximality principle). A *partial order* $\preceq$ on a set is a relation that is reflexive, symmetric and transitive; that is, for all $x, y, z \in \mathcal{S}$

- (i) $x \preceq x$,
- (ii) if $x \preceq y$ and $y \preceq x$, then $x = y$, and
- (iii) if $x \preceq y$ and $y \preceq z$, then $x \preceq z$.

We call S , or more precisely, $(S, \preceq)$ a *partially ordered set* or *poset*. A subset T of S is *totally ordered*, if for each $x, y \in T$ either $x \preceq y$ or $y \preceq x$. A totally ordered subset T is often called a *chain*. An upper bound z for a chain T is an element $z \in S$ such that $t \preceq z$ for all $t \in T$. A maximal element for S is $w \in S$ that has no successor; that is there does not exist an $s \in S$ such that $s \neq w$ and $w \preceq s$.

Theorem 21.6 (Zorn's Lemma). *Suppose S is a partially ordered set. If every chain in S has an upper bound, then S has a maximal element.*

Proof of Theorem 21.5. The idea is to show that the extension can be done one dimension at a time and then infer the existence of an extension to the whole space by appeal to Zorn's lemma. We may of course assume $\mathcal{M} \neq \mathcal{X}$. So, fix a vector $x \in \mathcal{X} \setminus \mathcal{M}$ and consider the subspace $\mathcal{M} + \mathbb{R}x \subset \mathcal{X}$. For any $m_1, m_2 \in \mathcal{M}$, by hypothesis,

$$L(m_1) + L(m_2) = L(m_1 + m_2) \leq p(m_1 + m_2) \leq p(m_1 - x) + p(m_2 + x).$$

Rearranging gives, for $m_1, m_2 \in \mathcal{M}$,

$$L(m_1) - p(m_1 - x) \leq p(m_2 + x) - L(m_2)$$

and thus

$$\sup_{m \in \mathcal{M}} \{L(m) - p(m - x)\} \leq \inf_{m \in \mathcal{M}} \{p(m + x) - L(m)\}.$$

Now choose any real number λ satisfying

$$\sup_{m \in \mathcal{M}} \{L(m) - p(m - x)\} \leq \lambda \leq \inf_{m \in \mathcal{M}} \{p(m + x) - L(m)\}.$$

In particular, for $m \in \mathcal{M}$,

$$\begin{aligned} L(m) - \lambda &\leq p(m - x) \\ L(m) + \lambda &\leq p(m + x). \end{aligned} \tag{5}$$

Let $\mathcal{N} = \mathcal{M} + \mathbb{R}x$ and define $L' : \mathcal{N} \rightarrow \mathbb{R}$ by $L'(m + tx) = L(m) + t\lambda$ for $m \in \mathcal{M}$ and $t \in \mathbb{R}$. Thus L' is linear and agrees with L on $\mathcal{M}$ by definition. We now check that $L'(y) \leq p(y)$ for all $y \in \mathcal{M} + \mathbb{R}x$. Accordingly, suppose $m \in \mathcal{M}$, $t \in \mathbb{R}$ and let $y = m + tx$. If $t = 0$ there is nothing to prove. If $t > 0$, then, in view of equation (5),

$$L'(y) = L'(m + tx) = t \left(L\left(\frac{m}{t}\right) + \lambda \right) \leq t p\left(\frac{m}{t} + x\right) = p(m + tx) = p(y)$$

and a similar estimate shows that $L'(m + tx) \leq p(m + tx)$ for $t < 0$.

We have thus successfully extended L to $\mathcal{M} + \mathbb{R}x$. To finish the proof, let $\mathcal{L}$ denote the set of pairs $(L', \mathcal{N})$ where $\mathcal{N}$ is a subspace of $\mathcal{X}$ containing $\mathcal{M}$, and L' is an extension of L to $\mathcal{N}$ obeying $L'(y) \leq p(y)$ on $\mathcal{N}$. Declare $(L'_1, \mathcal{N}_1) \preceq (L'_2, \mathcal{N}_2)$ if $\mathcal{N}_1 \subset \mathcal{N}_2$ and $L'_2|_{\mathcal{N}_1} = L'_1$. This relation $\preceq$ is a partial order on $\mathcal{L}$. An exercise shows, given any increasing chain $(L'_\alpha, \mathcal{N}_\alpha)$ in $\mathcal{L}$, it has as an upper bound $(L', \mathcal{N})$ in $\mathcal{L}$, where $\mathcal{N} := \bigcup_\alpha \mathcal{N}_\alpha$ and $L(n_\alpha) := L'_\alpha(n_\alpha)$ for $n_\alpha \in \mathcal{N}_\alpha$. By Zorn's lemma the collection $\mathcal{L}$ has a maximal element $(L', \mathcal{N})$ with respect to the order $\preceq$. Since it is always possible to extend to a strictly larger subspace, the maximal element must have $\mathcal{N} = \mathcal{X}$, and the proof is finished. $\square$

The proof is a typical application of Zorn's lemma - one knows how to carry out a construction one step at a time, but there is no clear way to do it all at once.

In the special case that p is a seminorm, since $L(-x) = -L(x)$ and $p(-x) = p(x)$ the inequality $L \leq p$ is equivalent to $|L| \leq p$.

Corollary 21.7. *Suppose $\mathcal{X}$ is a normed vector space over $\mathbb{R}$, $\mathcal{M}$ is a subspace, and L is a bounded linear functional on $\mathcal{M}$. If $C \geq 0$ and $|L(x)| \leq C\|x\|$ for all $x \in \mathcal{M}$, then there exists a bounded linear functional L' on $\mathcal{X}$ extending L such that $\|L'\| \leq C$.*

Proof. Apply the Hahn-Banach theorem with the Minkowski functional $p(x) = C\|x\|$. $\square$

Before obtaining further corollaries, we extend these results to the complex case. First, if $\mathcal{X}$ is a vector space over $\mathbb{C}$, then trivially it is also a vector space over $\mathbb{R}$, and there is a simple relationship between the $\mathbb{R}$ - and $\mathbb{C}$ -linear functionals.

Proposition 21.8. *Let $\mathcal{X}$ be a vector space over $\mathbb{C}$. If $L : \mathcal{X} \rightarrow \mathbb{C}$ is a $\mathbb{C}$ -linear functional, then $u(x) = \operatorname{Re} L(x)$ defines an $\mathbb{R}$ -linear functional on $\mathcal{X}$ and $L(x) = u(x) - iu(ix)$. Conversely, if $u : \mathcal{X} \rightarrow \mathbb{R}$ is $\mathbb{R}$ -linear then $L(x) := u(x) - iu(ix)$ is $\mathbb{C}$ -linear. If in addition $p : \mathcal{X} \rightarrow \mathbb{R}$ is a seminorm, then $|u(x)| \leq p(x)$ for all $x \in \mathcal{X}$ if and only if $|L(x)| \leq p(x)$ for all $x \in \mathcal{X}$.*

Proof. Problem 21.5. □

Theorem 21.9 (The Hahn-Banach Theorem, complex version). *Let $\mathcal{X}$ be a vector space over $\mathbb{C}$, p a seminorm on $\mathcal{X}$, and $\mathcal{M}$ a subspace of $\mathcal{X}$. If $L : \mathcal{M} \rightarrow \mathbb{C}$ is a $\mathbb{C}$ -linear functional satisfying $|L(x)| \leq p(x)$ for all $x \in \mathcal{M}$, then there exists a $\mathbb{C}$ -linear functional $L' : \mathcal{X} \rightarrow \mathbb{C}$ such that*

- (i) $L'|_{\mathcal{M}} = L$ and
- (ii) $|L'(x)| \leq p(x)$ for all $x \in \mathcal{X}$.

Proof. The proof consists of applying the real Hahn-Banach theorem to extend the $\mathbb{R}$ -linear functional $u = \operatorname{Re} L$ to a functional $u' : \mathcal{X} \rightarrow \mathbb{R}$ and then defining L' from u' as in Proposition 21.8. The details are left as an exercise. □

The following corollaries are quite important, and when the Hahn-Banach theorem is applied it is usually in one of the following forms:

Corollary 21.10. *Let $\mathcal{X}$ be a normed vector space.*

- (i) *If $\mathcal{M} \subset \mathcal{X}$ is a subspace and $L : \mathcal{M} \rightarrow \mathbb{K}$ is a bounded linear functional, then there exists a bounded linear functional $L' : \mathcal{X} \rightarrow \mathbb{K}$ such that $L'|_{\mathcal{M}} = L$ and $\|L'\| = \|L\|$.*
- (ii) (Linear functionals detect norms) *If $x \in \mathcal{X}$ is nonzero, there exists $L \in \mathcal{X}^*$ with $\|L\| = 1$ such that $L(x) = \|x\|$.*
- (iii) (Linear functionals separate points) *If $x \neq y$ in $\mathcal{X}$, there exists $L \in \mathcal{X}^*$ such that $L(x) \neq L(y)$.*
- (iv) (Linear functionals detect distance to subspaces) *If $\mathcal{M} \subset \mathcal{X}$ is a closed subspace and $x \in \mathcal{X} \setminus \mathcal{M}$, there exists $L \in \mathcal{X}^*$ such that*
 - (a) $L|_{\mathcal{M}} = 0$;
 - (b) $\|L\| = 1$; and
 - (c) $L(x) = \operatorname{dist}(x, \mathcal{M}) = \inf_{y \in \mathcal{M}} \|x - y\| > 0$.

Proof. (i): Consider the (semi)norm $p(x) = \|L\| \|x\|$. By construction, $|L(x)| \leq p(x)$ for $x \in \mathcal{M}$. Hence, there is a linear functional L' on $\mathcal{X}$ such that $L'|_{\mathcal{M}} = L$ and $|L'(x)| \leq p(x)$ for all $x \in \mathcal{X}$. In particular, $\|L'\| \leq \|L\|$. On the other hand, $\|L'\| \geq \|L\|$ since L' agrees with L on $\mathcal{M}$.

(ii): Let $\mathcal{M}$ be the one-dimensional subspace of $\mathcal{X}$ spanned by x . Define a functional $L : \mathcal{M} \rightarrow \mathbb{K}$ by $L(t \frac{x}{\|x\|}) = t$. In particular, $|L(y)| = \|y\|$ for $y \in \mathcal{M}$ and thus $\|L\| = 1$. By (i), the functional L extends to a functional (still denoted L) on $\mathcal{X}$ such that $\|L\| = 1$.

(iii): Apply (ii) to the vector $x - y$.

(iv): Let $\delta = \text{dist}(x, \mathcal{M})$. Since $\mathcal{M}$ is closed, $\delta > 0$. Define a functional $L : \mathcal{M} + \mathbb{K}x \rightarrow \mathbb{K}$ by $L(y + tx) = t\delta$. Since for $t \neq 0$ and $y \in \mathcal{M}$,

$$\|y + tx\| = |t| \|t^{-1}y + x\| \geq |t|\delta = |L(y + tx)|,$$

by Hahn-Banach we can extend L to a functional $L \in \mathcal{X}^*$ with $\|L\| \leq 1$. $\square$

Needless to say, the proof of the Hahn-Banach theorem is thoroughly non-constructive, and in general it is an important (and often difficult) problem, given a normed space $\mathcal{X}$, to find some concrete description of the dual space $\mathcal{X}^*$. Usually doing so means finding a Banach space $\mathcal{Y}$ and a bounded (or, better, isometric) isomorphism $T : \mathcal{Y} \rightarrow \mathcal{X}^*$.

Note that since $\mathcal{X}^*$ is a normed space, we can form its dual, denoted $\mathcal{X}^{**}$, and called the *bidual* or *double dual* of $\mathcal{X}$. There is a canonical relationship between $\mathcal{X}$ and $\mathcal{X}^{**}$. Each fixed $x \in \mathcal{X}$ gives rise to a linear functional $\hat{x} : \mathcal{X}^* \rightarrow \mathbb{K}$ via evaluation,

$$\hat{x}(L) := L(x).$$

Since $|\hat{x}(L)| = |L(x)| \leq \|L\| \|x\|$, the linear functional $\hat{x}$ is in $\mathcal{X}^{**}$ and $\|\hat{x}\| \leq \|x\|$.

Corollary 21.11. (*Embedding in the bidual*) The map $x \rightarrow \hat{x}$ is an isometric linear map from $\mathcal{X}$ into $\mathcal{X}^{**}$.

Proof. First, from the definition we see that

$$|\hat{x}(L)| = |L(x)| \leq \|L\| \|x\|$$

so $\hat{x} \in \mathcal{X}^{**}$ and $\|\hat{x}\| \leq \|x\|$. It is straightforward to check (recalling that the L 's are linear) that the map $x \rightarrow \hat{x}$ is linear. Finally, to show that $\|\hat{x}\| = \|x\|$, fix a nonzero $x \in \mathcal{X}$. From Corollary 21.10(i) there exists $L \in \mathcal{X}^*$ with $\|L\| = 1$ and $L(x) = \|x\|$. But then for this x and L , we have $|\hat{x}(L)| = |L(x)| = \|x\|$ so $\|\hat{x}\| \geq \|x\|$, and the proof is complete. $\square$

Definition 21.12. A Banach space $\mathcal{X}$ is called *reflexive* if the map $\hat{} : \mathcal{X} \rightarrow \mathcal{X}^{**}$ is surjective. $\triangleleft$

In other words, $\mathcal{X}$ is reflexive if the map $\hat{}$ is an (isometric) isomorphism of $\mathcal{X}$ with $\mathcal{X}^{**}$. For example, every finite dimensional Banach space is reflexive (Problem 21.6). Reflexive spaces often have nice properties. For instance, the distance from a point to a (closed) subspace is attained. On the other hand, by Propositions 21.1 and 21.2, c_0^{**} is isometrically isomorphic to ℓ^∞ . In Problem 21.7 you will show that c_0 is not isometrically isomorphic to ℓ^∞ and so c_0 is not reflexive. After we have studied the L^p and ℓ^p spaces in more detail, we will see that L^p is reflexive for $1 < p < \infty$.

We note in passing that if $\mathcal{X}$ is reflexive, then its dual $\mathcal{X}^*$ has a *unique predual*: that is, if $\mathcal{Y}$ is another Banach space and $\mathcal{Y}^*$ is isometrically isomorphic to $\mathcal{X}^*$, then in fact

The embedding into the bidual has many applications; one of the most basic is the following.

Proposition 21.13 (Completion of normed spaces). *If $\mathcal{X}$ is a normed vector space, then there is a Banach space $\overline{\mathcal{X}}$ and an isometric map $\iota : \mathcal{X} \rightarrow \overline{\mathcal{X}}$ such that the image $\iota(\mathcal{X})$ is dense in $\overline{\mathcal{X}}$.*

Proof. Embed $\mathcal{X}$ into $\mathcal{X}^{**}$ via the map $x \rightarrow \hat{x}$ and let $\overline{\mathcal{X}}$ be the closure of the image of $\mathcal{X}$ in $\mathcal{X}^{**}$. Since $\overline{\mathcal{X}}$ is a closed subspace of a complete space, it is complete. $\square$

The space $\overline{\mathcal{X}}$ is called the *completion* of $\mathcal{X}$. It is unique in the sense that if $\mathcal{Y}$ is another Banach space and $j : \mathcal{X} \rightarrow \mathcal{Y}$ embeds $\mathcal{X}$ isometrically as a dense subspace of $\mathcal{Y}$, then $\mathcal{Y}$ is isometrically isomorphic to $\overline{\mathcal{X}}$. The proof of this fact is left as an exercise.

21.3. Dual spaces and adjoint operators. Let $\mathcal{X}, \mathcal{Y}$ be normed spaces with duals $\mathcal{X}^*, \mathcal{Y}^*$. If $T : \mathcal{X} \rightarrow \mathcal{Y}$ is a linear transformation and $f : \mathcal{Y} \rightarrow \mathbb{K}$ is a linear functional, then $T^*f : \mathcal{X} \rightarrow \mathbb{K}$ defined by

$$(T^*f)(x) = f(Tx) \quad (6)$$

is a linear functional on $\mathcal{X}$. If T and f are both continuous (that is, bounded) then the composition T^*f is bounded, and more is true:

Theorem 21.14. *Let $T : \mathcal{X} \rightarrow \mathcal{Y}$ be a bounded linear transformation. For $f \in \mathcal{Y}^*$, define T^*f by the formula (6). Then:*

- i) T^*f belongs to $\mathcal{X}^*$, and T^* is a linear map from $\mathcal{Y}^*$ into $\mathcal{X}^*$.
- ii) $T^* : \mathcal{Y}^* \rightarrow \mathcal{X}^*$ is bounded and $\|T^*\| = \|T\|$.

Proof. Since T is assumed bounded, for a fixed $f \in \mathcal{Y}^*$ and all $x \in \mathcal{X}$

$$|T^*f(x)| = |f(Tx)| \leq \|f\| \|Tx\| \leq \|f\| \|T\| \|x\|.$$

It follows that T^*f is bounded on $\mathcal{X}$ (thus, belongs to $\mathcal{X}^*$) and

$$\|T^*f\| \leq \|f\| \|T\|. \quad (7)$$

Thus T^* maps $\mathcal{Y}^*$ into $\mathcal{X}^*$ and it is straightforward to verify that T^* is linear, which proves item (21.14). Moreover, the inequality of equation (7) also shows that T^* is bounded and $\|T^*\| \leq \|T\|$.

It remains to show $\|T^*\| \geq \|T\|$. Toward this end, let $0 < \epsilon < 1$ be given and choose $x \in \mathcal{X}$ with $\|x\| = 1$ and $\|Tx\| > (1 - \epsilon)\|T\|$. Now consider Tx . By the Hahn-Banach theorem (Corollary 21.10(i)), there exists $f \in \mathcal{Y}^*$ such that $\|f\| = 1$ and $f(Tx) = \|Tx\|$. For this f ,

$$\|T^*\| \geq \|T^*f\| \geq |T^*f(x)| = |f(Tx)| = \|Tx\| > (1 - \epsilon)\|T\|.$$

Hence, $\|T^*\| \geq (1 - \epsilon)\|T\|$. Since ϵ was arbitrary, $\|T^*\| \geq \|T\|$. $\square$

21.4. Duality for Sub and Quotient Spaces. The Hahn-Banach Theorem allows for the identification of the duals of subspaces and quotients of Banach spaces. Informally, the dual of a subspace is a quotient and the dual of a quotient is a subspace. The precise results are stated below for complex scalars, but they hold also for real scalars.

Given a (closed) subspace $\mathcal{M}$ of the Banach space $\mathcal{X}$, let π denote the map from $\mathcal{X}$ to the quotient $\mathcal{X}/\mathcal{M}$. Recall (see Problem 20.20), the quotient is a Banach space with the norm,

$$\|z\| = \inf\{\|y\| : \pi(y) = z\}.$$

In particular, if $x \in \mathcal{X}$, then

$$\|\pi(x)\| = \inf\{\|x - m\| : m \in \mathcal{M}\}.$$

It is evident from the construction that π is continuous and $\|\pi\| \leq 1$. Further, by Problem 20.18 (or see Proposition 21.15 below) if $\mathcal{M}$ is a proper (closed) subspace, then $\|\pi\| = 1$. In particular, $\pi^* : (\mathcal{X}/\mathcal{M})^* \rightarrow \mathcal{X}^*$ (defined by $\pi^*\lambda = \lambda \circ \pi$) is also continuous. Moreover, if $x \in \mathcal{M}$, then

$$\pi^*\lambda(x) = 0.$$

Let

$$\mathcal{M}^\perp = \{f \in \mathcal{X}^* : f(x) = 0 \text{ for all } x \in \mathcal{M}\}.$$

($\mathcal{M}^\perp$ is called the *annihilator* of $\mathcal{M}$ in $\mathcal{X}^*$.) Recall, given $x \in \mathcal{X}$, the element $\hat{x} \in \mathcal{X}^{**}$ is defined by $\hat{x}(\tau) = \tau(x)$, for $\tau \in \mathcal{X}^*$. In particular,

$$\mathcal{M}^\perp = \bigcap_{x \in \mathcal{M}} \ker(\hat{x})$$

and thus $\mathcal{M}^\perp$ is a closed subspace of $\mathcal{X}^*$. Further, if $\lambda \in (\mathcal{X}/\mathcal{M})^*$, then $\pi^*\lambda \in \mathcal{M}^\perp$.

Proposition 21.15 (The dual of a quotient). *The mapping $\psi : (\mathcal{X}/\mathcal{M})^* \rightarrow \mathcal{M}^\perp$ defined by*

$$\psi(\lambda) = \pi^*\lambda$$

is an isometric isomorphism; i.e., the mapping $\pi^ : (\mathcal{X}/\mathcal{M})^* \rightarrow \mathcal{X}^*$ is an isometric isomorphism onto $\mathcal{M}^\perp$.*

Informally, the proposition is expressed as $(\mathcal{X}/\mathcal{M})^* = \mathcal{M}^\perp$.

Proof. The linearity of ψ follows from Theorem 21.14 as does $\|\psi\| = \|\pi\| \leq 1$. To prove that ψ is isometric, let $\lambda \in (\mathcal{X}/\mathcal{M})^*$ be given. Automatically, $\|\psi(\lambda)\| \leq \|\lambda\|$. To prove the reverse inequality, fix $r > 1$. Let $q \in \mathcal{X}/\mathcal{M}$ with $\|q\| = 1$ be given. There exists an $x \in X$ such that $\|x\| < r$ and $\pi(x) = q$. Hence,

$$|\lambda(q)| = |\lambda(\pi(x))| = \|\psi(\lambda)(x)\| \leq \|\psi(\lambda)\| \|x\| < r \|\psi(\lambda)\|.$$

Taking the supremum over such q shows $\|\lambda\| \leq r \|\psi(\lambda)\|$. Finally, since $1 < r$ is arbitrary, $\|\lambda\| \leq \|\psi(\lambda)\|$.

To prove that ψ is onto, and complete the proof, let $\tau \in \mathcal{M}^\perp$ be given. Fix $q \in \mathcal{X}/\mathcal{M}$. If $x, y \in \mathcal{X}$ and $\pi(x) = q = \pi(y)$, then $\tau(x) = \tau(y)$. Hence, the mapping $\lambda : \mathcal{X}/\mathcal{M} \rightarrow \mathbb{C}$ defined by $\lambda(q) = \tau(x)$ is well defined. That λ is linear is left as an exercise. To see that λ is continuous, observe that

$$|\lambda(q)| = |\tau(x)| \leq \|\tau\| \|x\|,$$

for each $x \in \mathcal{X}$ such that $\pi(x) = q$. Taking the infimum over such x gives shows

$$|\lambda(q)| \leq \|\tau\| \|q\|.$$

Finally, by construction $\psi(\lambda) = \tau$. □

Since $\mathcal{M}^\perp$ is closed in $\mathcal{X}^*$, the quotient space $\mathcal{X}^*/\mathcal{M}^\perp$ is a Banach space. Let $\rho : \mathcal{X}^* \rightarrow \mathcal{X}^*/\mathcal{M}^\perp$ denote the quotient mapping. Suppose $\lambda \in \mathcal{M}^*$. By Corollary 21.10, there is an $f \in \mathcal{X}^*$ such that $f|_{\mathcal{M}} = \lambda$; that is f is a bounded extension of λ (and indeed f can be chosen such that $\|f\| = \|\lambda\|$). If f and g are two extensions of λ to bounded linear functionals on $\mathcal{X}^*$, then $f(x) - g(x) = 0$ for $x \in \mathcal{M}$. Hence $f - g \in \mathcal{M}^\perp$ or equivalently, $\rho(f) = \rho(g)$. Consequently, the mapping $\varphi : \mathcal{M}^* \rightarrow \mathcal{X}^*/\mathcal{M}^\perp$ defined by $\varphi(\lambda) = \rho(f)$ (where f is any bounded extension of λ to $\mathcal{X}$) is well defined. It is easily verified that φ is linear. Further, given $q \in \mathcal{X}^*/\mathcal{M}^\perp$, there is an $f \in \mathcal{X}^*$ such that $\rho(f) = q$. In particular, with $\lambda = f|_{\mathcal{M}}$ we have $\varphi(\lambda) = \rho(f)$. Therefore φ is onto.

Proposition 21.16 (The dual of a subspace). *The mapping $\varphi : \mathcal{M}^* \rightarrow \mathcal{X}^*/\mathcal{M}^\perp$ is an isometric isomorphism.*

Proof. It remains to show that φ is an isometry, a fact that is an easy consequence of the Hahn-Banach Theorem. Fix $\lambda \in \mathcal{M}^*$ and let $q = \varphi(\lambda)$. If f is any bounded extension of λ to $\mathcal{X}^*$, then $\|f\| \geq \|\lambda\|$. Hence,

$$\begin{aligned} \|\varphi(\lambda)\| &= \|q\| \\ &= \inf\{\|f\| : f \in \mathcal{X}^*, \rho(f) = q\} \\ &= \inf\{\|f\| : f \in \mathcal{X}^*, f|_{\mathcal{M}} = \lambda\} \\ &\geq \|\lambda\|. \end{aligned}$$

On the other hand, by the Hahn-Banach Theorem there is a bounded extension g of λ with $\|g\| = \|\lambda\|$. Thus $\|\lambda\| \leq \|q\|$. $\square$

A special case of the following useful fact was used in the proofs above. If $\mathcal{X}, \mathcal{Y}$ are vector spaces and $T : \mathcal{X} \rightarrow \mathcal{Y}$ is linear and $\mathcal{M}$ is a subspace of the kernel of T , then T induces a linear map $\tilde{T} : \mathcal{X}/\mathcal{M} \rightarrow \mathcal{Y}$. A canonical choice is $\mathcal{M} = \ker(T)$ in which case $\tilde{T}$ is one-one. If $\mathcal{X}$ is a Banach space, $\mathcal{Y}$ is a normed vector space and $\mathcal{M}$ is closed, then $\mathcal{X}/\mathcal{M}$ is a Banach space.

Lemma 21.17. *If $\mathcal{X}$ is a Banach space, $\mathcal{M}$ is a (closed) subspace, $\mathcal{Y}$ is a normed vector space and $T : \mathcal{X} \rightarrow \mathcal{Y}$ is continuous, then the mapping $\tilde{T}$ is bounded and $\|\tilde{T}\| = \|T\|$.*

Proof. Let $\pi : \mathcal{X} \rightarrow \mathcal{X}/\mathcal{M}$ denote the quotient map and observe that $\tilde{T}\pi = T$. Since the quotient map π has norm 1 (see Problem 20.20), we see that $\|\tilde{T}\| \leq \|T\|$. For the opposite inequality, let $0 < \epsilon < 1$ and choose $x \in \mathcal{X}$ such that $\|x\| = 1$ and $\|Tx\| > (1 - \epsilon)\|T\|$. Then $\|\pi(x)\| \leq 1$ and

$$\|\tilde{T}\| \geq \|\tilde{T}\pi(x)\| = \|Tx\| > (1 - \epsilon)\|T\|.$$

Letting ϵ go to zero finishes the proof. $\square$

21.5. Problems.

Problem 21.1. Prove, if $\mathcal{X}$ is any normed vector space, $\{x_1, \dots, x_n\}$ is a linearly independent set in $\mathcal{X}$, and $\alpha_1, \dots, \alpha_n$ are scalars, then there exists a bounded linear functional f

on $\mathcal{X}$ such that $f(x_j) = \alpha_j$ for $j = 1, \dots, n$. (Recall linear maps from a finite dimensional normed vector space to a normed vector space are bounded.)

Problem 21.2. Let $\mathcal{X}, \mathcal{Y}$ be normed spaces and $T : \mathcal{X} \rightarrow \mathcal{Y}$ a linear transformation. Prove T is bounded if and only if there exists a constant C such that for all $x \in \mathcal{X}$ and $f \in \mathcal{Y}^*$,

$$|f(Tx)| \leq C\|f\|\|x\|; \quad (8)$$

in which case $\|T\|$ is equal to the best possible C in (8).

Problem 21.3. Let $\mathcal{X}$ be a normed vector space. Show that if $\mathcal{M}$ is a closed subspace of $\mathcal{X}$ and $x \notin \mathcal{M}$, then $\mathcal{M} + \mathbb{K}x$ is closed. Use this result to give another proof that every finite-dimensional subspace of $\mathcal{X}$ is closed.

Problem 21.4. Prove, if $\mathcal{M}$ is a *finite-dimensional* subspace of a Banach space $\mathcal{X}$, then there exists a closed subspace $\mathcal{N} \subset \mathcal{X}$ such that $\mathcal{M} \cap \mathcal{N} = \{0\}$ and $\mathcal{M} + \mathcal{N} = \mathcal{X}$. (In other words, every $x \in \mathcal{X}$ can be written uniquely as $x = y + z$ with $y \in \mathcal{M}$, $z \in \mathcal{N}$.) *Hint:* Choose a basis $x_1, \dots, x_n$ for $\mathcal{M}$ and construct, using Problem 21.1 and the Hahn-Banach Theorem, bounded linear functionals $f_1, \dots, f_n$ on $\mathcal{X}$ such that $f_i(x_j) = \delta_{ij}$. Now let $\mathcal{N} = \bigcap_{i=1}^n \ker f_i$. (Warning: this conclusion can fail badly if $\mathcal{M}$ is not assumed finite dimensional, even if $\mathcal{M}$ is still assumed closed. Perhaps the first known example is that c_0 is not *complemented* in ℓ^∞ , though it is nontrivial to prove.)

Problem 21.5. Prove Proposition 21.8.

Problem 21.6. Prove every finite-dimensional Banach space is reflexive.

Problem 21.7. Let B denote the subset of ℓ^∞ consisting of sequences which take values in $\{-1, 1\}$. Show that any two (distinct) points of B are a distance 2 apart. Show, if C is a countable subset of ℓ^∞ , then there exists a $b \in B$ such that $\|b - c\| \geq 1$ for all $c \in C$. Conclude ℓ^∞ is not separable. Prove there is no isometric isomorphism $\Lambda : c_0 \rightarrow \ell^\infty$. As a corollary, conclude that c_0 is not reflexive. (Of course, saying $c_0 \neq \ell^\infty$ via the canonical embedding of Corollary 21.11 is much weaker than saying there is no isometric isomorphism between c_0 and ℓ^∞ .)

Problem 21.8. Prove, if μ is a finite regular (signed) Borel measure on a compact Hausdorff space, then the linear function $L_\mu : C(X) \rightarrow \mathbb{R}$ defined by

$$L_\mu(f) = \int_X f d\mu$$

is bounded (continuous) and $\|L_\mu\| = \|\mu\| := |\mu|(X)$. (See the Riesz-Markov Theorem for positive linear functionals.)

Problem 21.9. Let $\mathcal{X}$ and $\mathcal{Y}$ be normed vector spaces and $T \in L(\mathcal{X}, \mathcal{Y})$.

- Consider $T^{**} : \mathcal{X}^{**} \rightarrow \mathcal{Y}^{**}$. Identifying $\mathcal{X}, \mathcal{Y}$ with their images in $\mathcal{X}^{**}$ and $\mathcal{Y}^{**}$, show that $T^{**}|_{\mathcal{X}} = T$.
- Prove T^* is injective if and only if the range of T is dense in $\mathcal{Y}$.
- Prove that if the range of T^* is dense in $\mathcal{X}^*$, then T is injective; if $\mathcal{X}$ is reflexive then the converse is true.

- Problem 21.10.**
- a) Prove that if $\mathcal{X}$ is reflexive, then $\mathcal{X}^*$ is reflexive.
 - b) Prove that if $\mathcal{X}$ is reflexive and $\mathcal{M} \subset \mathcal{X}$ is a closed subspace, then $\mathcal{M}$ is reflexive.
 - c) Prove that a Banach space $\mathcal{X}$ is reflexive if and only if $\mathcal{X}^*$ is reflexive.
 - d) Prove that if $\mathcal{X}$ is reflexive and $\mathcal{Y}$ is another Banach space with $\mathcal{Y}^*$ isometrically isomorphic to $\mathcal{X}^*$, then $\mathcal{Y}$ is isometrically isomorphic to $\mathcal{X}$. (This conclusion can fail if $\mathcal{X}$ is not reflexive; see Problem 21.15.)

Problem 21.11. Prove, if $\mathcal{X}$ is a Banach space and $\mathcal{X}^*$ is separable, then $\mathcal{X}$ is separable. [Hint: let $\{f_n\}$ be a countable dense subset of $\mathcal{X}^*$. For each n choose x_n such that $\|x_n\| = 1$ and $|f_n(x_n)| \geq \frac{1}{2}\|f_n\|$. Show that the set of $\mathbb{Q}$ -linear combinations of $\{x_n\}$ is dense in $\mathcal{X}$.]

- Problem 21.12.**
- a) Prove there exists a bounded linear functional $L \in (\ell^\infty)^*$ with the following property: whenever $f \in \ell^\infty$ and $\lim_{n \rightarrow \infty} f(n)$ exists, then $L(f)$ is equal to this limit. (Hint: first show that the set of such f forms a subspace $\mathcal{M} \subset \ell^\infty$).
 - b) Show that such a functional L is not equal to L_g for any $g \in \ell^1$; thus the map $T : \ell^1 \rightarrow (\ell^\infty)^*$ given by $T(g) = L_g$ is not surjective.
 - c) Give another proof that T is not surjective, using Problem 21.11.

Problem 21.13. Let $\mathcal{X}$ be a normed space and let $K \subset \mathcal{X}$ be a convex set. (Recall, this means that whenever $x, y \in K$, then $\frac{1}{2}(x + y) \in K$; equivalently, $tx + (1 - t)y \in K$ for all $0 \leq t \leq 1$.) A point $x \in K$ is called an *extreme point* of K whenever $y, z \in K$, $0 < t < 1$, and $x = ty + (1 - t)z$, then $y = z = x$. (That is, the only way to write x as a convex combination of elements of K is the trivial way.)

- a) Let $\mathcal{X}$ be a normed space and let $B = \text{ball}(\mathcal{X})$ denote the (closed) unit ball of $\mathcal{X}$. Prove that $x \in B$ is *not* an extreme point of B if and only if there exists a nonzero $y \in B$ such that $\|x \pm y\| \leq 1$.
- b) Prove that if $\mathcal{X}$ and $\mathcal{Y}$ are normed spaces, and $T : \mathcal{X} \rightarrow \mathcal{Y}$ is a surjective linear isometry, (so that $\mathcal{X}$ and $\mathcal{Y}$ are isometrically isomorphic) then T induces a bijection between the extreme points of $\text{ball}(\mathcal{X})$ and $\text{ball}(\mathcal{Y})$.
- c) Let ℓ_n^p denote the (real) Banach space $\mathbb{R}^n$ equipped with the ℓ^p norm, $1 \leq p \leq \infty$. Prove that ℓ_2^1 and ℓ_2^∞ are isometrically isomorphic, but that there is *no* isometry between ℓ_3^1 and ℓ_3^∞ .

- Problem 21.14.**
- a) Show that the extreme points of the unit ball of ℓ^1 are precisely the points of the form λe_n where $|\lambda| = 1$ and e_n is the sequence which is 1 in the n^{th} entry and 0 elsewhere. (See Problem 21.13).
 - b) Determine the extreme points of the unit ball of ℓ^∞ .
 - c) Show that the unit ball of c_0 has *no* extreme points.

Problem 21.15. Let

$$c = \left\{ f : \mathbb{N} \rightarrow \mathbb{K} \mid \lim_{n \rightarrow \infty} f(n) \text{ exists} \right\},$$

and equip c with the supremum norm $\|f\|_\infty := \sup |f(n)|$.

- a) Show that $c^* \cong \ell^1$ isometrically.
- b) Prove that c is boundedly isomorphic to c_0 .
- c) Prove that c is not *isometrically* isomorphic to c_0 . (Hint: examine the extreme points of the unit balls of c and c_0 ; see Problems 21.13 and 21.14.)

(This problem provides an example of Banach spaces $\mathcal{X}$ and $\mathcal{Y}$ such that $\mathcal{X}$ and $\mathcal{Y}$ are not isometrically isomorphic, but $\mathcal{X}^*$ and $\mathcal{Y}^*$ are. So in general we cannot recover $\mathcal{X}$ (isometrically) from $\mathcal{X}^*$. In fact the situation is worse, ℓ^1 has isometric preduals which are not even boundedly isomorphic to c_0 , but the construction is more involved and outside the scope of these notes.)

22. THE BAIRE CATEGORY THEOREM AND APPLICATIONS

Recall, a set D in a metric space X is *dense* if $\overline{D} = X$. Thus, D is dense if and only if D^c does not contain a nonempty open set, if and only if D has nontrivial intersection with every nonempty open set. A topological space X is called a *Baire space* if it has the following property: if $(U_n)_{n=1}^\infty$ is a countable sequence of open dense subsets of X , then the intersection $\cap_{n=1}^\infty U_n$ is dense in X .

Theorem 22.1 (The Baire Category Theorem). *Every complete metric space X is a Baire space. In other words, if X is a complete metric space and if $(U_n)_{n=1}^\infty$ is a sequence of open dense subsets of X , then $\cap_{n=1}^\infty U_n$ is dense in X .*

Theorem 22.1 is true if X is a locally compact Hausdorff space and there are connections between the Baire Category Theorem and the axiom of choice.

A subset $E \subset X$ is *nowhere dense* if its closure has empty interior. Equivalently, $\overline{E}^c$ is open and dense. A set F in a metric space X is *first category* (or *meager*) if it can be expressed as the countable union of nowhere dense sets. In particular, a countable union of first category sets is first category. A set G is *second category* if it is not first category. For applications, the following corollary often suffices.

Corollary 22.2. *If X is a complete metric space, then X is not a countable union of nowhere dense sets; i.e., X is of second category in itself.*

Proof. Take complements and apply Theorem 22.1. □

Thus, the Baire property is used as a kind of pigeonhole principle: the “thick” Baire space X cannot be expressed as a countable union of the “thin” nowhere dense sets E_n . Equivalently, if X is Baire and $X = \bigcup_n E_n$, then at least one of the E_n is somewhere dense.

The following lemma should be familiar from advanced calculus.

Lemma 22.3. *Let X be a complete metric space and suppose (C_n) is a sequence of subsets of X . If*

- (i) *each C_n is nonempty;*
- (ii) *(C_n) is nested decreasing;*

- (iii) each C_n is closed; and
- (iv) $(\text{diam}(C_n))$ converges to 0,

then there is an $x \in X$ such that

$$\{x\} = \bigcap C_n.$$

Moreover, if $x_n \in C_n$, then (x_n) converges to some x .

Proof of Theorem 22.1. Let $(U_n)_{n=1}^\infty$ be a sequence of open dense sets in X and let $I = \bigcap U_n$. To prove I is dense, it suffices to show that I has nontrivial intersection with every nonempty open set W . Fix such a W . Since U_1 is dense, there is a point $x_1 \in W \cap U_1$. Since U_1 and W are open, there is a radius $0 < r_1 < 1$ such that the $\overline{B(x_1, r_1)}$ is contained in $W \cap U_1$. Similarly, since U_2 is dense and open there is a point $x_2 \in \overline{B(x_1, r_1)} \cap U_2$ and a radius $0 < r_2 < \frac{1}{2}$ such that

$$\overline{B(x_2, r_2)} \subset B(x_1, r_1) \cap U_2 \subset W \cap U_1 \cap U_2.$$

Continuing inductively, since each U_n is dense and open there is a sequence of points $(x_n)_{n=1}^\infty$ and radii $0 < r_n < \frac{1}{n}$ such that

$$\overline{B(x_n, r_n)} \subset B(x_{n-1}, r_{n-1}) \cap U_n \subset W \cap (\bigcap_{j=1}^n U_j).$$

The sequence of sets $(\overline{B(x_n, r_n)})$ satisfies the hypothesis of Lemma 22.3 and X is complete. Hence there is an $x \in X$ such that

$$x \in \bigcap_n \overline{B(x_n, r_n)} \subset W \cap I.$$

□

We now give three important applications of the Baire category theorem in functional analysis. These are the Principle of Uniform boundedness (also known as the Banach-Steinhaus theorem), the Open Mapping Theorem, and the Closed Graph Theorem. (In learning these theorems, keep careful track of what completeness hypotheses are needed.)

Theorem 22.4 (The Principle of Uniform Boundedness (PUB)). *Suppose $\mathcal{X}, \mathcal{Y}$ are normed spaces and $\{T_\alpha : \alpha \in A\} \subset B(\mathcal{X}, \mathcal{Y})$ is a collection of bounded linear transformations from $\mathcal{X}$ to $\mathcal{Y}$. Let B denote the set*

$$B := \{x \in X : M(x) := \sup_\alpha \|T_\alpha x\| < \infty\}. \quad (9)$$

If B is of the second category (thus not a countable union of nowhere dense sets) in X , then

$$\sup_\alpha \|T_\alpha\| < \infty.$$

In particular, if $\mathcal{X}$ is complete and if the collection $\{T_\alpha : \alpha \in A\}$ is pointwise bounded, then it is uniformly bounded.

Proof. For each integer $n \geq 1$ consider the set

$$V_n := \{x \in \mathcal{X} : M(x) > n\}.$$

Since each T_α is bounded, the sets V_n are open. (Indeed, for each α the map $x \rightarrow \|T_\alpha x\|$ is continuous from $\mathcal{X}$ to $\mathbb{R}$, so if $\|T_\alpha x\| > n$ for some α then also $\|T_\alpha y\| > n$ for all y sufficiently close to x .) Let E_n denote the complement of V_n and observe that $B = \bigcup_{n=1}^{\infty} E_n$. Since B is assumed to be of the second category, there is an N such that $(\bar{E}_N)^\circ$ is not empty. Since E_N is closed, it follows that E_N has nonempty interior; i.e., there is an $x_0 \in E_N$ and $r > 0$ so that $x_0 - x \in E_N$ for all $\|x\| < r$. Thus, for every α and every $\|x\| < r$,

$$\|T_\alpha x\| \leq \|T_\alpha(x - x_0)\| + \|T_\alpha x_0\| \leq N + N.$$

That is, if $\|x\| < r$, then $M(x) \leq 2N$. By rescaling we conclude that if $\|x\| < 1$, then $\|T_\alpha x\| \leq 2N/r$ for all α and thus $\sup_\alpha \|T_\alpha\| \leq 2N/r < \infty$. $\square$

Given a subset B of a vector space $\mathcal{X}$ and a scalar $s \in \mathbb{K}$, let $sB = \{sb : b \in B\}$. Similarly, for $x \in \mathcal{X}$, let $B - x = \{b - x : b \in B\}$. Let $\mathcal{X}, \mathcal{Y}$ be normed vector spaces and suppose $T : \mathcal{X} \rightarrow \mathcal{Y}$ is linear. If $B \subset \mathcal{X}$ and $s \in \mathbb{K}$ is nonzero, then $T(sB) = sT(B)$ and further, an easy argument shows $\overline{T(sB)} = s\overline{T(B)}$. It is also immediate that if B is open, then so is $B - x$.

Recall that if X, Y are topological spaces, a mapping $f : X \rightarrow Y$ is called *open* if $f(U)$ is open in Y whenever U is open in X . In particular, if f is a bijection, then f is open if and only if f^{-1} is continuous. In the case of normed linear spaces the condition that a linear map be open can be refined somewhat.

Lemma 22.5 (Translation and Dilation lemma). *Let $\mathcal{X}, \mathcal{Y}$ be normed vector spaces, let B denote the open unit ball of $\mathcal{X}$, and let $T : \mathcal{X} \rightarrow \mathcal{Y}$ be a linear map. The following are equivalent.*

- (i) *The map T is open;*
- (ii) *$T(B)$ contains an open ball centered to 0;*
- (iii) *there is an $s > 0$ such that $T(sB)$ contains an open ball centered to 0; and*
- (iv) *$T(sB)$ contains an open ball centered to 0 for each $s > 0$.*

Proof. This result is more or less immediate from the fact, for fixed z_0 and $r \in \mathbb{K}$, that the translation map $z \rightarrow z + z_0$ and the dilation map $z \rightarrow rz$ are continuous in a normed vector space. The implication (i) implies (ii) is immediate. The fact that $T(sB) = sT(B)$ for $s > 0$ readily shows (ii), (iii) and (iv) are equivalent.

To finish the proof it suffice to show (iv) implies (i). Accordingly, suppose (iv) holds and let $U \subset X$ be a given open set. To prove that $T(U)$ is open, let $y \in T(U)$ be given. There is an $x \in U$ such that $T(x) = y$. There is an $s > 0$ such that the ball $B(x, s)$ lies in U ; that is $B(x, s) \subset U$. The ball $sB = B(0, s) = B(x, r) - x$ is an open ball centered to 0. By hypothesis there is an $r > 0$ such that $B^\mathcal{Y}(0, r) \subset T(B(0, s))$. (Here we use $B^\mathcal{Y}$

to emphasize this ball is a subset of $\mathcal{Y}$.) By linearity of T ,

$$\begin{aligned} B^{\mathcal{Y}}(y, r) &= B^{\mathcal{Y}}(0, r) + y \subset T(B(0, s)) + y \\ &= T(B(0, s)) + T(x) = T(B(0, s) + x) = T(B(x, s)) \subset T(U). \end{aligned}$$

Thus $T(U)$ is open. $\square$

Theorem 22.6 (Open Mapping). *Suppose that $\mathcal{X}$ is a Banach space, $\mathcal{Y}$ is a normed vector space and $T : \mathcal{X} \rightarrow \mathcal{Y}$ is bounded. If the range of T is of second category, then*

- (i) $T(\mathcal{X}) = \mathcal{Y}$;
- (ii) $\mathcal{Y}$ is complete (so a Banach space); and
- (iii) T is open.

In particular, if $\mathcal{X}, \mathcal{Y}$ are Banach spaces, and $T : \mathcal{X} \rightarrow \mathcal{Y}$ is bounded and onto, then T is an open map.

Proof. Observe that (i) follows immediately from (iii). To prove (iii), let $B(x, r)$ denote the open ball of radius r centered at x in $\mathcal{X}$. Trivially $\mathcal{X} = \bigcup_{n=1}^{\infty} B(0, n)$ and thus $T(\mathcal{X}) = \bigcup_{n=1}^{\infty} T(B(0, n))$. Since the range of T is assumed second category, there is an N such that $T(B(0, N))$ is second category and hence not nowhere dense. In other words, $\overline{T(B(0, N))}$ has nonempty interior. By scaling (see Lemma 22.5), $\overline{T(B(0, 1))}$ has nonempty interior. Hence, there exists $p \in \mathcal{Y}$ and $r > 0$ such that $\overline{T(B(0, 1))}$ contains the open ball $B^{\mathcal{Y}}(p, r)$. (Here the superscript $\mathcal{Y}$ is used to emphasize this ball is in $\mathcal{Y}$.) It follows that for all $\|y\| < r$,

$$y = -p + (y + p) \in \overline{T(B(0, 2))}.$$

In other words,

$$B^{\mathcal{Y}}(0, r) \subset \overline{T(B(0, 2))}.$$

By scaling, it follows that, for $n \in \mathbb{N}$,

$$B^{\mathcal{Y}}(0, \frac{r}{2^{n+1}}) \subset \overline{T(B(0, \frac{1}{2^n}))}.$$

We will use the hypothesis that $\mathcal{X}$ is complete to prove $B^{\mathcal{Y}}(0, \frac{r}{4}) \subset T(B(0, 1))$. Accordingly let y such that $\|y\| < \frac{r}{4}$ be given. Since y is in the closure of $T(B(0, \frac{1}{2}))$, there is a $y_1 \in T(B(0, \frac{1}{2}))$ such that $\|y - y_1\| < \frac{r}{8}$. Since $y - y_1 \in B^{\mathcal{Y}}(0, \frac{r}{8})$ it is in the closure of $T(B(0, \frac{1}{4}))$. Thus there is a $y_2 \in T(B(0, \frac{1}{4}))$ such that $\|(y - y_1) - y_2\| < \frac{r}{16}$. Continuing in this fashion produces a sequence $(y_j)_{j=1}^{\infty}$ from $\mathcal{Y}$ such that,

- (a) $\|y - \sum_{j=1}^n y_j\| \leq \frac{r}{2^{n+2}}$; and
- (b) $y_n \in T(B(0, \frac{1}{2^n}))$

for all n . It follows that $\sum_{j=1}^{\infty} y_j$ converges to y . Further, for each j there is an $x_j \in B(0, \frac{1}{2^j})$ such that $y_j = Tx_j$. Since

$$\|x\| \leq \sum_{j=1}^{\infty} \|x_j\| < \sum_{k=1}^{\infty} 2^{-k} = 1,$$

the series $\sum_{j=1}^{\infty} x_j$ converges to some $x \in B(0, 1)$. It follows that $y = Tx$ by continuity of T . Consequently $y \in T(B(0, 1))$ and the proof of (iii) is complete.

To prove (ii), let $\mathcal{M}$ denote the kernel of T and $\tilde{T}$ the mapping $\tilde{T} : \mathcal{X}/\mathcal{M} \rightarrow \mathcal{Y}$ determined by $\tilde{T}\pi = T$. By Lemma 21.17, $\tilde{T}$ is continuous and one-one. Further its range is the same as the range of T , namely $\mathcal{Y}$, and is thus second category. Hence, by what has already been proved, $\tilde{T}$ is an open map. and consequently $\tilde{T}^{-1}$ is continuous. Hence $\mathcal{X}/\mathcal{M}$ and $\mathcal{Y}$ are isomorphic (though of course not necessarily isometrically isomorphic) as normed vector spaces. Therefore, since $\mathcal{X}/\mathcal{M}$ is complete, so is $\mathcal{Y}$. $\square$

Note that the proof of item (ii) in the Open Mapping Theorem shows, in the case that in the case that T is one-one and its range is of second category, that T is onto and its inverse is continuous. In particular, if $T : \mathcal{X} \rightarrow \mathcal{Y}$ is a continuous bijection and $\mathcal{Y}$ is a Banach space (so the range of T is second category), then T^{-1} is continuous.

Corollary 22.7 (The Banach Isomorphism Theorem). *If $\mathcal{X}, \mathcal{Y}$ are Banach spaces and $T : \mathcal{X} \rightarrow \mathcal{Y}$ is a bounded bijection, then T^{-1} is also bounded (hence, T is an isomorphism).*

To state the final result of this section, we need a few more definitions. Let $\mathcal{X}, \mathcal{Y}$ be normed spaces. The Cartesian product $\mathcal{X} \times \mathcal{Y}$ is a topological space in the *product topology*. A set is open in the product topology if and only if it can be written as a union of products of open sets. Alternately, a set O is open if and only if for each $z = (x, y) \in O$ there exists open sets $U \subset \mathcal{X}$ and $V \subset \mathcal{Y}$ such that $z \in U \times V \subset O$. It is not too hard to show that $\mathcal{X} \times \mathcal{Y}$ is metrizable (in fact the product topology can be realized by norming $\mathcal{X} \times \mathcal{Y}$, e.g. with the norm $\|(x, y)\| := \max(\|x\|, \|y\|)$). It is easy to see that a sequence $z_n = (x_n, y_n)$ converges in the product topology if and only if both (x_n) and (y_n) converge. Further, if $\mathcal{X}, \mathcal{Y}$ are both Banach spaces (complete), then $\mathcal{X} \times \mathcal{Y}$ is also complete and hence a Banach space. The space $\mathcal{X} \times \mathcal{Y}$ is equipped with the coordinate projections $\pi_{\mathcal{X}}(x, y) = x, \pi_{\mathcal{Y}}(x, y) = y$. It is clear from the definition of the product topology that these maps are continuous. (In fact the product topology is the coarsest topology such that the coordinate projections are continuous.)

Given a linear map $T : \mathcal{X} \rightarrow \mathcal{Y}$, its *graph* is the set

$$G(T) := \{(x, y) \in \mathcal{X} \times \mathcal{Y} : y = Tx\}$$

Observe that since T is a linear map, $G(T)$ is a linear subspace of $\mathcal{X} \times \mathcal{Y}$. The transformation T is *closed* if $G(T)$ is a closed subset of $\mathcal{X} \times \mathcal{Y}$. It is an easy exercise to show that $G(T)$ is closed if and only if whenever (x_n, Tx_n) converges to (x, y) , we have $y = Tx$. Problem 22.2 gives an example where $G(T)$ is closed, but T is not continuous. On the other hand, if $\mathcal{X}, \mathcal{Y}$ are complete (Banach spaces), then $G(T)$ is closed if and only if T is continuous.

Theorem 22.8 (The Closed Graph Theorem). *If $\mathcal{X}, \mathcal{Y}$ are Banach spaces and $T : \mathcal{X} \rightarrow \mathcal{Y}$ is closed, then T is bounded.*

Proof. Let π_1, π_2 be the coordinate projections $\pi_{\mathcal{X}}, \pi_{\mathcal{Y}}$ restricted to $G(T)$; explicitly $\pi_1(x, Tx) = x$ and $\pi_2(x, Tx) = Tx$. Note that π_1 is a bijection between $G(T)$ and $\mathcal{X}$ and

in particular $\pi_1^{-1}(x) = (x, Tx)$. By hypothesis $G(T)$ is a closed subset of a Banach space and hence a Banach space. Thus π_1 is a bounded linear bijection between Banach spaces and therefore, by Corollary 22.7, $\pi_1^{-1} : X \rightarrow G(T)$ is bounded. Since π_2 is bounded, $\pi_2 \circ \pi_1^{-1} : X \rightarrow Y$ is continuous. To finish the proof, observe $\pi_2 \circ \pi_1^{-1}(x) = \pi_2(x, Tx) = Tx$.

□

22.1. Problems.

Problem 22.1. Show that there exists a sequence of open, dense subsets $U_n \subset \mathbb{R}$ such that $m(\bigcap_{n=1}^{\infty} U_n) = 0$.

Problem 22.2. Consider the linear subspace $\mathcal{D} \subset c_0$ defined by

$$\mathcal{D} = \{f \in c_0 : \lim_{n \rightarrow \infty} |nf(n)| = 0\}$$

and the linear transformation $T : \mathcal{D} \rightarrow c_0$ defined by $(Tf)(n) = nf(n)$.

a) Prove T is closed, but not bounded. b) Prove T is bijective and $T^{-1} : c_0 \rightarrow \mathcal{D}$ is bounded (and surjective), but not open. c) What can be said of $\mathcal{D}$ as a subset of c_0 ?

Problem 22.3. Suppose $\mathcal{X}$ is a vector space equipped with two norms $\|\cdot\|_1, \|\cdot\|_2$ such that $\|\cdot\|_1 \leq \|\cdot\|_2$. Prove that if $\mathcal{X}$ is complete in both norms, then the two norms are equivalent.

Problem 22.4. Let $\mathcal{X}, \mathcal{Y}$ be Banach spaces. Provisionally, say that a linear transformation $T : \mathcal{X} \rightarrow \mathcal{Y}$ is *weakly bounded* if $f \circ T \in \mathcal{X}^*$ whenever $f \in \mathcal{Y}^*$. Prove, if T is weakly bounded, then T is bounded.

Problem 22.5. Let $\mathcal{X}, \mathcal{Y}$ be Banach spaces. Suppose (T_n) is a sequence in $B(\mathcal{X}, \mathcal{Y})$ and $\lim_n T_n x$ exists for every $x \in \mathcal{X}$. Prove, if T is defined by $Tx = \lim_n T_n x$, then T is bounded.

Problem 22.6. Suppose that $\mathcal{X}$ is a vector space with a countably infinite basis. (That is, there is a linearly independent set $\{x_n\} \subset \mathcal{X}$ such that every vector $x \in \mathcal{X}$ is expressed uniquely as a *finite* linear combination of the x_n 's.) Prove there is no norm on $\mathcal{X}$ under which it is complete. (Hint: consider the finite-dimensional subspaces $\mathcal{X}_n := \text{span}\{x_1, \dots, x_n\}$.)

Problem 22.7. The Baire Category Theorem can be used to prove the existence of (very many!) continuous, nowhere differentiable functions on $[0, 1]$. To see this, let E_n denote the set of all functions $f \in C[0, 1]$ for which there exists $x_0 \in [0, 1]$ (which may depend on f) such that $|f(x) - f(x_0)| \leq n|x - x_0|$ for all $x \in [0, 1]$. Prove the sets E_n are nowhere dense in $C[0, 1]$; the Baire Category Theorem then shows that the set of nowhere differentiable functions is second category. (To see that E_n is nowhere dense, approximate an arbitrary continuous function f uniformly by piecewise linear functions g , whose pieces have slopes greater than $2n$ in absolute value. Any function sufficiently close to such a g will not lie in E_n .)

Problem 22.8. Let $L^2([0, 1])$ denote the Lebesgue measurable functions $f : [0, 1] \rightarrow \mathbb{C}$ such that $|f|^2$ is in $L^1([0, 1])$. It turns out, as we will see later, that $L^2([0, 1])$ is a linear manifold (subspace of the vector space $L^1([0, 1])$), though this fact is not needed for this problem.

Let $g_n : [0, 1] \rightarrow \mathbb{R}$ denote the function which takes the value n on $[0, \frac{1}{n^3}]$ and 0 elsewhere. Show,

- (i) if $f \in L^2([0, 1])$, then $\lim_{n \rightarrow \infty} \int g_n f \, dm = 0$;
- (ii) $L_n : L^1([0, 1]) \rightarrow \mathbb{C}$ defined by $L_n(f) = \int g_n f \, dm$ is bounded, and $\|L_n\| = n$;
- (iii) conclude $L^2([0, 1])$ is of the first category in $L^1([0, 1])$.

Problem 22.9. A *Banach space of functions* on a set X is a vector subspace B of the space of complex-valued functions on X with a norm $\|\cdot\|$ making B a Banach space such that, for each $x \in X$, the mapping $E_x : B \rightarrow \mathbb{C}$ defined by $E_x(f) = f(x)$ is continuous (bounded) and if $f(x) = 0$ for all $x \in X$, then $f = 0$.

Suppose $g : X \rightarrow \mathbb{C}$. Show, if $gf \in B$ for each $f \in B$, then the linear map $M_g : B \rightarrow B$ defined by $M_g f = gf$ is bounded.

my solution: By the closed graph theorem, it suffices to show that the graph of M_g is closed. To this end, suppose (f_n, gf_n) is a convergent sequence from $G(M_g)$; that is that there exist $f, h \in B$ such that (f_n, gf_n) converges to (f, g) . In particular, (f_n) converges to f and (gf_n) converges to h in B . By continuity, for $x \in X$, we have $(f_n(x) = E_x(f_n))$ converges to $f(x) = E_x(f)$. Similarly, $(g(x)f_n(x))$ converges to $h(x)$. On the other hand, $(g(x)f_n(x))$ also converges to $g(x)f(x)$. Hence $h(x) = g(x)f(x)$ and $h = M_g f$. Thus $(f, h) = (f, M_g f) \in G(M_g)$ and the proof is complete.

Problem 22.10. Suppose $\mathcal{X}$ is a Banach space and $\mathcal{M}$ and $\mathcal{N}$ are closed subspaces. Show, if for each $x \in \mathcal{X}$ there exist unique $m \in \mathcal{M}$ and $n \in \mathcal{N}$ such that

$$x = m + n,$$

then the mapping $P : \mathcal{X} \rightarrow \mathcal{M}$ defined by $Px = m$ is bounded.