

OPERATOR-VALUED HERGLOTZ KERNELS AND FUNCTIONS OF POSITIVE REAL PART ON THE BALL

MICHAEL T. JURY

ABSTRACT. We describe several classes of holomorphic functions of positive real part on the unit ball; each is characterized by an operator-valued Herglotz formula. Motivated by results of J.E. McCarthy and M. Putinar, we define a family of weighted Cauchy-Fantappiè pairings on the ball and establish duality relations between certain pairs of classes, and in particular we identify the dual of the positive Schur class. We also establish the existence of self-dual classes with respect to this pairing, and identify some extreme points of the positive Schur class.

1. INTRODUCTION

It is well-known that a function f , holomorphic in the unit disk \mathbb{D} , has nonnegative real part if and only if there exists a finite, positive Borel measure μ on the unit circle \mathbb{T} such that

$$(1.1) \quad f(z) = \int_{\mathbb{T}} \frac{1 + z\bar{\zeta}}{1 - z\bar{\zeta}} d\mu(\zeta) + i\operatorname{Im} f(0).$$

This expression is called the *Herglotz representation* of f . No such simple characterization is available in higher dimensions. On the one hand, a result of Korányi and Pukánszky [5] shows that every holomorphic function of positive real part in the unit ball $\mathbb{B}^d \subset \mathbb{C}^d$ (call this set of functions $O^+(\mathbb{B}^d)$) admits an integral representation

$$(1.2) \quad f(z) = \int_{\partial\mathbb{B}^d} \left[\frac{2}{(1 - \langle z, \zeta \rangle)^d} - 1 \right] d\mu(\zeta) + i\operatorname{Im} f(0).$$

for some positive measure μ on the sphere. However, the integral kernel does not have positive real part when $d > 1$, and so not every measure gives an f of positive real part. The class of measures that do is characterized by a family of moment conditions, but these conditions are difficult to work with. In particular very little is known about the extreme rays of O^+ .

Date: October 19, 2008.

Partially supported by NSF grant DMS-0701268.

On the other hand, one may consider the class M^+ , consisting of all functions on \mathbb{B}^d of the form

$$(1.3) \quad f(z) = \int_{\partial\mathbb{B}^d} \frac{1 + \langle z, \zeta \rangle}{1 - \langle z, \zeta \rangle} d\mu(\zeta) + i\text{Im } f(0)$$

where μ is again a positive measure on the sphere. In this case the kernel has positive real part, so every measure gives an element of O^+ , but when $d > 1$ not every $f \in O^+$ can be represented in this way. (Note that the expressions (1.1)-(1.3) all coincide when $d = 1$.)

The present paper examines several subclasses of functions of positive real part in \mathbb{B}^d , which are shown to admit what one might call a “noncommutative Herglotz representation;” that is, a representation where the integral of the Herglotz kernel against a measure (as in (1.3)) is replaced by the value of an operator-valued Herglotz kernel (taking values in a C*-algebra A) under a positive functional on the C*-algebra. The classes we consider will turn out to lie strictly between M^+ and O^+ .

Indeed a result of this form (though not stated in quite these terms) was established by McCarthy and Putinar [7] for functions in the *positive Schur class* S^+ , which consists of all holomorphic functions f on \mathbb{B}^d for which the Hermitian kernel

$$\frac{f(z) + \overline{f(w)}}{1 - \langle z, w \rangle}$$

is positive semidefinite. In the noncommutative Herglotz formula for S^+ , the algebra $C(\partial\mathbb{B}^d)$ of continuous functions on $\partial\mathbb{B}^d$ is replaced by the *Cuntz algebra* \mathcal{O}_d , which is the universal C*-algebra generated by a family of isometries V_1, \dots, V_d satisfying the relations

$$V_i^* V_j = \delta_{ij}, \quad \sum_{j=1}^d V_j V_j^* = I.$$

The Herglotz kernel is replaced by the \mathcal{O}_d -valued holomorphic function

$$H(z, V) = 2(I - \sum_{j=1}^d z_j V_j)^{-1} - I.$$

We then have the following theorem:

Theorem 1.1 ([7]). *Let f be holomorphic in \mathbb{B}^d . Then $f \in S^+$ if and only if there exists a positive linear functional ρ on \mathcal{O}_d and a real number t such that*

$$(1.4) \quad f(z) = \rho(H(z, V)) + it.$$

(This result was later obtained independently by Popescu [10].) Letting O^+ denote the class of all holomorphic functions with positive real part in \mathbb{B}^d , McCarthy and Putinar established the inclusions

$$M^+ \subset S^+ \subset O^+$$

and showed that each is proper. They also introduced a pairing on O^+ (basically the pairing induced by the Fantappiè transform, analogous to the Cauchy pairing in one variable) such that the cones M^+ and O^+ are dual to each other, and the dual of S^+ is contained in S^+ . The main results of this paper continue this line of investigation: we introduce a fourth class R^+ which is shown to lie strictly between M^+ and S^+ , and show that R^+ and S^+ are dual to each other; this answers an open question of [7]. We also establish an operator-valued Herglotz representation for R^+ , show that R^+ is characterized by a von Neumann-type inequality, and describe some of the extreme rays of R^+ and S^+ .

2. POSITIVE CLASSES

Let O denote the set of all holomorphic functions on the unit ball $\mathbb{B}^d \subset \mathbb{C}^d$, and $O^+ \subset O$ the functions of positive real part. We equip O with the topology of uniform convergence on compact subsets of \mathbb{B}^d , which makes O into a locally convex topological vector space (in fact a Fréchet space). We will be interested in several subclasses of O^+ which all satisfy a number of functional-analytic conditions, codified by the following definition.

Definition 2.1. *A positive class on \mathbb{B}^d is a set of functions $\mathcal{P} \subset O^+$ which is a closed, convex cone in O and is closed under dilations, that is, if $f \in \mathcal{P}$ then $f_r \in \mathcal{P}$ for all $0 \leq r \leq 1$, where f_r is defined by*

$$f_r(z) = f(rz)$$

Evidently O^+ is a positive class; we will discuss further examples below (the classes M^+ , R^+ and S^+). It is also clear that the intersection of any collection of positive classes is a positive class, so given any subset $E \subset O^+$ we may speak of the positive class $\mathcal{P}(E)$ it generates.

We introduce a family of Cauchy-Fantappiè pairings analogous to the classical Cauchy pairing in the disk. In the case of the disk, if the functions are sufficiently smooth then the pairing is essentially the inner product in the Hilbert space H^2 . The analogous Hilbert space on \mathbb{B}^d is the Drury-Arveson space H_d^2 , which is most easily defined as the

reproducing kernel Hilbert space whose kernel is the Fantappiè kernel

$$k(z, w) = \frac{1}{1 - \langle z, w \rangle} = \sum_{\alpha} z^{\alpha} \overline{w^{\alpha}} \frac{|\alpha|!}{\alpha!}$$

Here we have used the usual multi-index notation:

$$\begin{aligned} \alpha &= (\alpha_1, \dots, \alpha_d) \in \mathbb{N}^d, \\ \alpha! &= \alpha_1! \alpha_2! \cdots \alpha_d!, \\ |\alpha| &= \alpha_1 + \cdots + \alpha_d, \\ z^{\alpha} &= z_1^{\alpha_1} \cdots z_d^{\alpha_d}. \end{aligned}$$

For functions $f, g \in H_d^2$ with Taylor expansions

$$f(z) = \sum_{\alpha \in \mathbb{N}^d} c_{\alpha} z^{\alpha}, \quad g(z) = \sum_{\alpha \in \mathbb{N}^d} d_{\alpha} z^{\alpha}$$

the inner product is given by

$$(2.1) \quad \langle f, g \rangle_{H_d^2} = \sum_{\alpha} c_{\alpha} \overline{d_{\alpha}} \frac{\alpha!}{|\alpha|!}$$

We may also express the inner product in an integral formula [1]. To state it we let \mathcal{R}_d denote the *radial derivative*:

$$(\mathcal{R}_d f)(z) := \sum_{j=1}^d z_j \frac{\partial f}{\partial z_j}$$

Then

$$(2.2) \quad \langle f, g \rangle_{H_d^2} = f(0) \overline{g(0)} + \frac{1}{d!} \int_{\mathbb{B}^d} (\mathcal{R}_d f)(z) \overline{g(z)} \frac{1}{|z|^{2d}} d\nu(z)$$

where ν is normalized volume measure on \mathbb{B}^d . Note that the singularity in the integrand of (2.2) is integrable near 0, since

$$|(\mathcal{R}_d f)(z)| \leq |z| \|Df(z)\|.$$

Now define a family of pairings $Q_r(\cdot, \cdot)$ on O as follows: given functions $f, g \in O$ with Taylor expansions

$$f(z) = \sum_{\alpha \in \mathbb{N}^d} c_{\alpha} z^{\alpha}, \quad g(z) = \sum_{\alpha \in \mathbb{N}^d} d_{\alpha} z^{\alpha}$$

define for each $0 < r < 1$

$$(2.3) \quad Q_r(f, g) = \sum_{\alpha} c_{\alpha} \overline{d_{\alpha}} r^{|\alpha|} \frac{\alpha!}{|\alpha|!} + f(0) \overline{g(0)}$$

Note that we regard $0 \in \mathbb{N}$, so that the term $f(0) \overline{g(0)}$ appears twice in the definition of Q_r . We will write $Q(f, g)$ for the pairing in the case $r = 1$, provided the series is absolutely convergent.

Lemma 2.2. *The series defining $Q_r(f, g)$ converges absolutely for all $f, g \in O$ and all $r \in [0, 1)$.*

Proof. Since the pairing $Q_r(f, g)$ is formally the same (up to the extra $f(0)\overline{g(0)}$ term) as the H_d^2 inner product of the dilated functions $f_{\sqrt{r}}$ and $g_{\sqrt{r}}$, by the Cauchy-Schwarz inequality it suffices to show that if $f \in O$ then all of its dilates f_r belong to H_d^2 . But this is immediate from the integral form of the H_d^2 inner product (2.2), since f_r is analytic across the boundary $\partial\mathbb{B}^d$ and so f_r and its derivatives are uniformly bounded on $\overline{\mathbb{B}^d}$. \square

Our interest is in the duality relationships that obtain between positive classes that admit representations by a generalized Herglotz formula, more specifically, in terms of a positive functional applied to an operator-valued Herglotz kernel, taking values in a (generally noncommutative) C^* -algebra. Using the pairings Q_r we can define the dual of a subset $\mathcal{C} \subset O$:

Definition 2.3. *For $\mathcal{C} \subset O$ define*

$$\mathcal{C}^* = \{g \in O \mid \operatorname{Re} Q_r(f, g) \geq 0 \text{ for all } f \in \mathcal{C} \text{ and all } r \in [0, 1)\}$$

This is slightly different from the dual \mathcal{C}^\dagger defined in [7]; the dual \mathcal{C}^\dagger is defined to be the set of all $g \in O$ such that $\operatorname{Re} Q(f, g) \geq 0$ for all $f \in \mathcal{C}$ for which the series $Q(f, g)$ is absolutely convergent. The present definition is somewhat more convenient technically, because of Lemma 2.2. However it is not hard to show that the two definitions coincide whenever \mathcal{C} is invariant under dilations $f \rightarrow f_r$, which is true by definition when \mathcal{C} is a positive class.

Lemma 2.4. *If \mathcal{C} is dilation invariant, then $\mathcal{C}^* = \mathcal{C}^\dagger$.*

Proof. Suppose $g \in \mathcal{C}^\dagger$. Then $Q_r(f, g) = Q(f_r, g)$ is absolutely convergent for all $f \in \mathcal{C}$ and all $r < 1$, and hence has nonnegative real part since $f_r \in \mathcal{C}$. Thus $g \in \mathcal{C}^*$. Conversely, if $g \in \mathcal{C}^*$, $f \in \mathcal{C}$ and $Q(f, g)$ is absolutely convergent, then $\operatorname{Re} Q(f, g) = \lim_{r \rightarrow 1} \operatorname{Re} Q_r(f, g) \geq 0$, so $g \in \mathcal{C}^\dagger$. \square

From the proof of Lemma 2.2, $Q(f, g)$ will be absolutely convergent whenever either f or g is holomorphic in a neighborhood of the closed ball. In particular, every continuous linear functional on O can be represented in this way:

Theorem 2.5. *A linear map $\varphi : O \rightarrow \mathbb{C}$ is continuous if and only if there exists $g \in O(\overline{\mathbb{B}^d})$ such that*

$$\varphi(f) = Q(f, g)$$

for all $f \in O$.

Proof. To prove that each $Q(\cdot, g)$ induces a continuous linear functional, it suffices to prove continuity at 0. Since g is holomorphic in a neighborhood of $\overline{\mathbb{B}^d}$, we can choose $r < 1$ so that the dilate $g_{\frac{1}{r}}$ is still holomorphic in a neighborhood of $\overline{\mathbb{B}^d}$. We may then write

$$Q(f, g) = Q(f_r, g_{\frac{1}{r}})$$

for all $f \in O$. If now f_n is any sequence of functions tending to 0 uniformly on compact sets, for each $r < 1$ the dilates $(f_n)_r$ converge to 0 uniformly on a neighborhood of $\overline{\mathbb{B}^d}$. It follows that this sequence also converges to 0 in H_d^2 . We then have by Cauchy-Schwarz

$$\begin{aligned} |Q(f_n, g)| &\leq |f_n(0)\overline{g(0)}| + |\langle (f_n)_r, g_{\frac{1}{r}} \rangle_{H_d^2}| \\ &\leq |f_n(0)\overline{g(0)}| + \|(f_n)_r\|_{H_d^2} \|g_{\frac{1}{r}}\|_{H_d^2} \end{aligned}$$

which converges to 0 as $n \rightarrow \infty$.

For the converse, let φ be a continuous linear functional on O . Let $C = C(\mathbb{B}^d)$ be the space of continuous functions on \mathbb{B}^d equipped with the topology of local uniform convergence. The dual of C is the space of finite Borel measures with compact support in \mathbb{B}^d . Since O is closed in C , by Hahn-Banach φ extends to a continuous functional on C . Thus there exists a measure with compact support $K \subset \mathbb{B}^d$ such that

$$\varphi(f) = \int_K f d\mu$$

for all $f \in O$. Now define a function $g \in O$ by

$$(2.4) \quad g(z) = \frac{1}{2} \int_K \frac{1 + \langle z, w \rangle}{1 - \langle z, w \rangle} d\bar{\mu}(w)$$

Since K is at positive distance from boundary $\partial\mathbb{B}^d$, g is holomorphic in a neighborhood of $\overline{\mathbb{B}^d}$. Since the power series for $(1 - \langle z, w \rangle)^{-1}$ converges uniformly on K , we can pass the integral inside the sum to obtain the power series expansion for g

$$g(z) = \frac{1}{2} \int_K d\bar{\mu}(w) + \sum_{\alpha \neq 0} z^\alpha \frac{|\alpha|!}{\alpha!} \int_K \bar{w}^\alpha d\bar{\mu}(w).$$

It is then readily verified that

$$(2.5) \quad \varphi(f) = \int_K f d\mu = Q(f, g).$$

(Indeed, the pairing $Q(\cdot, \cdot)$ is defined precisely so that (2.5) holds when g is of the form (2.4).) \square

We now introduce the positive M^+ , which consists of functions that can be written as Herglotz transforms of positive measures on $\partial\mathbb{B}^d$:

Definition 2.6. $f \in M^+$ if and only if there exists a finite, positive Borel measure μ , supported on $\partial\mathbb{B}^d$, such that

$$f(z) = \int_{\partial\mathbb{B}^d} \frac{1 + \langle z, \zeta \rangle}{1 - \langle z, \zeta \rangle} d\mu(\zeta) + i \operatorname{Im} f(0)$$

Unlike in one dimension, M^+ is a proper subclass of O^+ , however we do have the following duality relation (compare [7, Theorem 8.2]):

Theorem 2.7. $M^{+*} = O^+$ and $O^{+*} = M^+$.

Proof. By a calculation similar to that in the proof of Theorem 2.5, if $g \in M^+$ and $f \in O$ we have

$$Q_r(f, g) = \int_{\partial\mathbb{B}^d} f_r d\mu$$

The real part of this is positive for all μ and all $r < 1$ if and only if $f \in O^+$. The opposite duality will follow from Theorem 2.8 below. \square

Theorem 2.8. If $\mathcal{P} \supset M^+$ is a positive class, then \mathcal{P}^* is a positive class and $\mathcal{P}^{**} = \mathcal{P}$.

Proof. Since $\mathcal{P} \supset M^+$, we have $\mathcal{P}^* \subset O^+$. All of the remaining properties of a positive class now follow from the definition of the pairings Q_r .

It is immediate from definitions that $\mathcal{P} \subset \mathcal{P}^{**}$. The reverse inclusion is proved by a cone separation argument. If $h \in \mathcal{P}^{**} \setminus \mathcal{P}$, since \mathcal{P} is a closed, convex cone we can find a linear functional φ on O such that $\operatorname{Re} \varphi(f) \geq 0$ for all $f \in \mathcal{P}$ but $\operatorname{Re} \varphi(h) < 0$. By Theorem 2.5 there exists $g \in O$ such that $\varphi(f) = Q(f, g)$ for all $f \in O$. Since \mathcal{P} is a positive class, we have $Q_r(f, g) = Q(f_r, g) = \varphi(f_r)$ for all $f \in \mathcal{P}$; taking real parts it follows that $g \in \mathcal{P}^*$. But then $\operatorname{Re} Q_r(h, g) < 0$ for r sufficiently close to 1, contradicting $h \in \mathcal{P}^{**}$. Thus $\mathcal{P}^{**} \subset \mathcal{P}$. \square

Theorem 2.9. Suppose $\mathcal{P} \supset M^+$ is a positive class with $\mathcal{P} \subset \mathcal{P}^*$. Then there exists a positive class \mathcal{W} such that

$$\mathcal{P} \subset \mathcal{W} \subset \mathcal{P}^*$$

and $\mathcal{W} = \mathcal{W}^*$.

Proof. Let \mathcal{U} denote the collection of all positive classes \mathcal{W} such that $\mathcal{P} \subset \mathcal{W} \subset \mathcal{P}^*$ and $\mathcal{W} \subset \mathcal{W}^*$. The collection \mathcal{U} is nonempty (it contains \mathcal{P}) and partially ordered by inclusion. We apply Zorn's lemma to obtain a maximal element: if $\{\mathcal{W}_j\}_{j \in J}$ is an increasing chain in \mathcal{U} , define $\mathcal{W}_\infty = \overline{\cup_j \mathcal{W}_j}$. It is clear that \mathcal{W}_∞ is a closed, convex cone in

O^+ . Moreover, if $f \in \mathcal{W}_\infty$ and $r < 1$, we have a sequence $f_j \rightarrow f$ locally uniformly, with each f_j in some \mathcal{W}_j . Since each \mathcal{W}_j is a positive class, $(f_j)_r \in \mathcal{W}_j$ and $(f_j)_r \rightarrow f_r$, so $f_r \in \mathcal{W}_\infty$ and hence \mathcal{W}_∞ is a positive class. It remains to check that $\mathcal{W}_\infty \subset \mathcal{W}_\infty^*$. But it is straightforward to check that $\mathcal{W}_\infty^* = \bigcap_j \mathcal{W}_j^*$, and from this it follows that \mathcal{W}_∞ belongs to \mathcal{U} . Thus every increasing chain in \mathcal{U} has an upper bound, so Zorn's lemma gives a maximal element \mathcal{W} .

To see that $\mathcal{W} = \mathcal{W}^*$, we proceed by contradiction. If $f \in \mathcal{W}^* \setminus \mathcal{W}$, then we form a new positive class \mathcal{W}_f , the positive class generated by \mathcal{W} and f . We now check that $\mathcal{W}_f \subset \mathcal{W}_f^*$, contradicting the maximality of \mathcal{W} .

The set of functions of the form

$$g = g_0 + \sum_{j=1}^n c_j f_{r_j}$$

with $g_0 \in \mathcal{W}$, $c_j > 0$ and $r_j \in [0, 1)$, is dense in \mathcal{W}_f . We check that $\operatorname{Re} Q_r(g, h) \geq 0$ for all $g, h \in \mathcal{W}_f$ of the above form; positivity for all $g, h \in \mathcal{W}_f$ follows by taking limits. Letting

$$h = h_0 + \sum_{k=1}^m d_k f_{s_k}$$

we have for $r < 1$

$$(2.6) \quad Q_r(g, h) = Q_r(g_0, h_0) +$$

$$(2.7) \quad + Q_r(g_0, \sum_{k=1}^m d_k f_{s_k})$$

$$(2.8) \quad + Q_r(\sum_{j=1}^n c_j f_{r_j}, h_0)$$

$$(2.9) \quad + Q_r(\sum_{j=1}^n c_j f_{r_j}, \sum_{k=1}^m d_k f_{s_k})$$

The first term has positive real part because $\mathcal{W} \subset \mathcal{W}^*$, the next two because $f \in \mathcal{W}^*$, and the last from the definition of Q_r . \square

3. OPERATOR-VALUED HERGLOTZ KERNELS

A *row contraction* is a d -tuple of operators $T = (T_1, \dots, T_d)$ on a Hilbert space \mathcal{H} such that

$$I - T_1 T_1^* - \dots - T_d T_d^* \geq 0$$

If T is a row contraction and $z \in \mathbb{B}^d$, then the operator

$$\langle z, T \rangle := z_1 T_1^* + \cdots + z_d T_d^*$$

is a strict contraction. We may then define an operator-valued Herglotz kernel

$$\begin{aligned} H(z, T) &:= 2(I - \langle z, T \rangle)^{-1} - I \\ &= (I + \langle z, T \rangle)(I - \langle z, T \rangle)^{-1} \end{aligned}$$

which we view as an analytic function on \mathbb{B}^d taking values in $\mathcal{B}(\mathcal{H})$. A straightforward calculation shows

$$\operatorname{Re} H(z, T) = 2(I - \langle z, T \rangle)^{-1}(I - \langle z, T \rangle \langle z, T \rangle^*)(I - \langle z, T \rangle^*)^{-1} \geq 0$$

Hence if ρ is a positive linear functional on the C*-algebra generated by T , the holomorphic function

$$(3.1) \quad f(z) = \rho(H(z, T))$$

has positive real part on \mathbb{B}^d . We will call such f the *Herglotz transform* of the pair (ρ, T) . Theorem 1.1 characterizes the positive class S^+ in terms of Herglotz transforms.

Three particular row contractions (and the C*-algebras they generate) will be of interest. First, we may view the commutative C*-algebra $C(\partial\mathbb{B}^d)$ as the universal C*-algebra generated by a d -tuple of commuting normal elements $Z = (Z_1, \dots, Z_d)$ satisfying

$$I - Z_1 Z_1^* - \cdots - Z_d Z_d^* = 0$$

(these are called *spherical operators* in [2]; one should think of these “universal generators” simply as the coordinate functions ζ_1, \dots, ζ_d in $C(\partial\mathbb{B}^d)$). Second, we consider the row contraction $S = (S_1, \dots, S_d)$ where S_j is multiplication by z_j on the space H_d^2 . These generate the *d-variable Toeplitz algebra* \mathcal{T}_d . Finally, we have the d -tuple of *Cuntz isometries*, which are isometries with orthogonal ranges whose final projections sum to I . That is, d -tuples $V = (V_1, \dots, V_d)$ such that

$$V_i^* V_j = \delta_{ij}, \quad I - V_1 V_1^* - \cdots - V_d V_d^* = 0.$$

The *Cuntz C* algebra* \mathcal{O}_d is the universal C*-algebra generated by the Cuntz isometries. We now have three positive classes M^+ , R^+ and S^+ which can be expressed in terms of operator-valued Herglotz formulae. The classes M^+ and S^+ were described in [7].

Theorem 3.1. *Each of the following is a positive class on \mathbb{B}^d :*

- M^+ , consisting of all functions of the form

$$f(z) = \rho(H(z, Z)) + i \operatorname{Im} f(0)$$

where ρ is a positive linear functional on $C(\partial\mathbb{B}^d)$,

- R^+ , consisting of all functions of the form

$$f(z) = \rho(H(z, S)) + i\text{Im } f(0)$$

where ρ is a positive linear functional on \mathcal{T}_d ,

- S^+ , consisting of all functions of the form

$$f(z) = \rho(H(z, V)) + i\text{Im } f(0)$$

where ρ is a positive linear functional on \mathcal{O}_d .

Proof. We prove that R^+ is a positive class; it will be clear that an analogous proof works for the others.

It is clear from the definition that R^+ is a convex cone, since the set of positive functionals on a C^* -algebra is a convex cone. To see that R^+ is closed, let f_n be a sequence in R^+ converging uniformly on compact sets to a function $f \in O^+$ (we assume for simplicity that $f_n(0)$ is real for all n). Then there is a sequence of positive functionals ρ_n on \mathcal{T}_d such that

$$f_n(z) = \rho_n(H(z, S)).$$

Since the sequence $\|\rho_n\| = |\rho_n(I)| = |f_n(0)|$ is bounded, by the Banach-Alaoglu theorem ρ_n has a weak- $*$ cluster point ρ . But then we have

$$f(z) = \rho(H(z, S))$$

and hence $f \in R^+$.

To prove that R^+ is dilation invariant, we obtain an equivalent definition of R^+ (which will also be of use later). By the Gelfand-Naimark-Segal theorem applied to the functional ρ , there exist a Hilbert space \mathcal{H} , a vector $\xi \in \mathcal{H}$ and a representation $\pi : \mathcal{T}_d \rightarrow \mathcal{B}(\mathcal{H})$ such that

$$\rho(H(z, S)) = \langle H(z, \pi(S))\xi, \xi \rangle_{\mathcal{H}}$$

By Arveson's dilation theorem [2], we conclude that $f \in R^+$ if and only if there exists a Hilbert space \mathcal{H} , a vector $\xi \in \mathcal{H}$ and a commuting row contraction T such that (ignoring the harmless imaginary constant)

$$f(z) = \langle H(z, T)\xi, \xi \rangle_{\mathcal{H}}$$

Representing $f \in R^+$ in this way, we have

$$(3.2) \quad f_r(z) = \langle H(rz, T)\xi, \xi \rangle_{\mathcal{H}}$$

$$(3.3) \quad = \langle H(z, rT)\xi, \xi \rangle_{\mathcal{H}}$$

and hence $f_r \in R^+$ since rT is a commuting row contraction. \square

Of course, since a positive linear functional on $C(\partial\mathbb{B}^d)$ is just a finite Borel measure μ supported on $\partial\mathbb{B}^d$, the Herglotz formula for M^+ takes the form

$$f(z) = \int_{\partial\mathbb{B}^d} \frac{1 + \langle z, \zeta \rangle}{1 - \langle z, \zeta \rangle} d\mu(\zeta) + i\text{Im } f(0)$$

in accord with the original definition.

As in the proof of the theorem, by combining the GNS representation theorem with appropriate dilation theorems we can obtain equivalent descriptions of R^+ and S^+ that are sometimes useful. We first recall that the original definition of S^+ is the set of all $f \in O$ such that the kernel

$$(3.4) \quad \frac{f(z) + \overline{f(w)}}{1 - \langle z, w \rangle}$$

is positive. In [7] it is shown that $f \in S^+$ if and only if there exists a Cuntz isometry V acting on a space \mathcal{H} and a vector $\xi \in \mathcal{H}$ such that

$$f(z) = \langle H(z, V)\xi, \xi \rangle + \text{Im } f(0).$$

By Popescu's dilation theorem [9], every row contraction T dilates to a Cuntz isometry, and hence S^+ consists of all functions of the form

$$(3.5) \quad f(z) = \langle H(z, T)\xi, \xi \rangle + \text{Im } f(0).$$

where T is any row contraction. In contrast, the Arveson dilation theorem [2] says that every commuting row contraction dilates to a representation of the d -shift S , and hence R^+ consists of all functions f admitting a representation of the form (3.5) for some *commuting* row contraction T .

The formalism established thus far leads to an ‘‘abstract Herglotz formula’’ characterizing positive classes \mathcal{P} contained in S^+ :

Theorem 3.2. *To each positive class $\mathcal{P} \subset S^+$ on \mathbb{B}^d there corresponds a weak- $*$ closed, convex subset $\widehat{\mathcal{P}}$ of the state space of the Cuntz algebra \mathcal{O}_d , such that $f \in \mathcal{P}$ if and only if there exists a state $\rho \in \widehat{\mathcal{P}}$ and a real number $\lambda \geq 0$ such that*

$$(3.6) \quad f(z) = \lambda\rho(H(z, V)) + \text{Im } f(0)$$

Proof. Given \mathcal{P} , let $\widehat{\mathcal{P}}$ consist of all states representing functions $f \in \mathcal{P}$ as in (3.6). Since \mathcal{P} is a positive class, $\widehat{\mathcal{P}}$ is convex, and it is straightforward to check that if $\rho_n \rightarrow \rho$ weak- $*$, then $f_n \rightarrow f$ uniformly on compact sets, so $\widehat{\mathcal{P}}$ is closed. \square

It must be kept in mind, however, that in general the state ρ corresponding to f may be far from unique. This is even the case in M^+ ,

since the closed linear span of the Herglotz kernels and their adjoints is a proper subspace of $C(\partial\mathbb{B}^d)$ when $d > 1$.

We now return to the particular positive classes described by Theorem 3.1. Our main goal is to establish that R^+ and S^+ are dual to each other in the sense of Definition 2.3, and to show that the inclusion $R^+ \subset S^+$ is proper, resolving an open question of [7].

To begin with we recall the ‘‘symmetrized functional calculus’’ from [7]. If $T = (T_1, \dots, T_d)$ is a d -tuple of operators, we can make sense of a monomial z^α in T by averaging over all possible orderings of the T_i 's. Explicitly, for a multi-index $\alpha = (\alpha_1, \dots, \alpha_d)$ we define

$$(z^\alpha)^{sym}(T) = \frac{\alpha!}{|\alpha|!} \sum T_{i_1} T_{i_2} \cdots T_{i_{|\alpha|}}$$

where the sum is taken over all permutations of α_1 1's, α_2 2's, etc.

Theorem 3.3. $R^{+*} = S^+$ and $S^{+*} = R^+$.

Proof. By the above discussion, g belongs to R^+ if and only if there exists a commuting row contraction T on a Hilbert space \mathcal{H} and a vector $\xi \in \mathcal{H}$ such that

$$(3.7) \quad \frac{1}{2}[g(z) + \overline{g(0)}] = \langle (I - \langle z, T \rangle)^{-1} \xi, \xi \rangle$$

$$(3.8) \quad = \sum_{j=0}^{\infty} \langle \langle z, T \rangle^j \xi, \xi \rangle$$

$$(3.9) \quad = \sum_{\alpha} z^\alpha \langle (z^\alpha)^{sym}(T)^* \xi, \xi \rangle \frac{|\alpha|!}{\alpha!}$$

It follows that for any $f \in O$ and $r < 1$,

$$\begin{aligned} \operatorname{Re} Q_r(f, g) &= 2\operatorname{Re} \langle (f_r)^{sym}(T)^* \xi, \xi \rangle \\ &= 2\operatorname{Re} \langle f_r(T)^* \xi, \xi \rangle \\ &= 2\operatorname{Re} \langle f_r(T) \xi, \xi \rangle \end{aligned}$$

since T commutes. So $\operatorname{Re} Q_r(f, g) \geq 0$ for all $g \in R^+$ if and only if $\operatorname{Re} f_r(T) \geq 0$ for all commuting row contractions T . Taking $T = S$ shows that we must have $f_r \in S^+$ for all $r < 1$, so $f \in S^+$. But then $\operatorname{Re} f_r(T) \geq 0$ for all commuting row contractions T by von Neumann's inequality for S^+ . Thus, $R^{+*} = S^+$. The opposite duality follows from Theorem 2.8. \square

Our proof that $R^{+*} = S^+$ appealed to the characterization of S^+ in terms of a von Neumann inequality; the opposite duality $S^{+*} = R^+$ shows that a similar von Neumann inequality (for the symmetrized functional calculus) characterizes R^+ :

Theorem 3.4. *Let $f \in O$. Then $f \in R^+$ if and only if*

$$(3.10) \quad \operatorname{Re} (f_r)^{\operatorname{sym}}(T) \geq 0$$

for all $r < 1$ and all row contractions T .

Proof. We imitate the proof of Theorem 3.3, this time starting with $g \in S^+$ rather than R^+ . Arguing as before, g belongs to S^+ if and only if there exists a row contraction T and a vector ξ such that

$$\frac{1}{2}[g(z) + \overline{g(0)}] = \sum_{\alpha} z^{\alpha} \langle (z^{\alpha})^{\operatorname{sym}}(T)^* \xi, \xi \rangle \frac{|\alpha|!}{\alpha!}$$

Now, for any $f \in O$, we find that $\operatorname{Re} Q_r(f, g) = 2\operatorname{Re} \langle (f_r)^{\operatorname{sym}}(T)\xi, \xi \rangle$. Hence $f \in S^{+*} = R^+$ if and only if $\operatorname{Re} (f_r)^{\operatorname{sym}}(T) \geq 0$ for all row contractions T . \square

Corollary 3.5. *R^+ is a proper subset of S^+ .*

Proof. If $f \in R^+$, then by the preceding theorem

$$\operatorname{Re} (f_r)^{\operatorname{sym}}(V) \geq 0$$

for all Cuntz isometries V . Suppose by way of contradiction that this is the case for all $f \in S^+$. Let \mathcal{A} denote the set of all polynomials $p(S) \in \mathcal{T}_d$ and let \mathcal{S} be the operator system $\mathcal{A} + \mathcal{A}^*$. Our assumption would then imply that for every set of Cuntz isometries V acting on a Hilbert space \mathcal{H} , the map from \mathcal{S} to $\mathcal{B}(\mathcal{H})$ given by

$$p(S) + q(S)^* \rightarrow p^{\operatorname{sym}}(V) + q^{\operatorname{sym}}(V)^*$$

is positive. Then by [8, Corollary 2.8], the map sending $p(S)$ to $p^{\operatorname{sym}}(V)$ is contractive. But Davidson and Pitts [3, Example 3.4] give an explicit counterexample showing that this map is not contractive: in particular, for the polynomial $p(z) = z_1 + z_1 z_2$ we have $\|p(S)\| < \sqrt{2}$ but $\|p^{\operatorname{sym}}(V)\| = \sqrt{\frac{5}{2}}$. Thus $R^+ = S^{+*} \subsetneq S^+$. \square

Theorem 3.6. *The following chain of inclusions holds, and each is proper:*

$$M^+ \subset R^+ \subset S^+ \subset O^+.$$

Proof. The inclusion of R^+ in S^+ is given by the previous theorem. The proper inclusion of S^+ in O^+ is well known, e.g. proved in [7]. By Theorem 3.3 the inclusion $M^+ \subsetneq R^+$ is dual to $S^+ \subsetneq O^+$. \square

Theorem 2.9 guarantees the existence of a self-dual positive class lying between R^+ and S^+ , however we do not know any concrete example.

Problem 3.7. *Give an explicit example of a self-dual positive class lying between R^+ and S^+ .*

4. EXTREME POINTS

If \mathcal{P} is a positive class, we can normalize each function $f \in \mathcal{P}$ so that $f(0) = 1$; we denote the resulting subclass \mathcal{P}_0 . Since O_0^+ is compact, it follows that \mathcal{P}_0 is also compact and hence by the Krein-Milman theorem it is the closed convex hull of its extreme points. It would obviously be desirable to identify the extreme points of the classes R_0^+ and S_0^+ . It is not hard to see that the extreme points of M_0^+ are exactly the functions obtained as Herglotz integrals of point masses; that is the extreme points of M_0^+ are the functions

$$h_\zeta(z) = \frac{1 + \langle z, \zeta \rangle}{1 - \langle z, \zeta \rangle}, \quad \zeta \in \partial\mathbb{B}^d.$$

It is known that these functions are not extreme for O^+ ; for example when $d = 2$ the functions

$$\frac{1 + z_1}{1 - z_1} + z_2^2 h(z_1, z_2)$$

belong to O^+ whenever $|h| \leq \frac{1}{4}$ [11, Section 19.2.6]. We will show presently that the functions h_ζ are extreme points for S^+ ; but of course in S^+ there must be others as well. The following proposition is elementary:

Proposition 4.1. *If \mathcal{P}_0 is a normalized positive class lying between M_0^+ and S_0^+ and f is an extreme point of \mathcal{P}_0 , then there exists a pure state on the Cuntz algebra \mathcal{O}_d such that*

$$f(z) = \rho(H(z, V))$$

Proof. The proof is a routine application of the Krein-Milman theorem: let \mathcal{S} denote the set of all states ρ on \mathcal{O}_d such that $\rho(H(z, V))$ belongs to \mathcal{P}_0 . \mathcal{S} is a weak-* compact, convex set in the dual of \mathcal{O}_d . Let $\mathcal{S}_f \subset \mathcal{S}$ denote those states such that $\rho(H(z, V)) = f$. \mathcal{S}_f is again compact and convex, so by Krein-Milman is the closed hull of its extreme points. It is now not hard to see that if f is extreme then every extreme point of \mathcal{S}_f is in fact extreme in \mathcal{S} . Indeed, suppose f is extreme and ρ is extreme in \mathcal{S}_f . If $\rho_0, \rho_1 \in \mathcal{S}$ satisfy $t\rho_0 + (1-t)\rho_1 = \rho$ for some $t \in (0, 1)$, then taking Herglotz transforms shows $\rho_0, \rho_1 \in \mathcal{S}_f$ and hence $\rho_0 = \rho_1 = \rho$. \square

Of course, the converse does not hold; there are many examples of pure states ρ for which the associated f is not extreme in S_0^+ .

To prove that the functions h_ζ are extreme for S^+ , we make use of the following strengthened Schwarz lemma for the Schur class \mathcal{S}_d . Recall that a function φ belongs to \mathcal{S}_d if and only if the Hermitian kernel

$$k^\varphi(z, w) = \frac{1 - \varphi(z)\overline{\varphi(w)}}{1 - \langle z, w \rangle}$$

is positive semidefinite. A simple calculation shows that $\varphi \in \mathcal{S}_d$ if and only if its Cayley transform $\frac{1+\varphi}{1-\varphi}$ belongs to S^+ .

Lemma 4.2. *Suppose $g \in \mathcal{S}_d$, $g(0) = 0$ and*

$$\frac{\partial g}{\partial z_1}(0) = 1.$$

Then $g(z) = z_1$.

Proof. To say that $g \in \mathcal{S}_d$ means that the kernel

$$(4.1) \quad \frac{1 - g(z)\overline{g(w)}}{1 - \langle z, w \rangle}$$

is positive. Let G denote the restriction of g to the disk $D = \{(z_1, 0) : |z_1| < 1\}$. Then $G(0) = 0$, $G'(0) = 1$ and $|G(z)| \leq 1$ for all $|z| < 1$. By the one-variable Schwarz lemma, we have $G(z_1) = z_1$. Thus the restriction of g to D equals z_1 . But since restriction of the kernel (4.1) to D is

$$\frac{1 - G(z_1)\overline{G(w_1)}}{1 - z_1\overline{w_1}} \equiv 1$$

which trivially has finite rank, the function $g|_D$ has a unique extension to an \mathcal{S}_d function in \mathbb{B}^d . This forces $g \equiv z_1$. \square

Theorem 4.3. *For each $\zeta \in \partial\mathbb{B}^d$, the function*

$$h_\zeta(z) = \frac{1 + \langle z, \zeta \rangle}{1 - \langle z, \zeta \rangle}$$

is an extreme point of S^+ .

Proof. Without loss of generality we assume $\zeta = e_1 = (1, \dots, 0)$. Suppose there exist functions f, g in the Schur class \mathcal{S}_d and $t \in (0, 1)$ such that

$$h(z) = h_{e_1}(z) = \frac{1 + z_1}{1 - z_1} = t \frac{1 + f(z)}{1 - f(z)} + (1 - t) \frac{1 + g(z)}{1 - g(z)}$$

Restrict this equation to the disk $D = \{(z_1, 0) : |z_1| < 1\}$. Since $h(z) = h(z_1)$ is extreme for $O^+(\mathbb{D})$, it follows that f and g must equal z_1 when restricted to D . By the uniqueness in the Schwarz lemma for

\mathcal{S}_d , we must have $f(z) = g(z) = z_1$ for all $z \in \mathbb{B}^d$. Thus h is extreme in S^+ . \square

By Proposition 4.1, there exists a pure state ρ on \mathcal{O}_d such that

$$h_\zeta(z) = \rho(H(z, V)).$$

In fact it can be shown that in this case the state ρ is unique, and equal to the *Cuntz state* ω_ζ defined by

$$\omega_\zeta(V_{i_1} \cdots V_{i_m} V_{j_1}^* \cdots V_{j_n}^*) = \zeta_{i_1} \cdots \zeta_{i_m} \overline{\zeta_{j_1}} \cdots \overline{\zeta_{j_n}}.$$

It seems unlikely that there would be any very concrete parameterization of the extreme points of S_0^+ . Nonetheless at present we do not know any extreme points of S_0^+ besides the functions h_ζ , so it would be desirable to find other examples.

Problem 4.4. *Find other extreme points of R_0^+ and S_0^+ . Can the extreme points of R_0^+ or S_0^+ be classified? Which pure states on \mathcal{T}_d (resp. \mathcal{O}_d) give rise to extreme points?*

We conclude with a remark on the mean growth of functions in S^+ . One reason for historical interest in extreme points of O^+ was their connection with the inner function conjecture (namely, that there were no inner functions on the ball \mathbb{B}^d when $d > 1$); in fact it had been conjectured that functions $f \in O^+$ all belonged to the Hardy space H^1 (see [11, Chapter 19]). The inner function conjecture turned out to be false, but it is obviously of interest to ask the same questions of other classes, particularly S^+ . Recall that for $0 < p < \infty$, a function $f \in O$ belongs to the Hardy space H^p if and only if

$$(4.2) \quad \sup_{0 < r < 1} \int_{\partial \mathbb{B}^d} |f(r\zeta)|^p d\sigma(\zeta) < \infty,$$

where σ denotes surface measure on the sphere. When $p \geq 1$ the p^{th} root of this quantity defines a norm on H^p . By standard estimates (see [11, Proposition 1.4.10]), the extreme points h_ζ belong to H^p for all $p < d$. In fact this is characteristic of the growth of functions in S^+ :

Theorem 4.5. *$S^+(\mathbb{B}^d) \subset H^p(\mathbb{B}^d)$ for all $p < d$.*

Proof. Let $g \in S^+$ and write $g = \frac{1+\varphi}{1-\varphi}$ for $\varphi \in \mathcal{S}_d$. Define a holomorphic mapping $\psi : \mathbb{B}^d \rightarrow \mathbb{B}^d$ by

$$\psi(z) = (\varphi(z), 0 \dots 0)$$

Since $\varphi \in \mathcal{S}_d$, the kernel

$$\frac{1 - \langle \psi(z), \psi(w) \rangle}{1 - \langle z, w \rangle} = \frac{1 - \varphi(z) \overline{\varphi(w)}}{1 - \langle z, w \rangle}$$

is positive semidefinite. It follows that the composition operator

$$C_\psi : f \rightarrow f \circ \psi$$

is bounded on H^2 [4, Theorem 5], and hence bounded on H^p for all $p > 0$ [6, Corollary 1.2]. Thus

$$g = \frac{1 + \varphi}{1 - \varphi} = C_\psi \left(\frac{1 + z_1}{1 - z_1} \right)$$

belongs to H^p for all $p < d$. \square

In contrast, O^+ contains functions which do not belong to H^1 ; indeed if φ is an inner function then $g = \frac{1+\varphi}{1-\varphi} \notin H^1$. As a corollary we see that the Schur class \mathcal{S}_d does not contain any inner functions when $d > 1$.

REFERENCES

- [1] Daniel Alpay and H. Turgay Kaptanoğlu. Integral formulas for a sub-Hardy Hilbert space on the ball with complete Nevanlinna-Pick reproducing kernel. *C. R. Acad. Sci. Paris Sér. I Math.*, 333(4):285–290, 2001.
- [2] William Arveson. Subalgebras of C^* -algebras. III. Multivariable operator theory. *Acta Math.*, 181(2):159–228, 1998.
- [3] Kenneth R. Davidson and David R. Pitts. Nevanlinna-Pick interpolation for non-commutative analytic Toeplitz algebras. *Integral Equations Operator Theory*, 31(3):321–337, 1998.
- [4] Michael T. Jury. Reproducing kernels, de Branges-Rovnyak spaces, and norms of weighted composition operators. *Proc. Amer. Math. Soc.*, 135(11):3669–3675, 2007.
- [5] A. Korányi and L. Pukánszky. Holomorphic functions with positive real part on polycylinders. *Trans. Amer. Math. Soc.*, 108:449–456, 1963.
- [6] Barbara D. MacCluer. Compact composition operators on $H^p(B_N)$. *Michigan Math. J.*, 32(2):237–248, 1985.
- [7] John E. McCarthy and Mihai Putinar. Positivity aspects of the Fantappiè transform. *J. Anal. Math.*, 97:57–82, 2005.
- [8] Vern Paulsen. *Completely bounded maps and operator algebras*, volume 78 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2002.
- [9] Gelu Popescu. Isometric dilations for infinite sequences of noncommuting operators. *Trans. Amer. Math. Soc.*, 316(2):523–536, 1989.
- [10] Gelu Popescu. Free holomorphic functions and interpolation. *Math. Ann.*, 342(1):1–30, 2008.
- [11] Walter Rudin. *Function theory in the unit ball of \mathbf{C}^n* , volume 241 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Science]*. Springer-Verlag, New York, 1980.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, GAINESVILLE, FLORIDA 32611-8105

E-mail address: mjury@math.ufl.edu