

Chapter 2: Topological spaces and continuous functions

Section 12: Topological spaces


Many concepts in analysis (continuity, convergence, compactness) only require knowledge of the open sets.

Def A topology on a set X is a collection τ of subsets, called open sets, satisfying


- $\emptyset, X \in \tau$.
- Arbitrary unions of open sets are open:
 $U_\alpha \in \tau \quad \forall \alpha \in I \quad \Rightarrow \quad U_{\alpha \in I} U_\alpha \in \tau$.
- Finite intersections of open sets are open:
 $U_1, \dots, U_n \in \tau \quad \Rightarrow \quad U_1 \cap \dots \cap U_n \in \tau$.

We denote this topological space by (X, τ) or X .


Ex Which of the following are topologies on $X = \{a, b, c\}$?




Yes, indiscrete or trivial topology




Yes, $\tau = \mathcal{P}(X)$ is the discrete topology




No



Yes



No



Yes

Ex Every metric space is a topological space.

The open sets are unions of (open) balls.

Ex X is a set.

$\tau = \{ U \subset X \mid U = \emptyset \text{ or } X - U \text{ is finite} \}$. Let $U^c = X - U$.

Called the finite complement topology.

For example, if $X = \mathbb{R}$, then a nonempty open set is \mathbb{R} with at most a finite number of points removed:



PS

- $\emptyset, X \in \tau$
- Consider $\{ U_\alpha \}_{\alpha \in I}$, $U_\alpha \in \tau$. Want: $\bigcup_{\alpha \in I} U_\alpha \in \tau$.
WLOG, assume $U_\alpha \neq \emptyset$ for all $\alpha \in I$.
So $(U_\alpha)^c$ is finite.
Note $(\bigcup_{\alpha \in I} U_\alpha)^c = \bigcap_{\alpha \in I} (U_\alpha)^c$, which is finite.
So $\bigcup_{\alpha \in I} U_\alpha \in \tau$.
- Let $\emptyset \neq U_1, \dots, U_n \in \tau$. So $X - U_i$ is finite.
Note $(U_1 \cap \dots \cap U_n)^c = U_1^c \cup \dots \cup U_n^c$, which is finite.
So $U_1 \cap \dots \cap U_n \in \tau$.

Rmk There is also a countable complement topology.


Def If τ and τ' are two topologies on X with $\tau \subset \tau'$, then we say τ is coarser and τ' is finer.

Section 13: Basis for a topology

Instead of specifying all open sets in a topology τ , it is often convenient to specify a nice subset that generates τ .

Def A basis for a topology on X is a collection \mathcal{B} of subsets of X such that

- (1) $\forall x \in X \exists B \in \mathcal{B}$ with $x \in B$.
- (2) If $x \in B_1 \cap B_2$ with $B_1, B_2 \in \mathcal{B}$, then $\exists B_3 \in \mathcal{B}$ with $x \in B_3 \subset B_1 \cap B_2$.



Prop Given a basis \mathcal{B} for a topology on X .

Let $\tau = \{ U \subset X \mid \forall x \in U \exists B \in \mathcal{B} \text{ s.t. } x \in B \subset U \}$




Let $\tau' = \{ U \subset X \mid U \text{ is a union of sets in } \mathcal{B} \}$

- Then
- (a) τ is a topology on X
 - (b) $\tau' = \tau$.

Def Call this the topology generated by \mathcal{B} .

Ex X a metric space,
 $\mathcal{B} = \{ \text{open balls} \}$

Ex $X = \mathbb{R}^2$ $\mathcal{B} = \{ \text{open balls} \}$
 or $\mathcal{B}' = \{ \text{axis-aligned open rectangles} \}$
 These generate the same topology.



Ex X $\mathcal{B} = \{ \text{one point sets} \}$ is a basis
 for the discrete topology.

Ex Choosing $\mathcal{B} = \mathcal{T}$ always gives a basis, but it is
 more valuable to find bases $\mathcal{B} \subsetneq \mathcal{T}$.

Proof of Proposition

(a) • $\emptyset \in \mathcal{T}$ since condition is vacuously true.

$X = \bigcup_{B \in \mathcal{B}} B$ by (1), so $X \in \mathcal{T}$.

• $\{ U_\alpha \}_{\alpha \in I}$ with $U_\alpha \in \mathcal{T}$.

If $x \in \bigcup_{\alpha \in I} U_\alpha$,

then $\exists \alpha \in I$ with $x \in U_\alpha$,

so $\exists B \in \mathcal{B}$ with $x \in B \subset U_\alpha \subset \bigcup_{\alpha \in I} U_\alpha$.

• $U_1, \dots, U_n \in \mathcal{T}$. Let $x \in U_1 \cap \dots \cap U_n$.

Claim: $\exists B \in \mathcal{B}$ with $x \in B \subset U_1 \cap \dots \cap U_n$.

Use induction on n .



Base case $n=1$ by definition of \mathcal{T}

Assume true for n . Let $x \in U_1 \cap \dots \cap U_{n+1}$.

Then $x \in U_1 \cap \dots \cap U_n$. By induction $\exists B_1 \in \mathcal{B}$ with $x \in B_1 \subset U_1 \cap \dots \cap U_n$.

Also $x \in U_{n+1}$. By def of \mathcal{T} $\exists B_2 \in \mathcal{B}$ with $x \in B_2 \subset U_{n+1}$.

By (2) $\exists B_3 \in \mathcal{B}$ with $x \in B_3 \subset B_1 \cap B_2 \subset U_1 \cap \dots \cap U_{n+1}$ ✓



(b) $(\tau' \subset \tau)$ Consider an arbitrary element of τ' : $\bigcup_{\alpha \in I} U_\alpha$, $U_\alpha \in \mathcal{B}$.

Let $x \in \bigcup_{\alpha \in I} U_\alpha$. Then $\exists \alpha$ s.t. $x \in U_\alpha$.

So $x \in U_\alpha \subset \bigcup_{\alpha \in I} U_\alpha$. Hence $\bigcup_{\alpha \in I} U_\alpha \in \tau$.

$(\tau \subset \tau')$ Consider an arb elt of τ :

$U \subset X$ s.t. $\forall x \in U \exists B_x \in \mathcal{B}$ s.t. $x \in B_x \subset U$.

Then $U = \bigcup_{x \in U} B_x$. Hence $U \in \tau'$. \square

Lemma 13.2 Let (X, τ) be a topological space.

Let \mathcal{C} be a collection of open sets such that if $x \in U$ for $U \in \tau$, then $\exists C \in \mathcal{C}$ with $x \in C \subset U$.

Then \mathcal{C} is a basis for τ .

PS (1) Since $X \in \tau$, $\forall x \in X \exists C \in \mathcal{C}$ with $x \in C \subset X$. \checkmark

(2) If $x \in C_1 \cap C_2$ for $C_1, C_2 \in \mathcal{C} \subset \tau$, then $C_1 \cap C_2 \in \tau$,

so $\exists C_3 \in \mathcal{C}$ with $x \in C_3 \subset C_1 \cap C_2$. \checkmark



So \mathcal{C} is a basis.

By definition, \mathcal{C} generates the topology τ . \square

Lemma Let $\mathcal{B}, \mathcal{B}'$ be bases for the topologies τ, τ' on X .
 Then τ' is finer than τ ($\tau \subset \tau'$ allowing equality)
 $\Leftrightarrow \forall B \in \mathcal{B}$ and $x \in B, \exists B' \in \mathcal{B}'$ with $x \in B' \subset B$.

PF See book



Ex $X = \mathbb{R}^2$ $\mathcal{B} = \{\text{open balls}\}$
 and $\mathcal{B}' = \{\text{axis-aligned open rectangles}\}$
 generate the same topology.



Def $X = \mathbb{R}$

$\mathcal{B} = \{(a, b) \mid a < b\}$ generates the standard topology

$\mathcal{B}' = \{[a, b) \mid a < b\}$ generates the lower limit topology



Fact $\tau \neq \tau'$ (I.e., τ' is finer than τ ,
 and not vice-versa)

PF sketch Apply the prior lemma. Consider $(a, b) \in \mathcal{B}$
 and $x \in (a, b)$. Note $[x, b) \in \mathcal{B}'$ satisfies
 $x \in [x, b) \subset (a, b)$, as required.



However for $[a, b)$ and $x = a$ and any (a', b') containing a also
 contains $a - \varepsilon$ for some $\varepsilon > 0$ and $a - \varepsilon \notin [a, b)$.

An imperfect analogy

Vector spaces

\mathbb{R}^n
vector spaces
basis

Any vector is a sum of basis elements.

This description is unique.

A vector space has many bases.

All bases have the same size.

Topological spaces

open sets in \mathbb{R}^n
topological spaces
basis

Any open set is a union of basis elements.

Nope.

A topological space has many bases.

Nope.

Basic open sets $\xrightarrow{\text{unions}}$ topology

Certain open sets $\xrightarrow{\text{unions and finite intersections}}$ topology ?


Def A subbasis \mathcal{S} for X is a collection of sets whose union is X . The topology τ generated by subbasis \mathcal{S} is the collection of all unions of finite intersections of elements in \mathcal{S} .

Ex $X = \{0, 1, 2\}$

$\mathcal{S} = \{\{0, 1\}, \{0, 2\}\}$ is a subbasis but not a basis.

$\tau = \{\emptyset, \{0\}, \{0, 1\}, \{0, 2\}, \{0, 1, 2\}\}$.

One basis is $\mathcal{B} = \{\{0\}, \{0, 1\}, \{0, 2\}\}$



Lemma τ is indeed a topology

Pf sketch Show that the collection \mathcal{B} of all finite intersections of elements in \mathcal{S} is a basis.

(1) is easy.

(2) follows since if $B = S_1 \cap \dots \cap S_m$ and $B' = S'_1 \cap \dots \cap S'_m$ are two elements of \mathcal{B} , then $B \cap B'$ is also an element of \mathcal{B} .



$$|\mathcal{S}| = 3$$

$$|\mathcal{B}| = 6 \text{ if you include } \emptyset.$$


$$|\tau| = 2^5$$

Section 14: The order topology

Let X be a set with total order \leq .

For $a, b \in X$, define

- $(a, b) = \{x \in X : a < x < b\}$
- $[a, b) = \{x \in X : a \leq x < b\}$
- $(a, b] = \{x \in X : a < x \leq b\}$
- $[a, b] = \{x \in X : a \leq x \leq b\}$.




Def Let \mathcal{B} contain

- (1) all intervals (a, b)
 - (2) all intervals $[a_0, b)$ where a_0 is the smallest element (if any) in X
 - (3) all intervals $(a, b_0]$ where b_0 is the largest element (if any) in X .
- The collection \mathcal{B} forms a basis for the order topology on X .

Ex The order topology is the standard topology on \mathbb{R} .

Ex $\mathbb{R} \times \mathbb{R}$ with the lexicographic order:

$$a \times b < c \times d \iff a < c \text{ or } a = c, b < d.$$



These intervals actually form a basis on their own.

This is not the standard topology on \mathbb{R}^2 .

Ex The order topology on \mathbb{Z}_+ is the discrete topology

Note $\{n\} = (n-1, n+1)$ for $n > 1$, and $\{1\} = [1, 2)$.

Ex The order topology on $\{1, 2\} \times \mathbb{Z}_+$ (lexicographic order) is not the discrete topology, since any basis element containing 2×1 must contain some $1 \times n$.


$\begin{array}{ccc} & \vdots & \vdots \\ 1 \times 8 & \bullet & \bullet & 2 \times \\ 1 \times 7 & \bullet & \bullet & 2 \times 7 \\ 1 \times 6 & \bullet & \bullet & 2 \times 6 \\ 1 \times 5 & \bullet & \bullet & 2 \times 5 \\ 1 \times 4 & \bullet & \bullet & 2 \times 4 \\ 1 \times 3 & \bullet & \bullet & 2 \times 3 \\ 1 \times 2 & \bullet & \bullet & 2 \times 2 \\ 1 \times 1 & \bullet & \bullet & 2 \times 1 \end{array}$

Later: Note $1 \times n \xrightarrow{n \rightarrow \infty} 2 \times 1$ is a convergent sequence in this topology!

Ex Let X be an ordered set and $a \in X$.

Let $(a, \infty) = \{x \in X \mid x > a\}$

and $(-\infty, a) = \{x \in X \mid a < x\}$



be the open rays.

Show these are indeed open in the order topology.

Ans If X has a largest element b_0 , then

$(a, \infty) = (a, b_0]$ is a basis element, else

$(a, \infty) = \bigcup_{x > a} (a, x)$ is a union of basis elements.

Ex Do the open rays form a basis for \mathbb{R} ?

Ans No — consider $a < b$. No open ray is contained inside $(-\infty, b) \cap (a, \infty) = (a, b)$.

Ex Do the open rays form a subbasis for the order topology on X ?

Ans Yes.

They're open in the order topology, so the topology they generate is contained in the order topology.

Also, every basis element for the order topology is a finite intersection of open rays:

$$(a, b) = (-\infty, b) \cap (a, \infty)$$

$$(a, b_0] = (a, \infty) \quad \text{for } b_0 \text{ largest}$$

$$[a_0, b) = (-\infty, b) \quad \text{for } a_0 \text{ smallest}$$

So the reverse containment of topologies is also true.

Section 15: The product topology on $X \times Y$




Def For X and Y topological spaces, the product topology on $X \times Y$ is the topology generated by the basis \mathcal{B} with all sets of the form $U \times V$, with U open in X and V open in Y .

Check Is this a basis?

Note $X \times Y \in \mathcal{B}$.

Also, for $U_1 \times V_1, U_2 \times V_2 \in \mathcal{B}$,

$$(U_1 \times V_1) \cap (U_2 \times V_2) = (U_1 \cap U_2) \times (V_1 \cap V_2) \in \mathcal{B}$$




Question Is \mathcal{B} a topology?

No, the union $(U_1 \times V_1) \cup (U_2 \times V_2)$ is not in \mathcal{B}

Smaller bases are possible:

Thm If \mathcal{B} is a basis for X
and \mathcal{C} is a basis for Y ,
then $\mathcal{D} = \{B \times C \mid B \in \mathcal{B}, C \in \mathcal{C}\}$ is a basis for $X \times Y$.

PF



Check that \mathcal{D} is a basis. (easy)

Let W open in $X \times Y$. Let $x \times y \in W$.

By def of product top $\exists U$ open in X and $\exists V$ open in Y s.t. $x \times y \in U \times V \subset W$.

By def of basis, $\exists B \in \mathcal{B}$ and $C \in \mathcal{C}$ s.t.

$$x \times y \in B \times C \subset U \times V \subset W.$$

Therefore \mathcal{D} generates the product topology. \square

Given sets X, Y , there are projection maps

$$\begin{aligned} \pi_1: X \times Y &\longrightarrow X & \text{and} & \quad \pi_2: X \times Y \longrightarrow Y & \text{which are onto.} \\ (x, y) &\longmapsto x & & \quad (x, y) \longmapsto y \end{aligned}$$

Now assume X, Y are topological spaces.

Let U be an open set in X . $\pi_1^{-1}(U) = U \times Y$, which is open in $X \times Y$.

Similarly, for V open in Y $\pi_2^{-1}(V) = X \times V$, which is open in $X \times Y$.

Then $\pi_1^{-1}(U) \cap \pi_2^{-1}(V) = U \times V$

Theorem $\mathcal{S} = \{ \pi_1^{-1}(U) \mid U \text{ open in } X \} \cup \{ \pi_2^{-1}(V) \mid V \text{ open in } Y \}$
is a subbasis for the product topology on $X \times Y$.

Proof easy - see textbook. \square


Section 16: The subspace topology

Def Let (X, τ) be a topological space.

For $Y \subseteq X$, the collection


$$\tau_Y = \{U \cap Y \mid U \in \tau\}$$

is the subspace topology on Y .



Check it is a topology:

- $\emptyset = \emptyset \cap Y, \quad Y = X \cap Y \quad \checkmark$
- Arbitrary unions:
 $\bigcup_{\alpha \in J} (U_\alpha \cap Y) = (\bigcup_{\alpha \in J} U_\alpha) \cap Y \quad \checkmark$
- Finite intersections:
 $(U_1 \cap Y) \cap \dots \cap (U_n \cap Y) = (U_1 \cap \dots \cap U_n) \cap Y \quad \checkmark$



Ex Though $[0, 1)$ is not open in \mathbb{R} , it is open in the subspace topology on $[0, 2] \subset \mathbb{R}$.




Lemma If Y is an open subset of X then for $U \subset Y$,
 $U \in \tau_Y \Leftrightarrow U \in \tau$.
(open in Y) (open in X)

Proof (\Rightarrow) $U \in \tau_Y \Rightarrow \exists V \in \tau$ s.t. $U = V \cap Y \Rightarrow U \in \tau$

(\Leftarrow) Assume $U \in \tau$. Since $U = U \cap Y$, $U \in \tau_Y$. \square

Lemma If \mathcal{B} is a basis for the topology on X , then $\mathcal{B}_Y = \{B \cap Y \mid B \in \mathcal{B}\}$ is a basis for the topology on Y .



PS Given $U \cap Y$ open in Y (with U open in X)


and $y \in U \cap Y$, we can find $B \in \mathcal{B}$ with $y \in B \subset U$.

Note $y \in B \cap Y \subset U \cap Y$.

It follows from Lemma 13.2

that \mathcal{B}_Y is a basis for the topology on Y .

Thm If $A \subseteq X$ and $B \subseteq Y$, then the product topology on $A \times B$ is the same as the subspace topology on $A \times B \subset X \times Y$.



PS Consider first the product topology on the larger space $X \times Y$, which has as a basis all $U \times V$, U open in X , V open in Y .


So the subspace topology on $A \times B$ has as a basis all $(U \times V) \cap (A \times B) = (U \cap A) \times (V \cap B)$, which is a basis for the product topology on $A \times B$.

These topologies are the same since they have a common basis.

Rmk The order and subspace topologies are not compatible in general.

For example, let $Y = [0, 1) \cup \{2\} \subset \mathbb{R}$.

In the subspace topology, $\{2\}$ is open in Y .



But in the order topology, any basis element containing 2 is of the form

$(a, 2] := \{y \in Y \mid a < y \leq 2\}$ for some $a \in Y$
and it follows that $\{2\}$ is not open.


Def If X is totally ordered, a subset $Y \subset X$ is convex if $\forall a, b \in Y$ with $a < b$, the interval $(a, b) = \{x \in X \mid a < x < b\}$ is contained in Y .

Thm If X is an ordered set with the order topology and $Y \subset X$ is convex, then the order and subspace topologies on Y agree.

Section 17: Closed sets and limit points


Def A subset A of a topological space X is closed if $X-A$ is open.

Ex $[a, b]$ is closed in \mathbb{R} since $\mathbb{R} - [a, b] = (-\infty, a) \cup (b, \infty)$ is open.



Ex $[a, b] \times [c, d]$ is closed in \mathbb{R}^2 .

(Complement is union of four basic open sets.)



Ex In the finite complement topology on a set X , the closed sets are X , \emptyset , and all finite subsets of X .

Ex In the discrete topology, every set is closed.

Rmk closed \neq not open

Ex $[0, 2)$ is neither open nor closed in \mathbb{R} .

Ex Let $Y = [0, 2) \cup \{4\} \subset \mathbb{R}$ have the subspace topology.

Is $[0, 2)$ open in Y ? *Yes.*

Is $\{4\}$ open in Y ? *Yes.*

Is $[0, 2)$ closed in Y ? *Yes.*

Is $\{4\}$ closed in Y ? *Yes.*


Thm For X a topological space,

- \emptyset and X are closed
- arbitrary intersections of closed sets are closed
- finite unions of closed sets are closed.

Pf See book. $\left(X - \bigcap_{\alpha \in J} C_{\alpha} = \bigcup_{\alpha \in J} (X - C_{\alpha}) \right)$

Rmk Topological spaces could have instead been defined via closed sets.

Thm For $Y \subset X$ with the subspace topology, a set $A \subset Y$ is closed in $Y \iff A = B \cap Y$ for some closed set B in X




$X = \mathbb{R}^2$

Pf See book.

Def For X a topological space and $A \subset X$,

- the interior of A , denoted $\text{Int } A$, is the union of all open sets contained in A
- the closure of A , denoted $\text{Cl } A$ or \bar{A} , is the intersection of all closed sets containing A .




Ex For $X = \mathbb{R}$ and $A = [0, 2)$,
 $\text{Int } A = (0, 2)$ and $\bar{A} = [0, 2]$.

Thm X topological space with basis \mathcal{B} , $A \subset X$.

(a) $x \in \bar{A} \iff$ every open set containing x intersects A .

(b) $x \in \bar{A} \iff$ every $B \in \mathcal{B}$ containing x intersects A .



Munkres calls these neighborhoods of x .

Rmk An open set containing x is called an open neighborhood of x .
A set containing an open neighborhood of x is called a neighborhood of x .

Pf (a) (\implies) Assume \exists an open neighborhood U of x that doesn't intersect A

$\implies X - U$ is a closed set containing A

$\implies \bar{A} \subset X - U \implies x \notin \bar{A}$

(\impliedby) $x \notin \bar{A}$ means $X - \bar{A}$ is an open nbhd of x not intersecting A

(b) Use (a).

(\Rightarrow) Basis elements are open

(\Leftarrow) An open nbhd containing x contains a basis element containing x .

Def X topological space, $A \subset X$. A point $x \in X$ is a limit point of A if every open nbhd of x contains a point in A other than x .

(x may or may not be in A)

Ex $A = [0, 2) \subset \mathbb{R}$. The set of limit points is $[0, 2]$.

Ex $B = \{ \frac{1}{n} \mid n \in \mathbb{Z}_+ \} \subset \mathbb{R}$. The only limit point is 0.

Ex $\mathbb{Q} \subset \mathbb{R}$. The set of limit points is \mathbb{R} .

Thm X topological space, $A \subset X$.

Let A' be the set of limit points of A .

Then $\bar{A} = A \cup A'$.

Cor A subset of a topological space is closed \Leftrightarrow it contains all its limit points.

Def A topological space X is a Hausdorff space if
 \forall distinct $x, y \in X$, \exists open neighborhoods U of x
 and V of y with $U \cap V = \emptyset$



Thm In a Hausdorff space X , finite sets
 are closed.

Pf It suffices to show that $\{x\}$ is closed $\forall x \in X$,
 since finite unions of closed sets are closed.

So, let $y \neq x$ in X . By the Hausdorff assumption,
 \exists open neighborhood $V \ni y$ with $x \notin V$.




So y is not a limit point of $\{x\}$.
 By Theorem above, $y \notin \overline{\{x\}}$.


So $\overline{\{x\}} = \{x\}$, meaning $\{x\}$ is closed. \square

Def A sequence x_1, x_2, x_3, \dots converges to a point
 $x \in X$ if \forall open neighborhoods U of x , $\exists N \in \mathbb{Z}_+$
 such that $x_n \in U \quad \forall n \geq N$.

Ex $\frac{1}{4}, \frac{1}{3}, \frac{1}{2}, \dots \rightarrow 0$ in \mathbb{R}



Ex In the topological space




note that $\{b\}$ is not closed,

and note that the sequence b, b, b, b, b, \dots

converges not only to b , but also to a or to c !

Thm In a Hausdorff space, sequences converge to at most one point.


PS If $x_n \rightarrow x$ and $y \neq x$, then let $U \ni x$ and $V \ni y$ be disjoint nbhds. Note U contains all but finitely many elements of the sequence, and hence V cannot.



Thm A subspace of a Hausdorff space is Hausdorff.
The product of two Hausdorff spaces is Hausdorff.

Section 18: Continuous functions

Def X, Y topological spaces. A function $f: X \rightarrow Y$ is continuous if \forall open U in Y , $f^{-1}(U)$ is open in X .



Rmk It suffices to check this condition on basis elements of Y :

$$U = \bigcup_{\alpha \in J} B_{\alpha} \quad f^{-1}(U) = f^{-1}\left(\bigcup_{\alpha \in J} B_{\alpha}\right) = \bigcup_{\alpha \in J} \underbrace{f^{-1}(B_{\alpha})}_{\text{open}} \\ \underbrace{\qquad\qquad\qquad}_{\text{open}}$$

Rmk It suffices to check this condition on subbasis elements of Y :


$$B = S_1 \wedge \dots \wedge S_n \quad f^{-1}(B) = f^{-1}(S_1 \wedge \dots \wedge S_n) = f^{-1}(S_1) \wedge \dots \wedge f^{-1}(S_n)$$

Ex $\text{Id}: \mathbb{R}_e \rightarrow \mathbb{R}$ (defined by $\text{Id}(x) = x \quad \forall x \in \mathbb{R}_e$) is continuous since $\text{Id}^{-1}((a,b)) = (a,b)$ is open in \mathbb{R}_e .

$\text{Id}: \mathbb{R} \rightarrow \mathbb{R}_e$ is not continuous since $\text{Id}^{-1}([a,b)) = [a,b)$ is not open in \mathbb{R} .

Thm Let X and Y be topological spaces, and let $f: X \rightarrow Y$.
The following are equivalent:

- (1) f is continuous
- (2) $\forall A \subset X, f(\overline{A}) \subset \overline{f(A)}$
- (3) \forall closed sets B in $Y, f^{-1}(B)$ is closed in X .
- (4) $\forall x \in X$ and open nghds V of $f(x) \exists$ open nghd U of x s.t. $f(U) \subset V$
- (5) $\forall x \in X$ and basic open nghd V of $f(x) \exists$ basic open nghd U of x s.t. $f(U) \subset V$.



If X, Y are metric space then
(5) agrees with the ϵ - δ definition
of continuity.

Rmk If (4) holds at $x_0 \in X$ but not necessarily at all
points in X , then we say f is continuous at x_0

Proof We will show that (1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (1) and (1) \Rightarrow (4) \Rightarrow (5) \Rightarrow (1)

(1) \Rightarrow (2) Let $A \subset X$. Let $x \in \overline{A}$. Want: $f(x) \in \overline{f(A)}$.

Let V be an open nghd of $f(x)$. Since f cont, $f^{-1}(V)$ open X .

Since $x \in f^{-1}(V)$, $f^{-1}(V)$ is an open nghd of x .

Since $x \in \overline{A}$, $f^{-1}(V) \cap A \neq \emptyset$. Let $x' \in f^{-1}(V) \cap A$.

Then $f(x') \in V \cap f(A)$. Therefore $f(x) \in \overline{f(A)}$.

(2) \Rightarrow (3) Let B closed Y . Let $A = f^{-1}(B)$. Want: A is closed. i.e. $\overline{A} = A$.
 $f(A) = f(f^{-1}(B)) \subset B$. Let $x \in \overline{A}$.

Then $f(x) \in f(\overline{A}) \subset \overline{f(A)} \subset \overline{B} = B$. Thus $x \in f^{-1}(B) = A$.

Hence $\overline{A} \subset A$. Therefore $\overline{A} = A$.

(3) \Rightarrow (1) Let $U \subset_{\text{open}} Y$. Then $U^c = X - U \subset_{\text{closed}} Y$.

Hence $(f^{-1}(U))^c = f^{-1}(U^c) \subset_{\text{closed}} X$. Therefore $f^{-1}(U) \subset_{\text{open}} X$.

(1) \Rightarrow (4) Let $x \in X$. Let V be an open nghd of $f(x)$.

Let $U = f^{-1}(V)$. U is an open nghd of x s.t. $f(U) \subset V$.

(4) \Rightarrow (5) Let $x \in X$. Let C be a basic open nghd of $f(x)$.

Then \exists open nghd U of x s.t. $f(U) \subset C$.

Hence \exists basic open nghd B of x s.t. $B \subset U$.

Therefore $f(B) \subset f(U) \subset C$.

(5) \Rightarrow (1) Let $U \subset_{\text{open}} Y$. If $f^{-1}(U) = \emptyset$ we are done.


Assume $f(x) \in U$. Then \exists basic open nghd C of $f(x)$ with $C \subset U$.

Thus \exists basic open nghd B of x with $f(B) \subset C \subset U$.

Therefore $B \subset f^{-1}(U)$. Hence $f^{-1}(U) \subset_{\text{open}} X$.

Def A homeomorphism is a continuous bijection $f: X \rightarrow Y$ such that $f^{-1}: Y \rightarrow X$ is also continuous.


We say "X is homeomorphic to Y" and write $X \cong Y$.



Ex $f: (-1, 1) \rightarrow \mathbb{R}$ defined by $f(x) = \frac{x}{1-x^2}$ is a homeomorphism with inverse $f^{-1}: \mathbb{R} \rightarrow (-1, 1)$ defined by $f^{-1}(y) = \frac{2y}{1 + \sqrt{1+4y^2}}$.

So homeomorphisms need not preserve boundedness.

Non-Ex $f: [0, 2\pi) \rightarrow S^1$ defined by $f(t) = (\cos t, \sin t)$ is a continuous bijection that is not a homeomorphism.




Rmk A homeomorphism gives a bijection b/w the open sets of X and Y. So it preserves all topological properties.


Def An embedding $f: X \rightarrow Y$ is a continuous injective map s.t. $f: X \rightarrow f(X)$ is a homeomorphism
 ↑ image of f with subspace top. Munkres: embedding

Thm (Constructing continuous functions)


(a) Constant functions are cont.



(b) The inclusion of a subspace is cont.




(c) Compositions are continuous: If $f: X \rightarrow Y$, $g: Y \rightarrow Z$ are cont., then so is $g \circ f: X \rightarrow Z$.



(d) $f: X \rightarrow Y$ cont. and $A \subset X \Rightarrow f|_A$ cont.

Restrict domain




(e) $f: X \rightarrow Y$ cont. \Rightarrow $f: X \rightarrow Z$ cont. for $Y \subset Z$ (Extend codomain)
 \Rightarrow $f: X \rightarrow W$ cont. for $f(X) \subset W$ (Restrict codomain)

(f) $f: X \rightarrow Y$, $X = \bigcup_{\alpha} U_{\alpha}$, $f|_{U_{\alpha}}$ cont $\forall \alpha \Rightarrow f$ cont.

Locality of continuity

open




(g) (Pasting lemma) $X = A \cup B$, A, B closed in X .

$f: A \rightarrow Y$ and $g: B \rightarrow Y$ cont. and $f(x) = g(x) \forall x \in A \cap B$.

Then the function $h: X \rightarrow Y$ defined via

$$h(x) = \begin{cases} f(x) & x \in A \\ g(x) & x \in B \end{cases}$$

is continuous.



Proof: Exercise / see book.

Ex Why is $f: \mathbb{R} \rightarrow \mathbb{R}$ not continuous?





Thm Let $f: A \rightarrow X \times Y$ be given by $f(a) = (f_1(a), f_2(a))$.
 Then f is continuous $\iff f_1, f_2$ are continuous.

Pf Let $\pi_1: X \times Y \rightarrow X$ and

$$\pi_2: X \times Y \rightarrow Y.$$


Note π_1 is continuous since if U is open in X , then $\pi_1^{-1}(U) = U \times Y$ is open in $X \times Y$.
 And similarly for π_2 .

(\implies) f cont. implies $\pi_1 \circ f = f_1$ and $\pi_2 \circ f = f_2$ are continuous.



(\impliedby) For $U \times V$ a basic open set in $X \times Y$ (meaning U open in X , V open in Y), note $f^{-1}(U \times V) = f_1^{-1}(U) \cap f_2^{-1}(V)$ is open in A

open in A since f_1 cont. open in A since f_2 cont.



Section 19 Product topology

Recall

Def Given $\{X_\alpha\}_{\alpha \in J}$, the cartesian product $\prod_{\alpha \in J} X_\alpha$ is the set of all J-tuples $(x_\alpha)_{\alpha \in J}$ which are maps $x: J \rightarrow \bigcup_{\alpha \in J} X_\alpha$ with $x_\alpha := x(\alpha) \in X_\alpha$.

For each $\alpha \in J$ we have a "projection map" $\pi_\alpha: \prod_{\alpha \in J} X_\alpha \rightarrow X_\alpha$
 $x \mapsto x_\alpha$

Def The (less-important) box topology on $\prod_{\alpha \in J} X_\alpha$ has as its basis all sets $\left\{ \prod_{\alpha \in J} U_\alpha \mid U_\alpha \text{ open in } X_\alpha \forall \alpha \right\}$


Def The (more-important) product topology on $\prod_{\alpha \in J} X_\alpha$ has subbasis given by $\left\{ \pi_\alpha^{-1}(U) \mid \alpha \in J, U \subset_{\text{open}} X_\alpha \right\}$.

Remark This subbasis consists of exactly the subsets of $\prod_{\alpha \in J} X_\alpha$ that need to be open for the projection maps π_α to be continuous.

The product topology on $\prod_{\alpha \in J} X_\alpha$ has basis

$$\left\{ \prod_{\alpha \in J} U_\alpha \mid \begin{array}{l} U_\alpha \text{ open in } X_\alpha \\ U_\alpha = X_\alpha \text{ for all but finitely many } \alpha \end{array} \right\}$$

Rmk These topologies agree if J is finite.



If J infinite then the box topology is finer.

The previous theorem is a special case of the following.

Thm Let $f_\alpha: A \rightarrow X_\alpha \quad \forall \alpha \in J$

Define $f: A \rightarrow \prod_{\alpha \in J} X_\alpha$ by $a \mapsto (f_\alpha(a))_{\alpha \in J}$.

Let $\prod X_\alpha$ have the product topology.

Then f is continuous $\iff f_\alpha$ is continuous $\forall \alpha$.

Recall each projection $\pi_\beta: \prod_{\alpha \in J} X_\alpha \rightarrow X_\beta$ is continuous.

Observe: $\pi_\alpha \circ f = f_\alpha$

(\implies) f cont. $\implies f_\alpha = \pi_\alpha \circ f$ cont. $\forall \alpha$

(\impliedby) Consider a subbasic open set $\pi_\alpha^{-1}(U)$, $U \subset_{\text{open}} X_\alpha$

$f^{-1}(\pi_\alpha^{-1}(U)) = f_\alpha^{-1}(U)$ open since f_α is continuous \square

Important fact:

Rmk (\impliedby) need not be true if $\prod X_\alpha$ has the box topology. Let $\mathbb{R}^\omega = \prod_{n \in \mathbb{Z}^+} X_n$ with $X_n = \mathbb{R} \quad \forall n$.

Define $f: \mathbb{R} \rightarrow \mathbb{R}^\omega$ by $f(t) = (t, t, t, \dots)$

Each coordinate function $f_n: \mathbb{R} \rightarrow \mathbb{R}$ by $f_n(t) = t$ is continuous.

But, f is not continuous if \mathbb{R}^ω has the box topology,

since $B = (-1, 1) \times (-\frac{1}{2}, \frac{1}{2}) \times (-\frac{1}{3}, \frac{1}{3}) \times \dots$ is open in the

box topology, but $f^{-1}(B) = \{0\}$ is not open in \mathbb{R} .

Section 20: The metric topology

Def A metric on a set X is a function $d: X \times X \rightarrow \mathbb{R}$ s.t.



(1) $d(x,y) \geq 0$, $d(x,y) = 0 \Leftrightarrow x=y$

(2) $d(x,y) = d(y,x)$

triangle inequality


(3) $d(x,z) \leq d(x,y) + d(y,z)$

$B_r^d(x) = \{y \in X \mid d(x,y) < r\}$ is the
" r -ball centered at x .



Def Given a metric space (X,d) , $\{B_r(x) \mid x \in X, r > 0\}$ is a basis for a metric topology on X .

Check its a basis (2)



Rmk U is open in (X,d)

$\Leftrightarrow \forall x \in U \quad \exists x \in B_r(y) \subset U$

$\Leftrightarrow \forall x \in U \quad \exists x \in B_r(x) \subset U$

Def A topological space X is metrizable if \exists a metric on X that induces the topology on X .

Important Question Is a given topological space metrizable?

Ex For X a set, defining $d(x,y) = \begin{cases} 1 & x \neq y \\ 0 & x = y \end{cases}$

gives a metric inducing the discrete topology.

Ex Metrics on \mathbb{R}^n

For $1 \leq p \leq \infty$, let $d_p(x,y) = \|x-y\|_p$, where

where $\|x\|_2 = \sqrt{x_1^2 + \dots + x_n^2}$

$\|x\|_1 = |x_1| + \dots + |x_n|$

$\|x\|_\infty = \max\{|x_1|, \dots, |x_n|\}$



"taxicab" metric




"sup" metric

$\|x\|_p = (|x_1|^p + \dots + |x_n|^p)^{1/p}$ for all $1 \leq p < \infty$.

Note $B_1^{d_\infty}(\vec{0}) \supset B_1^{d_2}(\vec{0}) \supset B_1^{d_1}(\vec{0}) \supset B_{1/2}^{d_\infty}(\vec{0})$.


Hence the following lemma shows that all of these metrics induce the same topology on \mathbb{R}^n :



(and, moreover, this topology is the product topology)

Lemma Let X have metrics d, d' generating the topologies τ, τ' . Then τ' is finer than τ (i.e. $\tau = \tau'$) if $\forall B_r^d(x), \exists B_s^{d'}(x) \subset B_r^d(x)$.

Pf See book.



Def Given a metric space (X, d) , define $\bar{d}(x, y) = \min(d(x, y), 1)$. This is the standard bounded metric.

Thm \bar{d} is metric on X and induces the same topology as d .

Pf See book.

Cor Boundedness (a set having finite diameter) is a metric property but not a topological property.

Metrics for \mathbb{R}^n Ideas: $d(x, y) = \left[\sum_{i=1}^n (x_i - y_i)^2 \right]^{1/2}$, $\rho(x, y) = \sup_n |x_n - y_n|$

Problem: Make take value ∞ .

Def The uniform metric on \mathbb{R}^J is given by $\bar{\rho}(x, y) = \sup_{\alpha \in J} \{ \bar{d}(x_\alpha, y_\alpha) \}$

(Recall: $\bar{d}(a, b) = \min(|a - b|, 1)$)

Theorem

- (a) Uniform top on \mathbb{R}^J finer than product topology.
- (b) Uniform top on \mathbb{R}^J coarser than box topology.
- (c) For J infinite, all 3 are distinct.

Proof Let $x \in \mathbb{R}^J$.

(a) Let $\prod_{\alpha \in J} U_\alpha$ be a basic open nhd of x in prod. top.

Let $\alpha_1, \dots, \alpha_n$ be the indices for which $U_{\alpha} \neq \mathbb{R}$.

For $1 \leq i \leq n$, choose ε_i s.t. $B_{\varepsilon_i}^{\bar{d}}(x_i) \subset U_{\alpha_i}$.

Let $\varepsilon = \min \{ \varepsilon_1, \dots, \varepsilon_n \}$. Claim: $B_{\varepsilon}^{\bar{d}}(x) \subset \bigcap_{\alpha \in J} U_{\alpha}$.

Let $z \in \mathbb{R}^{\omega}$ s.t. $\bar{p}(x, z) < \varepsilon$. Then $\forall \alpha \in J$, $\bar{d}(x_{\alpha}, z_{\alpha}) < \varepsilon$.

$\therefore z \in \bigcap_{\alpha \in J} U_{\alpha}$. \checkmark

(b) Consider $B_{\varepsilon}^{\bar{d}}(x)$. Let $U = \prod_{\alpha \in J} (x_{\alpha} - \frac{\varepsilon}{2}, x_{\alpha} + \frac{\varepsilon}{2})$.

Then $x \in U$. Claim: $U \subset B_{\varepsilon}^{\bar{d}}(x)$. Let $y \in U$.

Then $\forall \alpha \in J$, $\bar{d}(x_{\alpha}, y_{\alpha}) < \frac{\varepsilon}{2}$. So $\bar{p}(x, y) \leq \bigvee_{\alpha} \frac{\varepsilon}{2}$. \checkmark

(c) Let $\underline{0}$ be the J -tuple with all coordinates equal to 0.

Then $B_{\frac{1}{2}}^{\bar{d}}(\underline{0})$ is open in uniform top but not in the product topology.

There are subsets of \mathbb{R}^J that are open in the box topology but not the uniform topology. Exercise. (HWK 4). \square

Thm The product topology on \mathbb{R}^{ω} is induced by the metric

$$D(x, y) = \sup \left\{ \frac{\bar{d}(x_i, y_i)}{i} \right\}$$

Pf See book.


Rmk More generally, countable products of metric spaces are metrizable. You will show this in HW4.

Rmk $\sup \left\{ \frac{\bar{d}(x_i, y_i)}{i} \right\}$ is not a metric

It can take the value ∞ .


Section 21: Metric Topology (continued)

Rmk Metric spaces are Hausdorff: If $x \neq y$, then $B_\varepsilon(x)$ and $B_\varepsilon(y)$ are disjoint for $0 < \varepsilon < \frac{1}{2}d(x,y)$ by the triangle inequality.




Thm $f: (X, d_x) \rightarrow (Y, d_y)$ continuous
 \iff

Given $x \in X$, $\varepsilon > 0$, $\exists \delta > 0$ s.t.
 $d_x(x, x') < \delta \implies d_y(f(x), f(x')) < \varepsilon$.



Lemma (Sequence Lemma) X topological space, $A \subset X$.
 If a sequence in A converges to x , then $x \in \bar{A}$.
 Converse holds if X metrizable.


Pf (\implies) $x_n \rightarrow x$ implies every nbhd of x contains a point in A , so $x \in \bar{A}$.



Pf (\impliedby) Let d be a metric giving the topology on X .
 $\forall n \in \mathbb{Z}_+$, choose $x_n \in B_{1/n}(x) \cap A$. Note $x_n \rightarrow x$.

Rmk For the converse of this (and the next) lemma, assumption " X metrizable" can be relaxed to X first countable, which means:

$\forall x \in X$, \exists countable collection of nbhds $\{U_n\}_{n \in \mathbb{Z}_+}$ such that
 \forall nbhds $U \ni x$, $\exists n \in \mathbb{Z}_+$ with $x \in U_n \subset U$.



Preview: Leave in notes, but skip in class.

Cor A space X with a subset $A \subset X$ and $x \in \bar{A}$ and no sequence in A converging to x is not metrizable.

Example $\mathbb{R}^{\mathbb{J}}$ not metrizable for \mathbb{J} uncountable. Indeed, let $A = \{x = (x_\alpha) \in \mathbb{R}^{\mathbb{J}} \mid x_\alpha = 1 \text{ for all but finitely many } \alpha \in \mathbb{J}\}$.

Define $\vec{0} \in \mathbb{R}^{\mathbb{J}}$ to be the point x with $x_\alpha = 0 \forall \alpha \in \mathbb{J}$.

Then $\vec{0} \in \bar{A}$ since any basic open set about $\vec{0}$ is \mathbb{R} in all but finitely many coordinates, hence intersects A .

But for any sequence $x^1, x^2, x^3, \dots \in A$,


\exists some $\beta \in \mathbb{J}$ with $x_\beta^n = 1 \forall n$ (since a countable union of finite sets is countable),

hence $\pi_\beta^{-1}\left(\left(-\frac{1}{2}, \frac{1}{2}\right)\right)$ is a nbhd about $\vec{0}$ containing no x^n , so no sequence in A can converge to $\vec{0}$.

Thm X, Y topological spaces, $f: X \rightarrow Y$. If f is continuous, then $\forall x_n \rightarrow x$, we have $f(x_n) \rightarrow f(x)$.

Converse holds if X is a metric space.

Pf (\Rightarrow) Given nbhd $V \ni f(x)$, note $f^{-1}(V)$ is a nbhd of x , so x_n eventually in $f^{-1}(V)$ implies that $f(x_n)$ is eventually in V .



(\Leftarrow) Suffices to show $f(\bar{A}) \subset \overline{f(A)}$ for any $A \subset X$.

If $x \in \bar{A}$, then by prior lemma (since X metrizable), $\exists x_n \in A$ with $x_n \rightarrow x$. By assumption, $f(x_n) \rightarrow f(x)$.

Since $f(x_n) \in \overline{f(A)} \forall n$, the prior lemma gives $f(x) \in \overline{f(A)}$.



Hence $f(\bar{A}) \subset \overline{f(A)}$ as desired.

Section 22: The quotient topology

Let X be a topological space, and let X^* be a partition of X , namely a collection of disjoint subsets whose union is X .


(In other words, suppose we have an equivalence relation on X .)

Ex $[0,1] \times [0,1] / \sim$




torus

$[0,1] \times [0,1] / \sim'$



Klein bottle

D^2 / S^1



sphere

From the topology on X , how do we get a topology on X^* ?
 Give X^* the finest topology such that $p: X \rightarrow X^*$
 is continuous. $x \mapsto [x]$

U open in $X^* \iff p^{-1}(U)$ open in X . (Coarsest such topology would give only the open sets \emptyset, X^*)

Def Let X be a topological space, Y be a set, $p: X \rightarrow Y$ be surjective. In the quotient topology on Y ,

U open in $Y \iff p^{-1}(U)$ open in X .

Check This is a topology.

$$p^{-1}(Y) = X \text{ open in } X \Rightarrow Y \text{ open in } Y,$$


$$p^{-1}(\emptyset) = \emptyset \text{ open in } X \Rightarrow \emptyset \text{ open in } Y.$$

$$p^{-1}\left(\bigcup_{\alpha} U_{\alpha}\right) = \bigcup_{\alpha} p^{-1}(U_{\alpha}) \text{ open in } X \Rightarrow \bigcup_{\alpha} U_{\alpha} \text{ open in } Y.$$

open in $Y \Rightarrow$ open in X

$$p^{-1}\left(\bigcap_{i=1}^n U_i\right) = \bigcap_{i=1}^n p^{-1}(U_i) \text{ open in } X \Rightarrow \bigcap_{i=1}^n U_i \text{ open in } Y.$$

Ex



torus




sphere



Ex $p: \mathbb{R} \rightarrow \{-1, 0, 1\}$ by $p(x) = \begin{cases} -1 & \text{if } x < 0 \\ 0 & \text{if } x = 0 \\ 1 & \text{if } x > 0 \end{cases}$

The induced quotient topology on $\{-1, 0, 1\}$ is




Let $p: X \rightarrow Y$ be surjective, X a topological space.
 What if Y already has a topology?

Def For X, Y topological spaces and $p: X \rightarrow Y$ surjective,
 p is a quotient map if
 U open in $Y \iff p^{-1}(U)$ open in X .

Ex All the examples above, where Y has quotient topology.

Non-Ex



$X = \{(x, y) \in \mathbb{R}^2 \mid xy = 1\} \cup \{(0, 0)\}$
 with the subspace topology

$Y = \mathbb{R}$

$p(x, y) = x$

Note $p: X \rightarrow Y$ is continuous and surjective, but not a quotient map since $p^{-1}(\{0\}) = \{(0, 0)\}$ is open in X ,
 but $\{0\}$ is not open in Y .

Remark $p: X \rightarrow Y$ is a quotient map if

- $U \subseteq_{\text{open}} Y \Rightarrow p^{-1}(U) \subseteq_{\text{open}} X$ i.e. p continuous, and
- $p^{-1}(U) \subseteq_{\text{open}} X \Rightarrow U \subseteq_{\text{open}} Y$ i.e. p sends open sets that are complete inverse images of a subset of Y to an open set.
 called saturated

Remark The product of quotient maps need not be a quotient map.
 See Example 7 in Ch 22 in Munkres.


Thm (Continuous maps out of quotient spaces)

X, Y, Z topological spaces, $p: X \rightarrow Y$ a quotient map.


Let $g: X \rightarrow Z$ be constant on each $p^{-1}(\{y\})$,

hence inducing a function $f: Y \rightarrow Z$ with $f \circ p = g$.

Then f continuous $\iff g$ continuous.



Pf (\implies) f cont. implies $f \circ p = g$ cont.



(\impliedby) Given V open in Z , $g^{-1}(V)$ open in X since g is continuous.
 $p^{-1}(f^{-1}(V))$

Now, p a quotient map implies $f^{-1}(V)$ open in Y , so f is continuous.