


# MAP 2302, Exam I, Fall 2015

Name: \_\_\_\_\_


Student signature: \_\_\_\_\_

Turn in all relevant work with final answers circled on separate sheets. Full work is required for full credit.

(1) [25] The graph of a function  $f$  is shown below.



- (a) [10] Sketch a direction field for the differential equation  $\frac{dy}{dx} = f(y)$ . Your sketch must show points with  $0 \leq x \leq 3$  and  $-2 \leq y \leq 2$  and show directions at each integer point. Slopes are computed by plugging the  $y$ -coordinate of the point into  $f$ . These are shown on the graph below.
- (b) [10] Estimate the constant solutions of the above differential equation. Draw them on your direction field. Constant solutions occur when  $\frac{dy}{dx} = 0$  or equivalently  $f(y) = 0$ . This happens when  $y \approx \pm 1.5$ . These are drawn in red on the graph below.
- (c) [5] Use your direction field to sketch two solution curves, one satisfying  $y(0) = 2$  and the other satisfying  $y(0) = 0$ . With the direction field drawn, we need only follow the arrows and avoid crossing already computed solutions. These are shown in blue on the below graph.



(2) [25] Find the most general family of solutions to the equation [15]

$$x \frac{dy}{dx} - 3y = x^4$$

What are the singular points of the equation [5]? Does the equation have a unique solution satisfying  $y(0) = 0$  [5]? Justify your answer.

First, we have to put this linear equation in standard form

$$\frac{dy}{dx} - \frac{3}{x}y = x^3$$

Since we have a singular point at  $x = 0$ , we can only solve on either  $(0, \infty)$  or  $(-\infty, 0)$ . In either case, an integrating factor is given by  $\mu(x) = e^{\int (-3/x) dx} = e^{\ln|x|^{-3} + C} = C|x|^{-3} = Cx^{-3}$ . We may choose  $C = 1$  so that  $\mu(x) = x^{-3}$  regardless of whether  $x > 0$  or  $x < 0$ . Then we can multiply through by the integrating factor to obtain.

$$\begin{aligned} x^{-3} \frac{dy}{dx} - 3x^{-4}y &= 1 \\ \frac{d}{dx} (x^{-3}y) &= 1 \\ x^{-3}y &= x + C \\ y &= x^4 + Cx^3 \end{aligned}$$

Since we have solved independently for  $x < 0$  and  $x > 0$  (albeit with the same technique) we can choose different constants for  $x < 0$  and  $x > 0$  so the most general family of solutions is given by the piecewise defined family

$$y = \begin{cases} x^4 + Cx^3 & \text{if } x > 0 \\ x^4 + Dx^3 & \text{if } x \leq 0 \end{cases}$$

Observe that this is well-defined at 0 as both sides would have  $y(0) = 0$  regardless of choice of  $C, D$ .

On the exam, I gave full credit for saying  $y = x^4 + Cx^3$  was the family of solutions,  $x = 0$  was a singular point, and the solution was not unique because  $y(0) = 0$  regardless of choice of  $C$ . Note that the full solution is a bit more complicated.

(3) [25] Find a solution to the IVP

$$x \frac{dy}{dx} = y + \sqrt{x^2 - y^2} ; y(1) = 0$$

You may use the fact that  $\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1}(x)$  without work.

After exhausting all other methods, we realize that the equation is homogeneous and use the substitution  $y = ux$  so that  $\frac{dy}{dx} = u + x \frac{du}{dx}$ . We will rewrite the equation by dividing by  $x$ . Once again, by doing this we will have to choose to solve on  $x > 0$  or  $x < 0$ . Since our initial value is given at  $x = 1$  we will choose  $x > 0$ . Division yields

$$\frac{dy}{dx} = \frac{y}{x} + \frac{\sqrt{x^2 - y^2}}{x}.$$

After substitution we have

$$\begin{aligned} u + x \frac{du}{dx} &= u + \frac{\sqrt{x^2 - x^2u^2}}{x} \\ x \frac{du}{dx} &= \frac{|x|\sqrt{1-u^2}}{x}. \end{aligned}$$

Since  $x > 0$ ,  $|x| = x$  and  $|x|/x = 1$ . After separating we have.

$$\begin{aligned} \frac{1}{\sqrt{1-u^2}} du &= \frac{1}{x} dx \\ \sin^{-1}(u) &= \ln(x) + C. \end{aligned}$$

In the last step, we have again used our assumption that  $x > 0$ . Then changing back from  $u$  to  $y$  we have

$$\sin^{-1}(y/x) = \ln(x) + C.$$

With the initial condition  $y(1) = 0$  we find  $C = 0$  so the solution is

$$\sin^{-1}(y/x) = \ln(x).$$

(4) [25] Find the most general family of solutions to the equation

$$(3x^2 + y) dx + (x^2y - x) dy = 0.$$

We can solve this using integrating factors for exact equations. Letting  $M = 3x^2 + y$  and  $N = x^2y - x$  we have that  $M_y = 1$  and  $N_x = 2xy - 1$ . Then

$$\frac{M_y - N_x}{N} = \frac{2 - 2xy}{x^2y - x} = \frac{-2(xy - 1)}{x(xy - 1)} = \frac{-2}{x}.$$

Consequently, we can find our integrating factor as

$$\mu(x) = e^{\int(-2/x)dx} = x^{-2}.$$

After multiplying through by  $\mu$  we have the equation

$$(3 + x^{-2}y) dx + (y - x^{-1}) dy = 0.$$

This equation is exact. Then we look for solutions of the form  $F(x, y) = C$ . We know that  $F_x = 3 + x^{-2}y$  and  $F_y = y - x^{-1}$ . Integrating  $F_x$  we get

$$F = 3x - x^{-1}y + g(y).$$

Taking the  $y$  partial derivative we have

$$F_y = x^{-1}y + g'(y)$$

Since  $F_y = y - x^{-1}$

$$y - x^{-1} = x^{-1}y + g'(y)$$

$$g'(y) = y$$

$$g(y) = \frac{y^2}{2}$$

Then  $F(x, y) = 3x - x^{-1}y + \frac{y^2}{2}$  and solutions are given implicitly by

$$3x - x^{-1}y + \frac{y^2}{2} = C.$$