

A General Accounting Framework for Ecological Systems: A Functional Taxonomy for Connectivist Ecology

BRUCE HANNON

*Department of Geography and Illinois Natural History Survey,
University of Illinois, Urbana, Illinois 61801*

AND

ROBERT COSTANZA AND ROBERT ULANOWICZ

*Chesapeake Biological Laboratory, Center for Environmental and
Estuarine Studies, University of Maryland,
Solomons, Maryland 20688-0038*

Received February 1, 1990

Accounting of material and energy flows has long been an important tool in ecosystem ecology. But each material is usually handled separately and independently. The connections *between* materials, energy, plants, animals, etc. have not been incorporated into the accounting framework, and "service" or information flows (such as flower pollination by bees) are usually ignored. We develop a general accounting framework that addresses this deficiency. In our framework, each connection (both physical and informational) can be unambiguously assigned, quantified, and qualified, and an input-output balance is easily checked and maintained for each product. Costly independent data collections can be integrated into this common framework to amplify their original usefulness and provide the investigator or ecosystem manager with enhanced understanding of the entire ecosystem from which they were taken. The integrated data also allow various ecosystem models to be constructed efficiently, without unnecessary and costly duplication of effort. We present detailed guidelines for construction of such a framework, followed by examples and applications. © 1991 Academic Press, Inc.

Contents

- I. *Introduction.* A. What is an accounting system? B. Why use an accounting system?
- II. *Development of issues for accounting systems in ecology.*
- III. *A general accounting framework.* A. Desiderata. B. The art of choosing stocks and flows. C. Quantifying the flows. D. Examples. E. Data quality control.
- IV. *Uses of the general accounting system.* A. Network analysis and energy intensities. B. Simulation modeling. C. Optimization.
- V. *Summary and Conclusions.*

I. INTRODUCTION

A. *What Is an Accounting System?*

In biology, species are identified and classified according to a general system first devised by Linnaeus. The assignment of any particular species to a place in the hierarchic system is based on genetic and morphological similarities between organisms. Classifications are often debated and reassignments are occasionally made, but progress in biology without such a system would not be possible.

In ecology, although functional connections between the many species and their environment have been observed for decades, there is no accepted general accounting system for noting and comparing these connections. We propose a functional taxonomy for quantitative ecology—a general ecological accounting system. It is a framework in which the quantified connections between organisms (individual species, collections of species) and their abiotic environment can be placed and balanced, without ambiguity, omission, or double counting exchanges, at any scale which an investigator chooses. By connections we mean any kind of exchange of product (e.g., nectar from a plant, pollination time from an insect) between ecological processes (e.g., insect and plant). A connectivist ecology is one in which the choice of process definition is largely discretionary, but once the choice is made, the interconnection flows are determined and are of primary importance in the ensuring system description. This is in contrast to the Linnaean taxonomy, in which the place of an organism in the system is determined by its structure and evolutionary history, and flows between it and the rest of the system are of secondary importance at best.

This general ecological accounting system is the product of years of debate among ecological modelers, and it also benefits from years of debate and experience gained in national economic accounting.

B. *Why Use an Accounting System?*

The principal advantage of a universal accounting system in ecology is that it allows the material, energy, and service flows between all the parts of an ecosystem to be systematically placed in a common framework. Ecologists have long been involved in material and energy flow accounting (Hannon, 1973; Finn, 1976), but in the past each type of material (e.g., nitrogen) or energy was accounted for independently of all the rest. To qualify as ecosystem accounting, the interconnections between all the material and energy (and service) flows must also be included. This is the purpose of the framework proposed below.

The concept of a common framework also appreciates the value of

extant data bases (gathered for and incorporated and commensurate with current data. Together, old and new provides the format for evaluation (Hannon and Joiris, 1989).

In more detail, the use of a system before assure ecosystem analysts that a particular ecosystem was defined by a boundary (in space and time) and service flows for each component achieved; 3. with quantified connections particular species of special interest; 4. of data from various researches of growing utility.

The first known accounting system developed in the 18th century was a system accounting framework where flows were measured was in the field of concept of net output of an economy of quantitatively comparing the flows between various products. Leontieff (1941) developed the concept of total output required to produce the whole economy. This problem was of great importance during World War II. Many countries used a system to look for production of their economies in specific approaches to allow unambiguous comparison. With Stone's contribution, the system was essentially complete. The use of this system (Koopmans, 1951) and their development (1979) have been reported.

We have been guided in our development of the system by the experience of the laws of thermodynamics. With this recognition, we have attempted to create a more common framework that incorporates thermodynamics and meets the needs of ecologists to integrate economic and management of large regions as so elegantly recommended

ing System?

classified according to a general
 nment of any particular species
 d on genetic and morphological
 ations are often debated and
 rogress in biology without such

ions between the many species
 For decades, there is no accepted
 comparing these connections. We
 tive ecology—a general ecologi-
 n which the quantified connec-
 es, collections of species) and
 d balanced, without ambiguity,
 any scale which an investigator
 d of exchange of product (e.g.,
 a, an insect) between ecological
 vist ecology is one in which the
 tionary, but once the choice is
 ined and are of primary impor-
 a. This is in contrast to the
 F an organism in the system is
 y history, and flows between it
 importance at best.

em is the product of years of
 so benefits from years of debate
 c accounting.

ing System?

ccounting system in ecology is
 vice flows between all the parts
 ced in a common framework.
 rial and energy flow accounting
 ast each type of material (e.g.,
 ependently of all the rest. To
 erconnections between all the
 st also be included. This is the

also appreciates the value of

extant data bases (gathered for unrelated purposes) in that they can be incorporated and commensurately scaled into the same framework as current data. Together, old and new data can reside in a framework that provides the format for evaluation of whole ecosystem function (e.g., Hannon and Joiris, 1989).

In more detail, the use of an ecological accounting system could therefore assure ecosystem analysts and managers that a research effort on a particular ecosystem was done: 1. with full awareness of the system boundary (in space and time); 2. such that balance of materials, energy, and service flows for each compartment of the ecosystem under study was achieved; 3. with quantified connection to the rest of that ecosystem of any particular species of special interest; 4. in a way that allowed the compilation of data from various researchers of differing interests into a framework of growing utility.

The first known accounting framework for physical systems was developed in the 18th century by Turgot and Quesnay. The first large system accounting framework in which quantified interconnecting flows were measured was in the field of economics. Kuznets (1946) defined the concept of net output of an economy (Gross National Product) as a way of quantitatively comparing the activity levels of the economies of various countries. Leontieff (1941) defined a matrix structure for collecting the flows between various producing sectors so that he could determine the total output required to produce a unit of net output of each sector of the whole economy. This procedure is of crucial importance in planned economies and was of great use to the U.S. government just prior to and during World War II. Many countries (especially Japan) use the Leontieff system to look for production "bottlenecks" and to plan the development of their economies in specified areas. Stone (1963) refined the Leontieff approach to allow unambiguous accounting of multiple inputs and outputs. With Stone's contribution, the *economic* accounting procedure concept was essentially complete. The use of such accounting systems for public policy (Koopmans, 1951) and their theoretical implications for ecology (Amir, 1979) have been reported.

We have been guided in our development of an *ecological* accounting system by the experience of these economists. However, the consequences of the laws of thermodynamics are more apparent in ecological systems. With this recognition, we have modified the economic approach in an attempt to create a more comprehensive, less arbitrary accounting system that incorporates thermodynamic constraints and that simultaneously meets the needs of ecologists and economists. An added benefit is the ability to integrate economic and ecological data for analysis, planning, and management of large regional systems and for global change studies, as so elegantly recommended by Roughgarden (1989).

II. DEVELOPMENT OF ISSUES INVOLVING ACCOUNTING SYSTEMS IN ECOLOGY

Quantifying a population's food sources and losses in terms of biomass and numbers can lead to useful insights. However, when the interaction of living organisms with their non-living surroundings are measured, it is only natural to seek a physical yardstick. The fact that a significant portion of an organisms' surroundings consists of other living populations leads us to seek consistency by applying the same physical yardstick to interbiotic transfers.

Lindeman (1942) first constructed a budget of the flows connecting the biotic and non-living components of a senescent Michigan lake ecosystem. Lindeman's work is considered by many to be the prototype of modern ecosystems accounting schemes. It emphasized the relational or connectivist view of ecology. Not only is a population related to its physical surroundings, but it is connected *in the same terms* to other populations. Through Lindeman, ecology experienced a hierarchical leap from the study of populations to a consideration of the wider scope and scale of ecosystems.

Also important were the units in which Lindeman's accounting ultimately were reported—those of energy. Energy is a physical characteristic and can be used to express the physical nature of a biological system in a way that is *non-reductionistic*. Furthermore, energy is a construct that derives from the phenomenological science of thermodynamics. By accounting for all energy entering and leaving each population, one is recapitulating the first law of thermodynamics—that energy is neither created nor destroyed. But the first law begs the invocation of the second, which says (in one of its myriad of equivalent forms) that useful output from any process cannot occur without the degradation of some of the process input into less useful forms. The inviolability of the second law impresses a pyramidal form upon any discernible trophic chain of energetic transfers. The amounts of energy available to be passed on to higher trophic links must become progressively smaller.

Eugene Odum (1953, 1959) recognized the hierarchical shift implied by Lindeman's budgets and did more than anyone else to popularize the ecosystems perspective in quantitative terms. Howard Odum also pressed for further applications of the second law in ecology (Odum and Pinkerton, 1955). Howard Odum was also intrigued by the cycles that appear in most community budgets. He argued (Odum, 1971) that such cycles are indicative of the cybernetic processes at work in structuring ecosystems. Those components and pathways that are more effective than others at recycling materials and energy unto themselves will eventually grow at the relative expense of those other members. Systems develop and evolve in the same sense as do their components.

The shift towards whole systems insights. Connecting elements possible to trace indirect causal number and complexity of such perceptual capacities. So to make one must resort to systematic analysis of indirect effects is hardly linear economics as well. It was an ecology fully applied the techniques of indirect effects. Some economists cal flows be included in the Le

Hannon (1973) demonstrated to ecological budgets. There for put" analysis to various ecosystems to address better various issues combined input-output analysis notion of an ecological niche to (1978) emphasized the probability spotlighted the Markovian assumption Finn (1976) studied the indirect upon themselves and showed system activity that is devoted

Amir (1979) laid out the ecological systems, including the interpretation and Kemp (1979) remarked how malized transfers (the technical tion on how much medium tra They used the successive mat

trophic interactions into Linde Matis and Patten (1981) stocks, associated with the system effects and sketched out what standard input-output analysis component, environ analysis change via an assumed set of the stability of such linear systems (either endogenous or external towards particular configurations

Virtually all of the early work exchange of a single medium. T or carbon, or nitrogen; but se wherein more than one medium involved no physical exchange

COUNTING SYSTEMS IN ECOLOGY

and losses in terms of biomass. However, when the interaction of fundings are measured, it is only fact that a significant portion of er living populations leads us to physical yardstick to interbiotic

get of the flows connecting the scent Michigan lake ecosystem. o be the prototype of modern sized the relational or connec- pulation related to its physical me terms to other populations. hierarchical leap from the study ne wider scope and scale of

which Lindeman's accounting. Energy is a physical charac- physical nature of a biological. Furthermore, energy is a con- cal science of thermodynamics. leaving each population, one is amics—that energy is neither s the invocation of the second, alent forms) that useful output ne degradation of some of the nviolability of the second law nible trophic chain of energetic le to be passed on to higher aller.

he hierarchical shift implied by anyone else to popularize the s. Howard Odum also pressed law in ecology (Odum and o intrigued by the cycles that argued (Odum, 1971) that such cesses at work in structuring ys that are more effective than nto themselves will eventually members. Systems develop and nents.

The shift towards whole system budgets afforded other connectivist insights. Connecting elements together in an overall budget made it possible to trace indirect causes in quantitative terms. However, the number and complexity of such indirect linkages easily overwhelms one's perceptual capacities. So to make any progress in evaluating indirect effects one must resort to systematic analytical techniques. Of course, the question of indirect effects is hardly limited to ecology. It is a significant issue in economics as well. It was an economist (Leontieff, 1941) who first successfully applied the techniques of matrix algebra to the task of accounting for indirect effects. Some economists (Daly, 1968) have suggested that ecological flows be included in the Leontieff approach.

Hannon (1973) demonstrated how Leontieff's methods could be applied to ecological budgets. There followed copious efforts to apply "input-output" analysis to various ecosystems and to modify the analytical techniques to address better various issues of ecological concern. Patten et al. (1976) combined input-output analysis with general systems theory to extend the notion of an ecological niche to include indirect impacts and effects. Barber (1978) emphasized the probabilistic nature of ecological transfers and spotlighted the Markovian assumptions underlying the Leontieff approach. Finn (1976) studied the indirect effects that many system components exert upon themselves and showed how to estimate the fraction of the total system activity that is devoted to recycle.

Amir (1979) laid out the economist's views on equilibrium in ecological systems, including the interpretations of value, cost, and price. Ulanowicz and Kemp (1979) remarked how the algebraic powers of the matrix of normalized transfers (the technical coefficients of economics) provide information on how much medium traverses trophic pathways of various lengths. They used the successive matrix powers to transform arbitrary webs of trophic interactions into Lindeman-like chains or pyramids of flows.

Matis and Patten (1981) attempted to incorporate the contents, or stocks, associated with the system components into the analysis of indirect effects and sketched out what they called "environ analysis." Unlike the standard input-output analysis of systems that must balance around each component, environ analysis treats systems that are unbalanced and change via an assumed set of linear dynamics. Hannon (1986) examined the stability of such linear systems and evaluated several control strategies (either endogenous or externally applied) that could guide the system towards particular configurations.

Virtually all of the early work in ecosystem accounting focused on the exchange of a single medium. That is, budgets were cast in terms of energy, or carbon, or nitrogen; but seldom was there an attempt to treat systems wherein more than one medium was flowing, or to include connections that involved no physical exchange (e.g., the effect of pheromones, shading,

pollination). Implicit in single medium analyses are the assumptions that all inputs to a compartment have equal effect upon the recipient population, and that only one product issues from each compartment (usually its biomass as food to predators.) The latter assumption is not valid because organisms produce by-products such as feces, urine, detritus, and other exudates. How does one treat these multiple products that issue from a population or process? But perhaps even more problematic is the likelihood that the various foods consumed by each predator differ in their utilities to the consumer. How do we weight the various inputs to a compartment to reflect their relative utilities? Fortunately, one solution to both these difficulties is suggested by recent work in input-output theory.

In economics, one view of the origin of prices is that they account for the cumulative value added to a particular item at every step in the economic process from raw materials to finished product.¹ In ecology it is sometimes possible to make analogous calculations (Costanza and Neill, 1984). If a system has only one external input of a type not produced in the system, it is then possible to trace back through the network to identify the amounts of that input medium that were necessary to create any given connection. That is, one can estimate the portion of the input that has been "embodied" into each connection. It has been shown (Costanza, 1980; Costanza and Herendeen, 1984) how all the energies contributing to various economic products are correlated to their market prices. In the special case of a network with but a single medium (e.g., all energy flows), a particular transfer between higher trophic levels may yield value when directly measured (a certain number of calories). However, the *de facto* value of the flow in a systems context would reflect more the amount of the external input (e.g., sunlight) that went into creating the given flow. The ratio of the latter quantity to the former is called an "intensity" and can be used to weight the various inputs to a population so as to allow a legitimate comparison among these inputs.

Similarly, intensities can be calculated for an ecosystem description which has multiple outputs as well as multiple inputs. That procedure is more complex (Costanza and Hannon, 1989) and requires the accounting system described below.

Amir (1987) has discussed the use of an ecological accounting system, both static and dynamic, to elaborate on the formal connections between ecology, thermodynamics, and economics. He points out that the "intensities" of which we speak are measured in terms of an external input and that we therefore must choose which external input should form the basis of the intensities. In the development of the general accounting system,

¹ This is the view we take in our paper. Amir (1979) points out the other economic theories on the origin of prices.

however, we are not proposing a model. We view the accounting as a kind of ecosystem model which can be used for a comparison of model results. The accounting system which we propose is reminiscent of economic price allocation problems, just as the

III. A GENERAL

We can now set forth, in general terms, an accounting framework for ecological and economic systems. These can be characterized by (1) comprehensiveness; (2) uniqueness; (3) uniqueness; and (4) uniqueness, in more detail below.

Generality

The accounting framework is characterized by generality, in the sense that it applies to any particular system or temporal scale, both ecological and economic, ranging from microcosms to the whole earth. It is characteristic so as to begin to model the interactions of systems and to start to bridge the gap between ecology and economics. We wish to apply it to geographic regions of the same size over a given time period.

Comprehensiveness

The accounting framework is characterized by comprehensiveness, in the sense that it should include the full range of phenomena that are relevant to the practical limitations will often be imposed by the accounting, but the accounting system is not an impediment. The accounting system should have several different types of flows which have several different types of flows. Equally should be able to include non-physical exchange between a plant, it receives nectar (physical flow) from other flowers (non-physical flow) should be included in the accounting system.

analyses are the assumptions that affect upon the recipient population in each compartment (usually its assumption is not valid because feces, urine, detritus, and other simple products that issue from a given more problematic is the by each predator differ in their the various inputs to a com- fortunately, one solution to both k in input-output theory.

prices is that they account for the n at every step in the economic duct.¹ In ecology it is sometimes (Costanza and Neill, 1984). If a pe not produced in the system, h the network to identify the necessary to create any given con- tion of the input that has been been shown (Costanza, 1980; l the energies contributing to to their market prices. In the medium (e.g., all energy flows), ic levels may yield value when alories). However, the de facto l reflect more the amount of the to creating the given flow. The called an "intensity" and can be population so as to allow a

for an ecosystem description multiple inputs. That procedure is 9) and requires the accounting

a ecological accounting system, the formal connections between He points out that the "inten- terms of an external input and al input should form the basis he general accounting system,

points out the other economic theories

however, we are not proposing the use of intensities or any other kind of model. We view the accounting system as a necessary underpinning for any kind of ecosystem model which claims to be consistent with any other kind. The accounting system which we propose forms the basis for consistent comparison of model results. Amir points out that these intensities are reminiscent of economic prices which arise from ecological resource allocation problems, just as they do in economics.

III. A GENERAL ACCOUNTING FRAMEWORK

A. *Desiderata*

We can now set forth, in general terms, the minimum desired criteria for an accounting framework capable of handling both ecological and economic systems. These can be summarized as (1) generality; (2) comprehensiveness; (3) uniqueness; and (4) quality control. These are described in more detail below.

Generality

The accounting framework should exhibit a high degree of universality or generality, in the sense that its usefulness should not be limited to any particular system or temporal or spatial scale. It should be applicable to both ecological and economic systems, on spatial and temporal scales ranging from microcosms to the planet as a whole. We desire this characteristic so as to begin to make valid comparisons between divergent systems and to start to bridge the wide chasm that currently exists between ecology and economics. We wish to be able to combine data from different geographic regions of the same ecosystem into the same framework, for a given time period.

Comprehensiveness

The accounting framework should exhibit a high degree of comprehensiveness, in the sense that it should, at least in theory, be able to include the full range of phenomena that occur in a system. We understand that practical limitations will often limit the achievable comprehensiveness of the accounting, but the accounting framework itself should not be the impediment. The accounting system should be able to handle processes which have several different types of inputs and outputs. The system equally should be able to include simultaneously a physical exchange and a non-physical exchange between processes. For example, when a bee visits a plant, it receives nectar (physical exchange) and provides cross pollination from other flowers (non-physical exchange). Both types of exchanges should be included in the accounting system.

Uniqueness

The accounting framework should provide a unique and unambiguous place for each measurable type of connection. The physical or thermodynamic quality of the connection must be recognized and treated consistently and the system boundaries should be physically definable in a discoverable way.

Quality control

The accounting framework should exhibit an ability to deal with information of varying quality in a simple and consistent way. The precision of the data for each connection should carry its own mark of reliability and the change in that mark should be specified as the data are mingled arithmetically with other data.

B. The Art of Choosing Stocks and Flows

When asked to list the important components of the system, it is extremely rare that two investigators will spontaneously enumerate the same elements. Of course, the subjective biases that determine what is actually measured will color the ensuing analysis. Critics of systems ecology are always quick to cite the relative nature of the results from network analyses. It is our opinion that in the end the ambiguities associated with which components to include in the accounting scheme (the lexical decisions) pale in comparison to the insights into system structure and function one obtains by performing the budgetary exercise. To be more specific, we *hypothesize* that the qualitative conclusions drawn when the accounting techniques are applied to ecosystems will differ very little (if at all) whenever the same ecosystem is parsed in different ways by various investigators. More important is the application of a consistent, systematic, and comprehensive framework of accounting for exchanges and storages, much as the one described below. We can also put this question to the test by reparsing and reanalyzing the ecological system if the accounting framework allows this kind of manipulation to be done conveniently. Of course, the different scales used by independent investigators must be considered when the data is transposed into a common framework.

Such apology having been made, it is nonetheless useful to elaborate the *art* of choosing the elements to include in the accounting. As in general systems theory, we feel the delimitation of the system is strictly at the discretion of the observer; i.e., the system boundaries and list of internal elements may be chosen at will. That is not to say that all choices are equally good.

Common sense should prevail in defining the boundaries of the system. Ideally, the physical boundaries are chosen to minimize the amount and

diversity of exchanges across more easily monitored. Interfaces such as air-water or water-lake. Topographic features, such as nearly as well. The duration and hierarchical boundaries on the and restrictive as the demarcation. Temporal limits devolves into more a full discussion of these issues. Suffice it to say that the duration enough to characterize statistically frequent sampling will be required changes.

The temporal limits are the compartments. What are the ecosystem? The answer, of course investigator or the source. Certainly, the experienced investigator abundant species in the system. tion of budget items. Microbiological stocks, but engender a disproportionate a species of marginal ecology. accounting by virtue of its ecological usefulness, or, in the case of energy discussed below, value is not a

Then there is the delicate seemingly insignificant species ecosystem processes. Control beyond the scope of this paper that is known to exert strong an ecologist is always at risk. observer, is exhibiting control be as tragic as it seems. For included in the accounting process that the influences of these un processes.

While the finitude of resources compartments that can be in historical data conversely may budget that otherwise would be the duration of the project. In by the quantity of data extant all that is necessary to piece

vide a unique and unambiguous
ction. The physical or thermo-
be recognized and treated con-
ld be physically definable in a

it an ability to deal with infor-
consistent way. The precision of
its own mark of reliability and
ified as the data are mingled

Stocks and Flows

omponents of the system, it is
l spontaneously enumerate the
biases that determine what is
alysis. Critics of systems ecology
re of the results from network
the ambiguities associated with
unting scheme (the lexical deci-
o system structure and function
exercise. To be more specific, we
as drawn when the accounting
ll differ very little (if at all)
in different ways by various
tion of a consistent, systematic,
ng for exchanges and storages,
also put this question to the test
ical system if the accounting
on to be done conveniently. Of
pendent investigators must be
o a common framework.

hetheless useful to elaborate the
the accounting. As in general
of the system is strictly at the
boundaries and list of internal
not to say that all choices are

g the boundaries of the system.
a to minimize the amount and

diversity of exchanges across them. Then the exogenous transfers can be more easily monitored. Interfaces that inhibit certain classes of transfers, such as air-water or water-land, are often convenient for this purpose. Topographic features, such as watershed or airshed boundaries, often work nearly as well. The duration and frequency of sampling for data also place hierarchical boundaries on the system description every bit as important and restrictive as the demarcation of spatial domain. The definition of temporal limits devolves into monitoring strategy, and space does not permit a full discussion of these issues (see, for example, Schubel et al., in press). Suffice it to say that the duration of data acquisition should be long enough to characterize statistically the slower biotic processes and the most frequent sampling will be required by those populations exhibiting rapid changes.

The temporal limits are thus inexorably connected with the choice of compartments. What are the important processes and products of the ecosystem? The answer, of course, will be colored by the inclinations of the investigator or the source of funding for the accounting endeavor. Certainly, the experienced investigator will rarely omit the visibly most abundant species in the system. Activity level is another criterion for selection of budget items. Microbial populations usually have small standing stocks, but engender a disproportionate fraction of system activity. Often a species of marginal ecological importance will be included into the accounting by virtue of its economic, aesthetic, recreational, or political usefulness, or, in the case of endangered species, its rarity per se. As will be discussed below, value is not always synonymous with magnitude.

Then there is the delicate issue of control. It may happen that a seemingly insignificant species is exerting a powerful control upon ecosystem processes. Control is a dynamical attribute, and as such is beyond the scope of this paper. But we would be foolish to exclude an item that is known to exert strong control on significant processes. Conversely, an ecologist is always at risk of excluding a species that, unknown to the observer, is exhibiting control upon the system. Such negligence may not be as tragic as it seems. For practical reasons not every agency can be included in the accounting process, and the investigator is forced to assume that the influences of these unknown dynamics are implicit in the chosen processes.

While the finitude of resources for research may abridge the number of compartments that can be included in the scheme, the availability of historical data conversely may allow the inclusion of some species into the budget that otherwise would be too costly or impossible to monitor under the duration of the project. In fact, the reader would probably be surprised by the quantity of data extant on his/her ecosystem of interest. Quite often, all that is necessary to piece together a system budget of surprisingly

good resolution is a couple of man-years of expert effort and persistence (cf. Costanza et al., 1983; Baird and Ulanowicz, 1989).

Finally, there is no reason to regard the lexical choices an investigator makes at the outset of an accounting project as cast in stone. Accounting projects allow for iterative reevaluation as more data accrue. Often the investigator may begin with order-of-magnitude guesses as to the intensity of various processes and use these estimates to gauge the values to be attached to each flow. As a result, certain processes might be dropped from further consideration or aggregated with more important elements.

C. Quantifying the Flows

Once the system boundary in space and time has been identified, and the processes with their input and output products have been chosen, one can begin the accounting procedure. There are six distinct types of connections, in addition to the stocks, that must be identified in analyzing any ecosystem: 1. the non-produced inputs; 2. the net outputs; 3. the product use record; 4. the product production record; 5. the total output; and 6. the waste heat flows. These product flows are shown in Fig. 1.

1. The Non-produced Inputs

We distinguish here between imported/exported products made within the defined ecosystem, and imports of special products which are not made within this system. One thinks of the import of the non-produced products as a constraint on the system's activity level. These are the products which are sometimes deemed as scarce or growth-limiting in some sense. Sunlight is a non-produced import but not the only possible one. For example,




FIG. 1. The product flows into and out of a typical process in an ecosystem. Product flows are mainly distinguished as going to other processes inside the system boundary or, crossing the boundary. Products imported from outside the system are separated depending on whether they are made in the system or not.

sunlight contains blue, red, and violet, and is used by the ecosystem. While violet is extraneous, it is possible to detect its presence in the ecosystem. If the ecologist chooses to account for radiation, radiation should be included as an input.

Ambient heating is also an input. Heat flows through the ecosystem. Ambient heating, from an economic point of view, is a departure from some ideal state. It is a factor in agricultural output and an input to the ecosystem. From a point of view, the heat flow is a primary controlling factor in many processes. It is largely cyclical and within a system. To a large extent, heat flows are not of primary importance. They are of course important in determining the development in the ecosystem. They are as a measured net inputs to the system.

2. The Net Outputs

The net output flows are computed by crossing the system boundary to itself and to the outside. There are three kinds of net outputs: 1. the net output of our accounting system.

First, we have the import and export of products. The processes in the defined system are not the same. One of the processes is likely to be a process that would be "algal biomass". Some of the products are system (export) or into the system.

The second kind of net output is the net output of a particular product which occurs within the system. We have assumed a static picture of the system. Changes in stocks explicitly as a flow across the time boundary. An increase in storage will be used to account for the change.


Thus, the requirements are for a system that is harder to describe. A change is a flow across the time boundary. A change of stock due to the natural processes is inherent in all products of the system. Thermodynamics? Everyone is a part of the system. These same processes are countered by the simultaneous processes.

of expert effort and persistence (Ulanowicz, 1989). The lexical choices an investigator select as cast in stone. Accounting as more data accrue. Often the magnitude guesses as to the intensity rates to gauge the values to be processes might be dropped from more important elements.

Flows

time has been identified, and the products have been chosen, one can six distinct types of connections, be identified in analyzing any the net outputs; 3. the product record; 5. the total output; and vs are shown in Fig. 1.

exported products made within al products which are not made rt of the non-produced products l. These are the products which limiting in some sense. Sunlight nly possible one. For example,



process in an ecosystem. Product flows inside the system boundary or, crossing the system are separated depending on

sunlight contains blue, red, and infrared radiation, all of which may be used by the ecosystem. While we sometimes think of infrared radiation as extraneous, it is possible to define this radiation as a product *within* the ecosystem. If the ecologist chooses not to define it in this way, then infrared radiation should be included as a non-produced input to the system.

Ambient heating is also an input to any system. In what sense is the flow of heat through the ecosystem an important theoretical consideration? Ambient heating, from an economic point of view, is very important. The departure from some ideal standard results in a drop in the value of agricultural output and an increase in fossil fuel use. From the ecologic point of view, the heat flow is also very important, since temperature is a primary controlling factor in many life processes. However, heat flows are largely cyclical and within a given range, they are predictable. To this extent, heat flows are not of primary importance in ecosystem studies. They are of course important in detailed studies of the variations in organism development in the ecosystem. We recommend that heat flows be included as a measured net inputs to the system.

2. The Net Outputs

The net output flows are connections from the defined ecosystem across the system boundary to itself at a future time and/or to other ecosystems. There are three kinds of net output flows which must be accommodated in our accounting system.

First, we have the import and export of products made and/or used by the processes in the defined system. For example, in an aquatic ecosystem, one of the processes is likely to be algae, and one of its output products would be "algal biomass". Some of this biomass could be washed from the system (export) or into the system (import) during a flood.

The second kind of net output flow is the change in the storage level of a particular product which occurs during the designated time period. Since we have assumed a static picture of the ecosystem, we must include the changes in stocks explicitly as flows. These flows might be thought of as flows across the time boundary of the chosen period. For example, an increase in storage will be used in the next or later time periods.

Thus, the requirements are not very surprising for the experienced ecologist. But there remains a third net output flow across the ecosystem boundary that is harder to describe. If, as we have just assumed, the stock change is a flow across the time boundary of the system, what about the change of stock due to the natural decay and stock replacement processes inherent in all products of the ecosystem as required by the second law of thermodynamics? Everyone is aware of the decay processes associated with death. These same processes are at work in the living system, but they are countered by the simultaneous restorative or anabolic processes we call life.

Thus, when one observes a quantity of living substance perceived to be of constant stock over the chosen time period, one is really seeing the balanced result of the decay and replacement processes. During the study time period, a certain amount of the living product actually decayed into its component parts and some of that lost stock was replaced. This decay-replacement process is known to occur because the products of decomposition and the net absorption of energy are both evident. If the stock size of the organism does not change during the chosen time period, what is lost or gained by the organism? It is known from thermodynamics that the *order* of the structure of the organism was lost for that stock of product which decayed during the time period. Yet, for the case where the total product stock size did not change, that lost order was exactly made up by what we call the replacement process. One must record the contribution of this replacement process in some manner. It is a product contribution across the time boundary of the system in the same manner as is the net change of a product stock.

Decay rates can be measured directly by isolating a sample of the product in question and observing the loss of order with time. Frequently this isolation is not possible or practical, however, and an estimate of the magnitude of the decay rate for living products (e.g., biomass) must be substituted. The basal metabolism of living products can be used as an estimate of decay rates, since it represents the replacement flow required just to compensate for decay processes when the organism is at rest.

In a similar way, abiotic products in the ecosystem (e.g., solution ammonia) might also degrade during the time period, and in the process, give off heat. We need to know what the normal degradation rate would have been for this substance at the system temperature. Such data for most substances can be found in standard chemical reference handbooks. We accept these standard data as the *replacement* rate of the non-living substances in question. If one wishes to account for the waste heat associated with the decomposition of these abiotic compounds, one could add another process to the accounting system which describes the input and output of the compound and its degradation (therefore its replacement rate) and record the associated waste heat. Of course, some of the non-living substances might already be at their lowest energy state, at equilibrium with the system environment (e.g., carbon dioxide), and no decay need be considered.

The assignment of the basal or standard metabolism to the net output of the framework to represent the replacement of the depreciated stock is explicit acknowledgement of the second law of thermodynamics. The remainder of the respiratory flows are not counted as a part of the net output since they are not of the same quality as the flows along that row of the matrix. These flows are recorded only to provide a check on the

balance of the total flow of the process does not double count.

3. The Product Use Record

We have chosen the "Use" matrix and modified it to meet the special needs. The "Use" matrix is the record of the "Make" matrix is the product of the next section.

The first accounting step is to identify the process by which process. A "Use" or "U" matrix is placed at the head of a process and placed at the beginning of a row of a product (product "c") by a particular process (process "c") period (i.e., the flow of "c" into "c") lies at the intersection of row "c" and column "c". The substances consumed by predator are recorded. The picture of the U matrix appears as follows:

The units of measure of product "c" is used, but these need not be the same as for other products. For example, product "c" is measured per square meter and product "d" is measured per unit of time. So long as the units are consistent, the products can be measured in any units.

It is clear to ecologists that the products are measured in type and quantity with time, and with the availability of resources.

	Process a	Process b	Process c
Product a			
Product b			
Product c			
Product d			

The "Use"

FIG. 2. The uses of products by the U matrix.

iving substance perceived to be period, one is really seeing the ent processes. During the study g product actually decayed into lost stock was replaced. This occur because the products of energy are both evident. If the e during the chosen time period, is known from thermodynamics anism was lost for that stock of period. Yet, for the case where the hat lost order was exactly made s. One must record the contribu- manner. It is a product contribu- em in the same manner as is the

by isolating a sample of the s of order with time. Frequently however, and an estimate of the products (e.g., biomass) must be ng products can be used as an s the replacement flow required en the organism is at rest.

n the ecosystem (e.g., solution time period, and in the process, normal degradation rate would temperature. Such data for most mical reference handbooks. We ment rate of the non-living sub- unt for the waste heat associated mpounds, one could add another escribes the input and output of efore its replacement rate) and ource, some of the non-living est energy state, at equilibrium dioxide), and no decay need be

metabolism to the net output of ent of the depreciated stock is law of thermodynamics. The counted as a part of the net out- y as the flows along that row of uly to provide a check on the

balance of the total flow of that substance in the system. The assignment process does not double count any of the flows.

3. The Product Use Record

We have chosen the "Use" and "Make" matrices approach (Stone, 1963) and modified it to meet the special problems involved with ecological flows. The "Use" matrix is the record of which process uses what product. The "Make" matrix is the product production record and is discussed in the next section.

The first accounting step is to record where each product is used and by which process. A "Use" or U matrix is constructed as follows: Each process is placed at the head of a column in the U matrix; each product is placed at the beginning of a row in this matrix. The use of a product (e.g., product "c") by a particular process (e.g., process "d") for the chosen time period (i.e., the flow of "c" into "d"), is the number placed on the cell which lies at the intersection of row "c" and column "d". The quantities of the substances consumed by predators for example, appear in the U matrix. The picture of the U matrix appears in Fig. 2.

The units of measure of product *c* must be the same wherever product *c* is used, but these need not be the same as the measure for any of the other products. For example, product *c* may be measured in grams-carbon per square meter and product *d* may be measured in kcal per square meter. So long as the units are consistent along the row of *U*, the products may be measured in any units.

It is clear to ecologists that the inputs to a particular species may vary in type and quantity with time, with age of the individuals in that species, and with the availability of resources. It is theoretically possible to include




FIG. 2. The uses of products by the various processes are tabulated in the Use or U matrix.

such variation in a single descriptive Use matrix by devising a column for each time, age, and resource condition. This matrix is then controlled by a time-varying vector of factors, each between zero and one, which select the proper portions of each of the columns involved with that species. The Make matrix (discussed below) may require a similar modification.

Besides allowing greater convenience in data gathering, the tolerance of the U matrix for any kind of unit of measure lets the experimentalists incorporate units of "service" into the matrix. For example, the pollination of flowers by bees may be measured in "pollination bee-seconds" per square meter for the chosen time period. We know that if some pollutant reduced the bee population by half, flower productivity would decrease. But there is hardly any physical transfer from bees to flowers, except that measurable by the service units of pollination. The "bee-seconds" measure is the collective visitation time spent by bees in flowers. In the U matrix, the process *plant* would have "used" so many bee-seconds of pollination service. So the term "product" as we use it here may mean an actual physical product *or* a service. Negative services, such as the effect of allelopathic chemicals, can also be included.

4. Product Production Record

In the general accounting system, a second matrix is needed to show where the products are made. Such a matrix is called a "Make" or V matrix and it has the same configuration as the U matrix: the column heads are the processes and the product names are shown at the beginning of each row. The quantified list of substances produced by prey for example, are listed in the production matrix. The elements of the V matrix are for example, the amount of product *b* which is produced or made by process *d*; see Fig. 3.




FIG. 3. The products made by the various processes are tabulated according to where they are made.

The units of a product must be the same as the units of the U matrix but can vary from p

The products of a species i are compared with the available portions of i for each of the year, energy, water, and other resources. An organism requires these inputs to produce biomass and the type and amount of vectors can be assigned to a species. A varying control vector, whose components are inputs, can be used to the app

5. The Total Output Flow

The sum of the entries in a row of the net outputs is the total output, which must equal to the sum of the corresponding entries in the balance here provides a check

6. The Waste Heat Flow

The respiratory or total was heat from the process and the seeking or predator-avoidance the accounting procedure, we process into a vector. We consider lack of utility to any of the other heat were used by certain components of heat would be classed as appropriate components in the heat could be used as a measure that quantity of heat would appear made by the same component in this manner. Note that the not associated with the basal representing basal metabolism processes and appear in the ne

To complete the accounting with the matrices of product inputs, the non-produced inputs to the system are best assembled at the lower end and export products, the channels of production. If all products are all products of the system, we therefore associate the matrix of inputs with the U (Use) matrix for convenience in balancing. The way

matrix by devising a column for
This matrix is then controlled by
between zero and one, which select
s involved with that species. The
ire a similar modification.
n data gathering, the tolerance
measure lets the experimentalists
rix. For example, the pollination
n "pollination bee-seconds" per
We know that if some pollutant
ver productivity would decrease.
from bees to flowers, except that
tion. The "bee-seconds" measure
bees in flowers. In the U matrix,
many bee-seconds of pollination
here may mean an actual physi-
such as the effect of allelopathic

second matrix is needed to show
matrix is called a "Make" or V
n as the U matrix: the column
ames are shown at the beginning
stances produced by prey for
ix. The elements of the V matrix
b which is produced or made by

The amount of Product b
made by Process d
in the chosen time period.

cesses are tabulated according to where

The units of a product must be consistent across the row of the V and U matrix but can vary from product to product.

The products of a species or an organism in the ecosystem can vary with the available portions of the net input. For example, at various times of the year, energy, water, and nitrogen may be in short supply. Since the organism requires these inputs in fixed proportion for growth, the rate of biomass and the type and amount of excretion products will vary. A variety of vectors can be assigned to a particular organism or species and a time-varying control vector, whose zero to one factors depend on the mix of net inputs, can be used to the appropriate mix of outputs through time.

5. The Total Output Flow

The sum of the entries in a row of U plus the corresponding row of the net outputs is the total output flow for that product. This sum is also equal to the sum of the corresponding row of the V or "Make" matrix. The balance here provides a check on the various product entries.

6. The Waste Heat Flow

The respiratory or total waste heat is composed of the basal metabolic heat from the process and the organism's muscle and bone friction, prey-seeking or predator-avoidance effort, and reproductive effort. To complete the accounting procedure, we collect all of the heat given off by each process into a vector. We consider this heat as lost to the system due to its lack of utility to any of the other organisms in the system. If some of this heat were used by certain components in the ecosystem, then that quantity of heat would be classed as a product produced and used by the appropriate components in the U and V matrices. If some of this "waste" heat could be used as a measure of reproductive service for example, then that quantity of heat would appear in the U and V matrices as used and made by the same component. All of the "waste" heat could be assigned in this manner. Note that the waste heat flow defined in this section is not associated with the basal metabolic processes: those substances representing basal metabolism have been assigned to represent the decay processes and appear in the net output.

To complete the accounting framework, we augment Figs. 2 and 3 with the matrices of product inputs and outputs to the system. Since the non-produced inputs to the system are products used by the system, they are best assembled at the lower edge of the U matrix. The produced import and export products, the changes in product stocks, and the depreciation products are all products of the system but they cross the system boundary. We therefore associate the matrix of these three types of flows (the net outputs matrix) with the U (Use) matrix on its right-hand side for convenience in balancing. The waste heat from the processes of the system is




FIG. 4. The general accounting system describing the assignment of all product flows to where they are made and used.

tabulated at the bottom edge of the V (Make) matrix as a notational reminder that they have been accounted for. The complete ecological accounting system therefore looks as shown in Fig. 4.

As a matter of accounting convenience, the U and V matrices are overlaid in tabular form and kept in the same table, with the Make or V matrix entries shown immediately below the Use or U matrix entries. Also note that the U and V matrices will most likely be rectangular and not square, because the number of products used will not necessarily equal the number of products made.

The general accounting procedure is shown in the following examples.

D. Examples


The following simplified examples will serve to illustrate the accounting system. We have chosen to use simplified examples rather than data from actual systems. The application of a simplified form of the accounting system to the Southern North Sea ecosystem has been published (Hannon and Joiris, 1989; Costanza and Hannon, 1989). Our goal in this paper is to demonstrate the full accounting system in the simplest possible form so that each of the principles can be clearly explained. We know of no ecosystem for which a sufficient amount of data has been collected and codified for application of the full accounting procedure.

The first example is a hypothetical ecosystem consisting of three internal processes (abiotic chemical processes, primary producer processes, and consumer processes) and five products (nutrients, plant biomass, animal biomass, waste heat, and sunlight). The "use" or inputs, "make" or out-

TABLE I
Hypothetical Ecosystem Network Illustrating the General Accounting Framework

Products (units)	Internal Processes			External Processes		Totals	
	Abiotic Chemical Processes	Biotic processes		Change in stock	Exports/ imports		Depre- ciation
		Producers	Consumers				
		Net Outputs					

product c
of product c
on of product c
al amount of product c used and made



the assignment of all product flows to

(Make) matrix as a notational
d for. The complete ecological
n in Fig. 4.

ce, the U and V matrices are
ame table, with the Make or V
e Use or U matrix entries. Also
t likely be rectangular and not
ed will not necessarily equal the

own in the following examples.

s

erve to illustrate the accounting
examples rather than data from
implified form of the accounting
m has been published (Hannon
.989). Our goal in this paper is
in the simplest possible form so
ly explained. We know of no
of data has been collected and
ng procedure.

system consisting of three internal
primary producer processes, and
utrients, plant biomass, animal
use" or inputs, "make" or out-

TABLE I
Hypothetical Ecosystem Network Illustrating the General Accounting Framework

Products (units)	Internal Processes				External Processes			Totals
	Abiotic Chemical Processes	Biotic processes			Change in stock	Net Outputs		
		Producers	Consumers			Exports/ imports	Depre- ciation	
Nutrients (g/t)	use	20	10	0	0	5	2	37
	make	20	5	5	—	7	—	37
	stock	500	0	0	—	—	—	500
Plant biomass (g/t)	use	1	5	6	1	0	1	14
	make	0	14	0	—	0	—	14
	stock	0	100	0	—	—	—	100
Animal biomass (g/t)	use	1	1	5	—	0	1	8
	make	0	0	8	—	0	—	8
	stock	0	0	10	—	—	—	10
Waste heat (Cal/t)	use	—	—	—	—	—	100	100
	make	5	50	45	—	—	—	100
Sunlight (Cal/t)	use	0	100	0	—	—	—	100
	make	—	—	—	—	100	—	100

Internally produced

Externally produced

puts, and "stock" of each of these products by (or in) each of the internal processes are shown in table I and Fig. 5. The table and figure are equivalent representations of the system. Figure 6 shows the nutrient product flows isolated from the system. Most network analysis in ecosystems has in the past concentrated on just such a one-product abstraction of the system. Our general accounting framework is intended to take more of the full complexity of ecological networks into account as shown in Fig. 5 and table I.




FIG. 5. Flows and stocks of five different products in a hypothetical ecosystem with three internal processes.





FIG. 6. Hypothetical ecosystem flow diagram.

Table I represents a particular abstraction of the complex flows shown in Fig. 5 that abiotic chemical products, primary producers use 10, and all three processes, since plants and decompose. In contrast, plant producers and consumers, respectively.

A fourth internal product is made by all the internal processes in addition to the four internal products of this ecosystem. This is sunlight.

by (or in) each of the internal
The table and figure are equivalent
e 6 shows the nutrient product
work analysis in ecosystems has in
product abstraction of the system.
ended to take more of the full
account as shown in Fig. 5 and



s in a hypothetical ecosystem with three




FIG. 6. Hypothetical ecosystem flows and stocks of the single product "nutrients".

Table I represents a particularly compact, yet easily readable representation of the complex flows shown in Fig. 5. For example, one can read from Fig. 5 that abiotic chemical processes use 20 g per unit time of nutrients, primary producers use 10, and consumers use 0. Nutrients are "made" by all three processes, since plants and animals release nutrients when they die and decompose. In contrast, plant and animal biomass are "made" only by producers and consumers, respectively.

A fourth internal product is waste heat. This product is unique in that it is made by all the internal processes but is used by none of them. In addition to the four internal products, one "external" product is noted for this ecosystem. This is sunlight, which is used by producers to drive the

system and is not made anywhere in the system. Any product that is not produced anywhere in the system under study but is necessary for the system's survival is called an externally produced product.

There are also three "external processes" shown, change in stock, exports/imports, and depreciation replacement, along with their use and make of the system's products. As already mentioned, a change in stock is equivalent to an export to another time period, and depreciation is equivalent to export from the system. So, taken together, these three external processes represent net outputs from the system of one kind or another.

Note that for this system (and for any system) mass and energy balance are maintained for each product. For example the total use of nutrients in the system of 37 g/t (use by internal processes plus use by external processes in the form of change in stock, exports and depreciation) is equal to the total production of nutrients by internal processes plus imports (also 37 g/t). Product use must balance product make for each product in the system.

"Products" in our accounting system can also be services or information. To illustrate this we present an example system in which bee pollination services are an important product (table II). In this example the products are bee biomass, honey, plant biomass, and been pollination effort, in addition to the ubiquitous waste heat and sunlight. The internal processes are bees, and two species of plants that the bees pollinate. Bees in this system produce four different products; (1) bee biomass (7 g/t); (2) honey (35 g/t); (3) pollination effort (10 bee s/t); and waste heat (5 Cal/t). Note that both bee biomass and honey are used and made exclusively by bees. Bee pollination effort is used more by plant 1 (6 bee s/t) than by plant 2 (3 bee s/t). Plant 1 also has twice the biomass stock of plant 2 (200 vs. 100) and uses more sunlight (40 vs. 30 Cal/t).

E. Data Quality Control

All numerical estimates have some uncertainty associated with them, and the normal assumption that they are measured with total precision is an inaccurate approximation. In the case of well-established experimental or field data, there are standard statistical techniques available for representing data quality and how varying quality of data affects results of accounting systems and models on them (cf. Bullard and Sebald, 1977). But statistically valid samples are usually available for only a very small percentage of the entries needed in the proposed accounting system. What we need is some other means to communicate and manipulate data quality when that quality falls below the level that standard statistics can deal with. We need a way to use all the scarce information available on systems connections in a clear and straightforward way. In the absence of such a

TABLE II
Hypothetical Ecosystem Network Illustrating the Accounting of Service Type Products

Products (units)	Internal processes			External processes			Totals
	Bees	Plant 1	Plant 2	Change in stock	Net outputs exports/ imports	Depre- ciation	
		Internally produced					
Bee biomass	5	0	0	0	1	0	1
use	7	0	0	0			7
make							

system. Any product that is not study but is necessary for the produced product.

uses" shown, change in stock, ment, along with their use and mentioned, a change in stock is period, and depreciation is equivalent together, these three external system of one kind or another.

system) mass and energy balance ample the total use of nutrients processes plus use by external exports and depreciation) is equal rnal processes plus imports (also t make for each product in the

also be services or information. system in which bee pollination t). In this example the products and been pollination effort, in sunlight. The internal processes the bees pollinate. Bees in this) bee biomass (7 g/t); (2) honey and waste heat (5 Cal/t). Note 1 and made exclusively by bees. nt 1 (6 bee s/t) than by plant 2 ass stock of plant 2 (200 vs. 100)

Control

certainty associated with them, measured with total precision is of well-established experimental stical techniques available for quality of data affects results of (cf. Bullard and Sebald, 1977). available for only a very small oposed accounting system. What ate and manipulate data quality standard statistics can deal with. ration available on systems conway. In the absence of such a

TABLE II
Hypothetical Ecosystem Network Illustrating the Accounting of Service Type Products

Products (units)	Internal processes			External processes			Totals	
	Bees	Plant 1	Plant 2	Change in stock	Net outputs			
					exports/ imports	Depre- ciation		
Bee biomass (g/t)	use	5	0	0	1	0	1	7
	make	7	0	0	—	0	—	7
	stock	35	0	0	—	—	—	35
Honey (g/t)	use	30	0	0	0	0	5	35
	make	35	0	0	—	0	—	35
	stock	0	100	0	0	0	0	100
Plant biomass (g/t)	use	5	25	35	3	0	5	73
	make	0	33	40	—	0	—	73
	stock	0	200	100	—	—	—	300
Bee pollination Effort (bee s/t)	use	0	6	3	—	0	1	10
	make	10	0	0	—	0	—	10
	stock	10	0	0	—	—	—	10
Waste heat (Cal/t)	use	—	—	—	—	—	70	70
	make	5	30	35	—	—	—	70
Sunlight (Cal/t)	use	0	40	30	—	—	—	70
	make	—	—	—	—	70	—	70

Internally produced

Externally produced

system, once a number is obtained and exported to the general scientific community it is labelled either "good" or "bad," and the record of its degree of quality is all but forgotten except by the experts familiar with the details of the measurement methods. What is needed is a formal, systematic way of communicating to the non-expert the "degree of goodness" or "quality" associated with each number. This can be imagined as composed of its degrees of precision and accuracy; that is, how little "spread" or scatter there is in its underlying data, and how reliable it is as an estimate of what it is measuring. Thus, we can think of all quantitative measurements as having associated truth-value modifiers, which we will call the "grades" of numbers. We chose this term since everyone is familiar with the grading process through their academic experience, and the process of grading quantitative measurements is conceptually similar to grading students or grading almost anything else. Also, stating a grade with each number connotes the evaluative and qualitative nature of the measure of uncertainty we propose, which is in keeping with its correct interpretation.

Such systems for grading and manipulating data of sub-statistical quality are currently under development (e.g., Funtowitz and Ravitz, 1989). We lack the space to adequately treat them here, except to say that they are a necessary adjunct to the proposed accounting system to allow information the *quality* of numerical estimates to be carried through the accounting and subsequent manipulations of complex systems built up from measurements of highly varying quality. They will ultimately allow a more honest and rational use of our numerical resources.

IV. USES OF THE GENERAL ACCOUNTING SYSTEM

The accounting framework described above allows the collection of ecosystem data in a form that is useful for a number of analytical purposes. Some of these uses and their importance are described in detail in a number of recent works (e.g.) Hannon and Joiris, 1989, Costanza and Hannon, 1989, Fasham, 1984; Hannon, 1986; MacDonald, 1983; Ulanowicz, 1986; and Wulff *et al.*, 1989). Below we briefly describe some examples that demonstrate existing and potential uses, but the presence of a consistent accounting framework should allow an explosion of new uses not mentioned or conceived of here.

Most uses of the framework require some assumptions about the functional relationships between process inputs and outputs. The examples below are organized in order of increasing sophistication of these assumptions, from static, linear ones, to dynamic, nonlinear ones, to more elaborate optimization functions.

A. Network Analysis

The "mixed units" problem in economics. We have elaborated on the problem of mixed units (multiple products, but most analytical methods require a single unit, or convenient) when only a single unit is available (Hannon, 1989). Therefore is the conversion of mixed units to an equivalent single-product network. There are various ways to calculate intensities (e.g., Hannon, 1989; Costanza, 1985; Costanza, and Hannon, 1989) that allow all the systems to be compared on a common currency (Hannon, Costanza, 1985, Costanza, and Hannon, 1989). The problem has only one non-produced input and there are an equal number of produced outputs. It is possible to calculate the input-output matrix for a non-produced input "embodied in the system" (Costanza and Hannon, 1989). The system can be converted into a single product system ("sunlight") that is amenable to network analysis that can be used for ecological networks is given in a recent paper (Hannon, 1989).

The general accounting framework can be used to connect geographical ecosystems. The first ecosystem would be arranged in a matrix of the second, adjoining the first one, centered on the exchanges between the two ecosystems. The cell in the matrix to the right of the first one would be the net and total output and of net input. The matrix is in length to accommodate those ecosystems. The combined result could then be used to calculate the matrix. For example, the above matrix could reflect the combination of the two ecosystems for the same product from each ecosystem.

B. Simulation

Another possible use for the framework is as the basis for ecological simulation. The dynamic mathematical functions can be used to model the research that would itself benefit from the format like the proposed accounting framework.

ported to the general scientific "bad," and the record of its by the experts familiar with the s needed is a formal, systematic the "degree of goodness" or s can be imagined as composed that is, how little "spread" or how reliable it is as an estimate of all quantitative measurements which we will call the "grades" one is familiar with the grading e, and the process of grading similar to grading students or ng a grade with each number ture of the measure of uncer- u its correct interpretation.

g data of sub-statistical quality ntowitz and Ravitz, 1989). We e, except to say that they are a ng system to allow information ied through the accounting and ms built up from measurements ately allow a more honest and

ACCOUNTING SYSTEM

above allows the collection of a number of analytical purposes. e described in detail in a num- s, 1989, Costanza and Hannon, Donald, 1983; Ulanowicz, 1986; describe some examples that ut the presence of a consistent xplosion of new uses not men-

ne assumptions about the func- ts and outputs. The examples sophistication of these assump- mic, nonlinear ones, to more

A. Network Analysis and Energy Intensities

The "mixed units" problem is a critical one in both ecology and economics. We have elaborated an accounting system which allows multiple products, but most analytical work requires (or at least is much more convenient) when only a single product is tracked. One area of research therefore is the conversion of the full multi-product network into an equivalent single-product network. Work in this area has so far centered on various ways to calculate intensity factors (analogous to prices in economic systems) that allow all the system's products to be converted into a common currency (Hannon, Costanza, and Herendeen, 1986; Hannon and Costanza, 1985, Costanza, and Hannon, 1989). If the system under study has only one non-produced input (usually sunlight for ecological systems) and there are an equal number of products and processes it becomes possible to calculate the input intensities that represent the amount of the non-produced input "embodied" in each of the system's products (Costanza and Hannon, 1989). These energy intensities can then be used to convert the system into a single product network (i.e., in "embodied sunlight") that is amenable to further network analysis. The range of possible network analysis that can be performed on single product ecological networks is given in a recent compendium (Wulff *et al.*, 1989).

The general accounting framework will also allow analysis of different but connected geographical ecosystems. The matrix of use and make for the first ecosystem would be arranged in the manner stated above. The use matrix of the second, adjoining ecosystem would be appended to the use matrix of the first one, centered on the extended diagonal of the first. Any exchanges between the two ecosystems would be placed in the appropriate cell in the matrix to the right of or below the first matrix. The vectors of net and total output and of net input of the first matrix would be extended in length to accommodate those same vectors from the second system. The combined result could then be analyzed as though it were a single system matrix. For example, the above mentioned energy intensities would then reflect the combination of the two systems with possibly differing intensities for the same product from each system.

B. Simulation Modeling

Another possible use for the data in our accounting framework is as the basis for ecological simulation models. These models require that a dynamic mathematical function be used to link process inputs and outputs. The mathematical form of these "production functions" is an active area of research that would itself benefit from the existence of a consistent data format like the proposed accounting system.

Ecosystem simulation has been by far the most popular form of analysis of whole ecosystems. Thousands of ecosystem simulation models have been constructed over the years and we could not begin to summarize them here. Most of these modeling studies involved using either an explicit or an implicit, ad hoc accounting framework as a preliminary step. Our point is that the proposed framework will provide a consistent base from which to start, a minimum ideological baggage, and a minimum of unnecessary duplication of data collection. This step would facilitate ecosystem modeling of all kinds.

C. Optimization

Network analysis, simulation, and the related techniques help to portray the effects of the entire system upon individual processes and product flows. But the data required by energy budgets also can be used to gauge how the system as a whole is acting. In order to quantify such system tendencies some investigators have attempted to formulate objective functions that might portray some overall preferred state or goal. For example, H. T. Odum (1971) placed great stress on the maximization of useful output as a criterion for a population's survival (an idea which has much earlier been articulated by Lotka (1922)). Odum suggested that the Lotka maximum power principle also might be applicable to the ecosystem in aggregate. Hannon (1979) and Hannon *et al.* (1986) have posited that the scalar product of the energy intensities with the net outputs of each system is an appropriate objective function.

Ulanowicz (1980, 1986) defined the network "ascendency" as an appropriate objective function. It is calculated entirely from the exchanges transpiring among the system elements and with their environment. Ascendency is purported to rise monotonically as the system grows and develops.

Fontaine (1981), Loehle (1988), and Herendeen (1991) have attempted to compare the efficacies of the various objective functions as descriptors of system development, but to date, there is no agreement as to the best quantitative description of system development. The problem in comparing the various whole-system hypotheses has been a paucity of data on exchanges and storages in actual ecosystems. The potentially enormous advantages resulting from the discovery of a principle for ecosystem development should highlight the urgent need for more data collected under the accounting schema described above.

Amir (1987) also describes a dynamic optimization process for ecosystems in mathematical detail. He discusses the use of a biowealth function which is said to be maximized by the ecosystem resource allocation solution.

We have elaborated the ne system as an ecological research portion of a large ecosystem strategy. In so doing, they ma ment the future work of other strategy. They may also leave was unimportant to their own and checking it against the re these scientists are assuring th usable to others. In this way, th understanding of the larger ec pieces of the larger system, t scientists will be able to synthe of the larger system function.

The authors thank Craig Loehle and

- AMIR, S. 1979. Economic interpretation of ecosystem models. *Biol. Struct.* 2, 293-314.
- AMIR, S. 1987. Energy pricing, biomass, and the environment. In *Environmental Economics: The Analysis of* pp. 53-103, Leimbreger, Geneva.
- BAIRD, D., AND ULANOWICZ, R. E. 1985. The energy flow in a marine ecosystem, *Ecol. Monographs* 59(4).
- BARBER, M. 1978. A retrospective Mar. 1978, 5, 125-35.
- BULLARD, C., SEBALD, A. 1977. Effects of input-output models, *Rev. Econ. S.*
- COSTANZA, R. 1980. Embodied energy in the United States economy: 1963, 1967, and 1972.
- COSTANZA, R., AND HERENDEEN, R. A. 1984. The energy flow in the United States economy: 1963, 1967, and 1972.
- COSTANZA, R., AND NEILL, C. 1984. The energy flow in the United States economy: 1963, 1967, and 1972.
- COSTANZA, R., NEILL, C., LEIBOWITZ, S. 1983. "Ecological Models of the United States Economy." U.S. Fish and Wildlife Service, Washington, DC FWS/OBS-82/68.

The most popular form of analysis in simulation models have been not begin to summarize them using either an explicit or an preliminary step. Our point is a consistent base from which to and a minimum of unnecessary could facilitate ecosystem modeling

on

ated techniques help to portray al processes and product flows. o can be used to gauge how the quantify such system tendencies formulate objective functions d state or goal. For example, on the maximization of useful vival (an idea which has much Odum suggested that the Lotka applicable to the ecosystem in al. (1986) have posited that the the net outputs of each system

network "ascendency" as an ted entirely from the exchanges and with their environment. cally as the system grows and erendeen (1991) have attempted ctive functions as descriptors of s no agreement as to the best ent. The problem in comparing s been a paucity of data on ms. The potentially enormous of a principle for ecosystem need for more data collected ve.

mic optimization process for scusses the use of a biowealth d by the ecosystem resource

V. SUMMARY AND CONCLUSIONS

We have elaborated the need for and style of a general accounting system as an ecological research tool. When field ecologists investigate one portion of a large ecosystem, they usually design their own research strategy. In so doing, they may miss the opportunity to significantly augment the future work of other scientists by slight changes in their research strategy. They may also leave off some stock or flow measurement which was unimportant to their own needs. By reviewing their research proposal and checking it against the requirements of a general accounting system, these scientists are assuring themselves that their work will be in a form usable to others. In this way, their research is contributing to a network of understanding of the larger ecosystem. As the research progresses on the pieces of the larger system, the time approaches when they and other scientists will be able to synthesize the collective information into a model of the larger system function.

ACKNOWLEDGMENTS

The authors thank Craig Loehle and G. A. Swanson for their useful comments.

REFERENCES

- AMIR, S. 1979. Economic interpretations of equilibrium concepts in ecological systems, *J. Soc. Biol. Struct.* **2**, 293-314.
- AMIR, S. 1987. Energy pricing, biomass accumulation, and project appraisal, in "Environmental Economics: The Analysis of a Major Interface" (G. Pillet and T. Murota, Eds.), pp. 53-103, Leimbreger, Geneva.
- BAIRD, D., AND ULANOWICZ, R. E. 1989. The seasonal dynamics of the Chesapeake Bay ecosystem, *Ecol. Monographs* **59**(4), 329-364.
- BARBER, M. 1978. A retrospective Markovian model for ecosystem resource flow, *Ecol. Model* **5**, 125-35.
- BULLARD, C., SEBALD, A. 1977. Effects of parametric uncertainty and technological change on input-output models, *Rev. Econ. Stat.* **59**(1), 75-81.
- COSTANZA, R. 1980. Embodied energy and economic valuation, *Science* **210**, 1219-1224.
- COSTANZA, R., AND HERENDEEN, R. A. 1984. Embodied energy and economic value in the United States economy: 1963, 1967, and 1972, *Resour. Energy* **6**, 129-164.
- COSTANZA, R., AND NEILL, C. 1984. The energy embodied in the products of ecological systems: a linear programming approach, *J. Theor. Biol.* **106**, 41-57.
- COSTANZA, R., NEILL, C., LEIBOWITZ, S. G., FRUCI, J. R., BAHR, L. M., JR., AND DAY, J. W., JR. 1983. "Ecological Models of the Mississippi Deltaic Plain Region: Data Collection and Presentation." U.S. Fish and Wildlife Service, Division of Biological Services, Washington, DC FWS/OBS-82/68.

- COSTANZA, R., AND HANNON, B. 1989. Dealing with the "mixed" units problem in ecosystem analysis, in "Flow Analysis of Marine Ecosystems" (F. Wulff, J. G. Field, and K. H. Mann, Eds.), Chap. 5, Ecological Studies Series, Springer-Verlag, Berlin/New York.
- DALY, H. 1968. Economics as life science, *J. Polit. Econ.* **76**, 392-401.
- FASHAM, M. J. R. 1984. "Flows of Energy and Materials in Marine Ecosystems," Plenum, New York.
- FINN, J. 1976. Measure of ecosystem structure and function derived from the analysis of flows, *J. Theor. Biol.* **56**, 363-380.
- FONTAINE, T. D. 1981. A self-designing model for testing hypotheses of ecosystem development, in "Progress in Ecological Engineering and Management by Mathematical Modelling" (S. E. Jorgensen, Ed.), pp. 281-291, Elsevier, Amsterdam.
- FUNTOWITZ, S., AND RAVITZ, J. 1989. "Uncertainty and Quality in Science for Policy," Kluwer, Amsterdam.
- HANNON, B. 1973. The structure of ecosystems, *J. Theor. Biol.* **41**, 535-546.
- HANNON, B. 1979. Total energy costs in ecosystems, *J. Theor. Biol.* **80**, 271-93.
- HANNON, B. 1986. Ecosystem control theory, *J. Theor. Biol.* **121**, 417-437.
- HANNON, B. M., AND COSTANZA, R. 1985. Ecosystems with multiple products. *ISEM Journal* **7**, 27-49.
- HANNON, B., COSTANZA, R., AND HERENDEEN, R. 1986. Measures of energy cost and value in ecosystems, *J. Environ. Econ. Management* **13**, 391-401.
- HANNON, B., AND JOIRIS, C. 1989. A seasonal analysis of the Southern North sea ecosystem, *Ecology* **70**(6), 1916-1934.
- HERENDEEN, R. 1991. Do economics-like principles predict ecosystem behavior under changing resource constraints? in "Theoretical Studies Ecosystems: The Network Perspective" (T. Burns and M. Higashi, Eds), pp. 261-287, Cambridge Univ. Press.
- KOOPMANS, T. C. (Ed.) 1951. "Activity Analysis of Production and Allocation," Wiley, New York.
- KUZNETS, S. 1946. Problems of interpretation of national accounts, in "National Income: A Summary of Findings," pp. 111-129, National Bureau of Economic Research, New York.
- LEONTIEFF, W. 1941. "The Structure of the American Economy, 1919-1939," Oxford Univ. Press, New York.
- LINDEMAN, R. L. 1942. The trophic-dynamic aspect of ecology, *Ecology* **23**, 399-418.
- LOEHLE, C. 1988. Evolution: The missing ingredient in systems ecology, *Amer. Nat.* **132**, 884-899.
- LOTKA, A. J. 1922. Contribution to the energetics of evolution, *Proc. Natl. Acad. Sci. U.S.A.* **8**, 147-155.
- MACDONALD, N. 1983. "Trees and Networks in Biological Models," Wiley, Chichester.
- MATIS, J. H., AND PATTEN, B. C. 1981. Environ analysis of linear compartmental systems: The static, time invariant case, *Bull. Int. Stat. Inst.* **48**, 527-565.
- ODUM, E. P. 1953. "Fundamentals of Ecology," Saunders, Philadelphia.
- ODUM, H. T. 1971. "Environment, Power, and Society," Wiley, New York.
- ODUM, E. P., AND ODUM, H. T. 1959. "Fundamentals of Ecology," 2nd ed., Saunders, Philadelphia.
- ODUM, H. T., AND PINKERTON, R. C. 1955. Time's speed regulator: The optimum efficiency for maximum power output in physical and biological systems, *Amer. Sci.* **43**, 331-343.
- PATTERN, B., BOSSERMAN, R., FINN, J. AND CALE, W. 1976. Propagation of cause in ecosystems, in "Systems Analysis and Simulation in Ecology," Vol. 4, pp. 457-479, Academic Press, New York.
- ROUGHGARDEN, J., 1989, The U.S. needs an ecological survey, *Bioscience* **39**(1), 5.
- SCHUBEL, J., *et al.*, in press, Report of the Chesapeake Bay Panel, Marine Environmental Monitoring Committee, National Research Council, Washington, DC.

- STONE, R. 1963. Input-output relations, Vol. 3, pp. 11-22, MIT Press, Cambridge.
- ULANOWICZ, R. E. 1980. An hypothesis, *Biol.* **85**, 223-245.
- ULANOWICZ, R. E. 1986. "Growth and complexity," Springer-Verlag, New York.
- ULANOWICZ, R., AND KEMP, W. 1979. *Ecology* **60**, 871-883.
- WULFF, F., FIELD, J. G., AND MANN, K. H. 1989. "Flow Analysis of Marine Ecosystems," Ecological Studies Series, Springer-Verlag, Berlin.

the "mixed" units problem in ecosystem
 oms" (F. Wulff, J. G. Field, and K. H.
 Springer-Verlag, Berlin/New York.
on. **76**, 392-401.
 erials in Marine Ecosystems," Plenum,
 ction derived from the analysis of flows,
 for testing hypotheses of ecosystem
 ing and Management by Mathematical
 Elsevier, Amsterdam.
 y and Quality in Science for Policy,"
eor. Biol. **41**, 535-546.
J. Theor. Biol. **80**, 271-93.
r. Biol. **121**, 417-437.
 s with multiple products. *ISEM Journal*
 6. Measures of energy cost and value in
 1-401.
 s of the Southern North sea ecosystem,
 dict ecosystem behavior under changing
 ecosystems: The Network Perspective"
 mbridge Univ. Press.
 roduction and Allocation," Wiley, New
 onal accounts, in "National Income: A
 ureau of Economic Research, New York.
 a Economy, 1919-1939," Oxford Univ.
 of ecology, *Ecology* **23**, 399-418.
 t in systems ecology, *Amer. Nat.* **132**,
 evolution, *Proc. Natl. Acad. Sci. U.S.A.*
 ogical Models," Wiley, Chichester.
 is of linear compartmental systems: The
 527-565.
 ders, Philadelphia.
 y," Wiley, New York.
 ntals of Ecology," 2nd ed., Saunders,
 speed regulator: The optimum efficiency
 gical systems, *Amer. Sci.* **43**, 331-343.
 e, W. 1976. Propagation of cause in
 ion in Ecology," Vol. 4, pp. 457-479,
 l survey, *Bioscience* **39**(1), 5.
 Lake Bay Panel, Marine Environmental
 cil, Washington, DC.

- STONE, R. 1963. Input-output relationships 1954-1966, in "A Programme for Growth,"
 Vol. 3, pp. 11-22, MIT Press, Cambridge, MA.
 ULANOWICZ, R. E. 1980. An hypothesis on the development of natural communities. *J. Theor.*
Biol. **85**, 223-245.
 ULANOWICZ, R. E. 1986. "Growth and Development: Ecosystems Phenomenology," Springer-
 Verlag, New York.
 ULANOWICZ, R., AND KEMP, W. 1979. Toward canonical trophic aggregation, *Amer. Nat.* **114**,
 871-883.
 WULFF, F., FIELD, J. G., AND MANN, K. H. 1989, "Network Analysis in Marine Ecology,"
 Ecological Studies Series, Springer-Verlag, Berlin/New York.